



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

November 23, 2015

Jennifer J. Klika
HACCO, Inc.
110 Hopkins Drive
Randolph, WI 53956

Subject: Notification per PRN 98-10 – To add marketing claims to the label
Product Name: Ramik Green AG
EPA Registration Number: 61282-63
Application Date: November 18, 2015
Decision Number: 511256

Dear Ms. Klika:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped “Notification” and will be placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you have any questions, you may contact Gene Benbow at 703-347-0235 or via email at Benbow.Gene@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "M. Lewis". The signature is written in a cursive style with a large initial "M" and a distinct "L".

Marianne Lewis, Acting Product
Manager 7
Invertebrate & Vertebrate Branch 3
Registration Division (7505P)
Office of Pesticide Programs

RESTRICTED USE PESTICIDE
DUE TO HAZARD TO NON-TARGET SPECIES
For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

Ramik® green AG
Fish Flavored, Weather-Resistant Rodenticide
Kills Rats and Mice
For Control of California Ground Squirrels, Plains Pocket Gophers, and Moles*

ACTIVE INGREDIENT:

Diphacinone	
(2-Diphenylacetyl-1,3-Indandione)	0.005%
OTHER INGREDIENTS:	99.995%
TOTAL:	100.000%

EPA Reg. No. 61282-63

EPA Est. No. 61282-WI-01

KEEP OUT OF REACH OF CHILDREN
CAUTION

Net Contents:

- 4 pounds**
- 8 pounds**
- 20 pounds**
- 50 pounds**

Net Weight:

- 4 pounds**
- 8 pounds**
- 20 pounds**
- 50 pounds**

This Product May Not Be Sold in Packaging that Holds Less Than 4 Pounds of Bait.
***Not Approved for this Use Pattern in the following states: California, Hawaii, New Hampshire, and Indiana.**

For use only against the eastern mole (*Scalopus aquaticus*) and the hairy-tailed mole (*Parascalops breweri*) in North Carolina. Not for use against the star-nosed mole (*Condylura cristata parva*) in North Carolina.

Manufactured By:
HACCO, Inc.
110 Hopkins Drive
Randolph, WI 53956

NOTIFICATION

61282-63

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

11/23/2015

FIRST AID	
HAVE LABEL WHEN OBTAINING TREATMENT ADVICE	
If Swallowed:	<ul style="list-style-type: none"> • Call a poison control center, or doctor, or 1-800-498-5743 immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor.
If on Skin or Clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center, or doctor, or 1-800-498-5743 immediately for treatment advice.
If in Eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center, or doctor, or 1-800-498-5743 immediately for treatment advice.
NOTE TO PHYSICIAN	
<p>If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K₁, intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.</p>	
TREATMENT FOR PET POISONING	
<p>If animal eats bait, call veterinarian or 1-800-498-5743 at once.</p>	
NOTE TO VETERINARIAN	
<p>Anticoagulant Diphacinone: For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K₁.</p>	
<p>For 24-hour emergency information on this product, call 1-800-498-5743 (US & Canada) or 1-651-523-0318 (all other areas).</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Keep away from children, domestic animals and pets.

Personal Protective Equipment (PPE):

All handlers (including applicators) must wear long-sleeved shirt, long pants, shoes, socks and water-proof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear waterproof gloves.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying the bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to mammals, birds and other wildlife. Dogs, cats and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not allow bait to be exposed on soil surface. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment wash water or rinsate.

ENDANGERED SPECIES CONSIDERATIONS

Notice: It is a federal offense to use any pesticide in a manner that results in the death of a member of an endangered species. The use of this product may pose a hazard to Federally designated endangered/threatened species.

In California, obtain a copy of the Interim Measures Bulletin for your county from the County Commissioner's office or from the following internet site:

<http://www.cdpr.ca.gov/docs/es/index.htm>

Salt Marsh Harvest Mouse (*Reithrodontomys raviventris*)

Do not use this product in critical habitat with 0.5 miles of salt marsh vegetation and/or brackish water wetlands which are located: 1) near or adjacent to San Pablo Bay and San Francisco Bay, or 2) in the Sacramento River below or adjacent to confluence of the Sacramento River and the San Joaquin River (California).

Morro Bay Kangaroo Rat (*Dipodomys heermanni morroensis*)

Do not use this product in critical habitat within 2.5 miles of Baywood Park which is located on the Morro Bay (California).

Giant Kangaroo Rat (*Dipodomys ingrens*)

Follow the Interim Measures for protecting endangered species in the California Endangered Species Bulletin for the following California counties: Fresno, Kern, Kings, Merced, Monterey, San Benito, San Luis Obispo, Santa Barbara, and Tulare (California).

To protect endangered kangaroo rats, use one of the modified station designs (inverted "T" with "elbows" or platform stations) developed for this purpose in associated with the California Department of Food and Agriculture and the California Department of Fish and Game. A publication is available at the website indicated above.

Tipton Kangaroo Rat (*Dipodomys nitratoides nitratoides*)

Follow the Interim Measures for protecting endangered species in the California Endangered Species Bulletin for the following California counties: Kern, Kings, and Tulare (California).

Fresno Kangaroo Rat (*Dipodomys nitratoides exilis*)

Follow the Interim Measures for protecting endangered species in the California Endangered Species Bulletin for the following California counties: Kern, Kings, Madera, and Merced (California).

Stephens' Kangaroo Rat (*Dipodomys stephens*)

Follow the Interim Measures for protecting endangered species in the California Endangered Species Bulletin for the following California counties: San Bernardino, San Diego, and Riverside (California).

Pygmy Rabbit (*Brachylagus idahoensis*)

Do not use this within one mile of alfalfa fields.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

Use only for the sites, pests, and application methods described on this label.

IMPORTANT: Do not expose children, pets or other non-target animals to rodenticides. To help to prevent accidents:

Note: Bait stations are mandatory for outdoor-above-ground use.

Tamper-resistant bait stations must be used if children, pets, non-target mammals, or birds may access the bait.

USES ASSOCIATED WITH BUILDINGS AND STRUCTURES TO CONTROL RATS AND MICE

IMPORTANT: Do not expose children, pets or other non-target animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger stations are needed in areas open to hooved livestock, raccoons, bears or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled and unconsumed bait as specified on this label.

Note: Bait stations are mandatory for outdoor-above-ground use. Tamper-resistant bait stations must be used if children, pets, non-target mammals, or birds may access the bait.

USE RESTRICTIONS:

In and Around Buildings and Structures

This product may only be used to control Norway rats, roof rats, and house mice in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port or terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure as defined above, is prohibited. This product must not be applied directly to food or feed crops. Do not broadcast bait.

Inside Burrows

This product may also be applied to active rodent burrows to control Norway rats and roof rats *within or beyond* 100 feet of buildings and man-made structures, provided that infestations of these rodents have been confirmed. Because Norway/roof rat infestations may occur in areas farther than 100 feet from buildings and man-made structures when the rodents have ample supplies of food and cover, efforts should be made to remove food trash, garbage, clutter, and debris. Bait must be placed not less than 6 inches into active Norway/roof rat burrows. Do not broadcast bait.

TERRESTRIAL, NONFOOD / NONFEED AREAS:

This product may also be used in the following areas: river banks, irrigation ditches, gullies, railroad tracks, fences, and sanitary landfills. All placements must be in tamper-resistant bait stations or deeply into rat burrows.

Note: This weather-resistant bait may be placed at the above sites, even if wet or damp. Do not place bait in areas where there is a possibility of contaminating food/feed or surfaces that come in direct contact with food/feed. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats or house mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS

RATS: Place 4 to 16 ounces of bait per placement. Space placements at intervals of 15 to 30 feet in infested areas. Maintain an uninterrupted supply of fresh bait for at least 10 days or until there no longer are signs of new feeding by rats.

MICE: Apply ¼ to ½ ounces (1-2 level tablespoons) of bait at each placement location. Space placements at 8- to 12-foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high house mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until there are no longer signs of new feeding by house mice.

FOLLOW-UP: Replace contaminated or spoiled bait immediately. Using waterproof gloves collect and dispose of all dead, exposed animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

CALIFORNIA GROUND SQUIRRELS

USE RESTRICTIONS:

This product may be used to control California ground squirrels (*Spermophilus beecheyi*) in parks, golf courses, fruit tree orchards (dormant season only), non-crop rights-of-way, other non-crop areas (for manual or secured bait station applications) and, in secured bait-station applications only, in or near levee or ditch banks, around farm buildings, along fence lines, in orchards, and in or near crops. **DO NOT** apply this bait directly to food or feed crops or use it in any manner other than prescribed by this label. Do not use this product where nontarget species are likely to be adversely affected or where threatened or endangered species potentially at risk are present. Do not apply bait over roads, walkways or water. Do not allow young children, pets, domesticated animals or personal not associated with the bait application to be in areas where the bait is being applied. Wear waterproof gloves when applying bait. With detergent and hot water, wash all implements used for applying bait. Do not use these implements for mixing, holding or transferring foods or feeds.

APPLICATION DIRECTIONS:

Bait Stations:

Secure bait stations at locations where they will be readily accessible to target California ground squirrels, preferably near active burrows. Secure stations 20 to 100 feet apart, depending upon the density of the California ground squirrel population. Use 2 to 4 pounds of bait per station.

Check bait stations frequently to insure that an uninterrupted supply of bait is always available. Maintain an uninterrupted supply of fresh bait for at least 30 days or until all signs of feeding have stopped. Wearing waterproof gloves collect and dispose of all dead, exposed animals and leftover bait properly.

Bait stations may be maintained in areas on permanent basis, but must only be loaded with toxic bait when California ground squirrels are active and accepting grain-based foods in numbers sufficient to promote effective control

Manual Baiting:

Manually scatter ¼ cup of bait (40 grams) near each burrow opening being used by California ground squirrels. (See more information in *Endangered Species Considerations*). Bait spoons, measuring cups or other suitable implements may be used to measure and distribute the bait. The area around each burrow over which bait is distributed may be no larger than 50 square feet (e.g. a circular area no more than 8 feet in diameter). Re-apply bait to the same area at intervals of 2 to 3 days for a total of 3 to 4 applications, if ground squirrels remain active and continue to consume bait. A continuous supply of bait is needed for 6 to 12 days to ensure that all squirrels present have opportunity to feed on bait for at least 5 consecutive days. Do not exceed the maximum application rate of ¼ cup per burrow per application.

Surveillance and Follow-up:

Dead ground squirrels should begin to appear in or near treated areas within 4 to 5 days after the first bait application. Visit the application site to monitor the effects of treatment 4 days after the first application and at subsequent intervals of 1 or 2 days. Wearing gloves, collect and properly dispose of visible carcasses if ground squirrels and other species found in or near treated areas. Dead animals buried on site must be buried a minimum of 18 inches below the ground surface, preferably deeper. Continue to collect and dispose of dead ground squirrels and search for non-target animals for at least 2 weeks after the last bait application, or longer if carcasses are still being found at that time. Carcass collection and burial should be performed in the late afternoon, near sundown, to reduce the potential for nocturnal scavengers or predators to find carcasses on the ground surface. Wear gloves and use leak-proof plastic bags or other suitable containers for transporting carcasses not buried on site. Where there is a continuous source of reinfestation, establish permanent bait stations and replenish bait as needed.

PLAINS POCKET GOPHERS

USE RESTRICTIONS: For control of plains pocket gophers (*Geomys bursarius*) on lawns, golf courses, forest areas, parks, nurseries, Rangelands, grain fields, forage crops, hay crops, timothy crops, alfalfa crops, vineyards, grape vineyards, in pome fruit (apple, pear), stone fruit (peach, cherry, apricot, plum, prune, nectarine), citrus fruit, avocado and nut tree orchards (almonds, walnuts, pistachios, pecans, filberts, and macadamia), tree plantations, and vegetable crops. Bait must be applied directly into pocket gopher's underground burrow systems. Do not apply bait above ground.

HAND BAITING: Follow the directions indicated below to locate and treat pocket gophers' burrow systems.

Locating Underground Runways: Use a specially designed gopher probe, a metal rod, a strong smooth stick, or other suitable implement to probe the ground 10 to 15 inches away from fan-shaped gopher mounds. Begin probing on the flat side of the fan. When the main runway has been entered, the probe will drop about 2 inches due to decreased resistance from soil.

Applying Bait to Runways: Using a long-handled spoon or mechanical probe, drop 4 ounces of bait into the main runway at each baited point. Cover the opening with sod, a rock, or soil to exclude light. Take care not to cover bait with soil. Depending upon mound density, make two to three bait placements per burrow system. Recover and bury all spilled bait. Applied in this manner, one pound of bait will treat one to eight acres, depending upon the number of pocket gopher burrow systems present.

MOLES

not approved for the control of Moles in the following states:

California, Hawaii, Indiana, and New Hampshire

For use only against the eastern mole (Scalopus aquaticus) and the hairy-tailed mole (Parascalops breweri) in North Carolina. Not for use against the star-nosed mole (Condylura cristata parva) in North Carolina.

USE RESTRICTIONS: For control of moles in lawns, parks, and golf courses. This product may only be used to control moles in manual below ground applications. Do not apply in gardens where food or feed may become contaminated. Do not apply this product by use of an artificial burrow builder. Any person who retrieves carcasses or unused bait following application of this product must wear waterproof gloves.

Specific Directions for Moles: The presence of moles may be indicated by a network of surface ridges in the turf, by a series of conical mounds of earth pushed up from deep burrows, or by damage to flower bulbs.

Surface Ridges: Prior to treatment determine which burrows are active according to one of the following two methods: Either press down a small section of the tunnel or remove a one inch section of the tunnel's roof. Mark these sections and recheck in 24 hours. The burrow is considered active if the flattened runway has been raised up or if the roof has been repaired. Only treat active burrows. Carefully, with a long-handled spoon, punch a hole in the top of active tunnels, drop a teaspoon amount of this product and carefully close the hole with sod or stone. Do not collapse the runway again or allow loose soil to cover the bait. After a few days check to see if the burrows are still active according to one of the two methods described above. If so, repeat treatment.

Conical Mounds: With a long handled spoon carefully remove the center plug from each conical mound. Drop 4 ounces of bait into the underground tunnel. Close the opening with a rock or sod. Do not allow loose dirt to cover bait. If mole activity persists after one treatment, a second treatment may be made. If mole activity persists after two treatments, abandon use of this product and seek another method for controlling moles.

Flower Bulb Protection (Fall Planted): Pack about one-quarter inch of soil against each flower bulb. Place several pellets at sides of and around the flower bulb. Then cover.

FRUIT TREE ORCHARDS (BEARING AND NON-BEARING) FOR PINE AND MEADOW VOLE CONTROL

USE RESTRICTIONS: For control of pine and meadow voles in pome fruit (apples, pears) and stone fruit (peaches, nectarines) tree orchards in the states listed below. Aerial application is only permitted in those following states where aerial is listed next to the state name.

Connecticut	Michigan/aerial
Georgia	Missouri
Idaho/aerial	Montana/aerial
Massachusetts	North Carolina
New Hampshire	South Carolina
Ohio/aerial	Utah/aerial
Oregon/aerial	Virginia/aerial
Pennsylvania/aerial	Vermont/aerial
Washington/aerial	West Virginia/aerial

Timing: Apply after fall harvest (including drops), before new spring growth, and during which three days of rain-free and snow-free weather are expected.

Grazing: Do not graze animals in treated areas.

Hay: Do not use hay cut after application for feed or bedding.

Bare Ground and Unspecified Sites: Do not apply over bare ground or to orchards other than apple, pear, peach or nectarine.

Application Type: Apply only by hand spot baiting and ground broadcasting and aerial broadcast (for those states indicated).

PRE-APPLICATION DIRECTIONS:

Prior to application, examine orchard floor to locate trails and runways systems to be treated.

APPLICATION DIRECTIONS:

Hand Spot Baiting: Place about 1-1/2 oz. of bait in each active hole, trail, or runway at each tree site (10 lbs/acre). When practical, cover bait artificially (e.g., mats, boards) or by pulling overhanging grass over bait.

Ground & Aerial Broadcasting: For aerial application only in those states specified above. For ground application use a commercially-made seed or fertilizer spreader. Determine the vole species that is present in the orchard. Uniformly apply 20 lb of bait per acre rate for Pine Voles and 10 lb of bait per acre rate for Meadow Voles. To avoid exposing nontarget organisms, clean up any bait in piles and return bait to application equipment or dispose of as specified under "Pesticide Disposal."

POST-APPLICATION INSTRUCTIONS:

If populations are high, make a second application 1-2 months after the first application.

NUT ORCHARDS (PECANS, ALMONDS, WALNUTS, FILBERTS/HAZELNUTS) FOR PINE AND MEADOW VOLE CONTROL

USE RESTRICTIONS: For control of pine and meadow voles in Nut Orchards in the following states:

Idaho	Ohio	Montana
Oregon	Washington	

Timing: Apply after fall harvest (including drops), before new spring growth, and when three, rain-free days are expected. For other USE RESTRICTIONS, PRE-APPLICATION, APPLICATION, AND POST-APPLICATION DIRECTIONS, see use directions for pine and meadow voles in fruit orchards above.

CHRISTMAS TREE FARMS, COMMERCIAL NURSERIES, AND TREE PLANTATIONS FOR PINE AND MEADOW VOLE CONTROL

USE RESTRICTIONS: For control of pine and meadow voles in Christmas Tree Farms, Commercial Nurseries, and Tree Plantations in the following states:

Georgia	Michigan	Missouri
North Carolina	South Carolina	Washington

Timing: Apply when three, rain-free days are expected. For other USE RESTRICTIONS, PRE-APPLICATION, APPLICATION, AND POST-APPLICATION DIRECTIONS, see use directions for pine and meadow voles in fruit orchards above.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

Pesticide Disposal: Dispose of wastes resulting from the use of this product in trash or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Plastic Containers: Triple rinse (or equivalent) then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or incineration.

Paper Containers: Dispose of empty container by placing in trash, at an approved waste disposal facility or by incineration.

DISCLAIMER

DISCLAIMER: To the extent permitted by applicable law, seller makes no representation or warranty, either expressed or implied, for results due to misuse, improper handling or improper storage of this material. Nor does Seller assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or improper storage of this material.

OR

LIMITED WARRANTY

To the extent permitted by applicable law, Neogen Corporation makes no warranty concerning uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label. To the extent permitted by applicable law, Neogen Corporation shall not be liable for (1) any consequential, incidental or special damages related in any way to this product or its uses, or (2) any damages related in any way to resistance to pesticides.

OR

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

To the extent permitted by applicable law Manufacturer and Seller warrant that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law this warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Manufacturer, and Buyer and User assume the risk of any such use. To the extent permitted by applicable law, MANUFACTURER AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent permitted by applicable law, in no event shall Manufacturer or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT PERMITTED BY APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF THE MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

Manufacturer and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitations of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Manufacturer.

For non-emergency (e.g., current product information),
Call: **1-800-621-8829**

*Not Approved for this Use Pattern in the following states: California, Hawaii, New Hampshire, and Indiana.

For use only against the eastern mole (*Scalopus aquaticus*) and the hairy-tailed mole (*Parascalops breweri*) in North Carolina. Not for use against the star-nosed mole (*Condylura cristata parva*) in North Carolina.

Code (RR.RL.) RR.RL.

Ramik® is a Registered Trademark of HACCO, Inc.

Optional Marketing Statements: Made in the U.S.A.
<American Flag Graphic>

California Ground Squirrel Graphic



Mouse Silhouette Graphic

Mole Silhouette Graphic

California Ground Squirrel Silhouette Graphic

Gopher Silhouette Graphic

Weather-Resistant Bait

Palatable Bait

Palatable Pellets

Weather-Resistant, Palatable Bait

Easy Application

Contains Diphacinone

Helps Stop Damage Caused by California Ground Squirrels

Helps Stop Damage Caused by Commensal Rodents

Helps Stop Damage Caused by Rats and Mice

Helps Stop Damage Caused by Voles

Helps Stop Damage Caused by Moles

Helps Stop Damage Caused by Pocket Gophers

