

Reg # 61451-1

PM-22

TC 1051

19 11
7



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

APR 15 1994

Robert M. Sielaty
COMPLIANCE SERVICES INTERNATIONAL
AGENT FOR: VETERANS ILEX INC
2001 JEFFERSON DAVIS HWY, #1010
ARLINGTON, VA 22202

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Subject: Label Amendment Submission of 10/07/93 in Response to PR Notice 93-7
EPA Reg. No. 61451-1
TCIPN

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.



Recycled/Recyclable
Printed with Soy/Canola Ink on paper that
contains at least 50% recycled fiber

29 11


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

39 11
7

AGENT FOR: VETERANS ILEX INC.

61451-1 10/07/93

TCIPN

Original Submission

One or more of the statements in the Non-Agricultural Use Requirements box is not found on your original label. Delete the crossed-out sentence(s). If you wish to retain the sentence(s) you must submit an amendment request to the Product Manager. If there are no remaining requirements in the box after you delete the sentence(s), delete the entire Non-Agricultural Use Requirements box. Please refer to the instructions starting on page 45 of Supplement Three to PR Notice 93-7 (Main Labeling Guidance).

Correct the typographical errors circled on your proposed label.

(REVISED PROPOSED LABEL)

4911

(Highlighted items show ADDITIONS to comply with Worker Protection Standards)

TCIPN

ACCEPTED
with COMMENTS
In EPA Letter Dated

APR 15 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
61451-1

ACTIVE INGREDIENT:

Chlorothalonil (Tetrachloroisophthalonitrile) 40.4%

INERT INGREDIENTS: 59.6%

TOTAL 100.0%

Contains 4.2 Pounds Chlorothalonil Per Gallon (500 grams per liter)

KEEP OUT OF REACH OF CHILDREN

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water. Get medical attention.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferable mouth-to-mouth. Get medical attention.

IF ON SKIN: Wash with soap and water.

IF SWALLOWED: Get medical attention.

NOTE TO PHYSICIAN: Persons having an allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids.

See additional Precautionary Statements.

Prior to Use, TCIPN containers should be inverted several times to assure complete mixture of ingredients.

VETERANS-ILEX

Veterans-ILEX, Inc.
320 108th Ave. NE
Bellevue, WA 98004
EPA REG. NO. 61451-1

NET CONTENTS 2 1/2 Gallons

BEST AVAILABLE COPY

59 11

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS WARNING

Avoid contact with skin or clothing. Harmful if swallowed. Fatal if inhaled. Avoid breathing spray mist. Do not get in eyes. In case of contact with eyes, flush with plenty of water immediately for at least 15 minutes. Note: TCPN may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician. Affected persons respond to treatment with antihistamines or steroid creams and/or systemic steroids.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

• Coveralls over short-sleeved shirt and short pants;

• Waterproof gloves;

• Chemical-resistant footwear plus socks;

• Protective eyewear;

• Chemical-resistant headgear for overhead exposure;

• Protective equipment when cleaning equipment, mixing, or loading;

• For mixing and application: a respirator with either an organic vapor cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-22C) or a cartridge approved for pesticides (MSHA/NIOSH approval number prefix TO-14D);

• For application without a dust/fume filtering respirator (MSHA/NIOSH approval number prefix TC-21C);

• Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d) (4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

• Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet;

• Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing;

• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. Do not apply directly to water to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas.

GENERAL USE DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply this product only as specified on this label. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific requirements for labeling and packaging in addition to those contained in other Federal pesticide laws. This Standard also covers protective equipment (PPE) and PPE requirements. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not use or store material on any area treated area during the restricted entry interval (REI) of 48 hours.

PPE required for entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls over short-sleeved shirt and short pants; Waterproof gloves; Chemical-resistant footwear plus socks; Protective eyewear; Chemical-resistant headgear for overhead exposure.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

• Keep children and pets out of the treated area until sprays have dried.

BEST AVAILABLE COPY

6711

TCIPN is a broad spectrum fungicide especially formulated for use to control plant diseases of agricultural crops as well as turf and ornamental plantings. **TCIPN** readily mixes with water to form an easily applied suspension which effectively wets a wide variety of plant surfaces. An essential factor in **TCIPN**'s performance is through end uniform coverage (see the table below for spray volume recommendations).

Prior to use, **TCIPN** containers should be inverted several times to assure complete mixture of ingredients. When adding **TCIPN** to the dilute spray solution, pour the required amount slowly into the tank as water is being added. For concentrate sprays, pre-mix the recommended amount of **TCIPN** with water in a clean container and add to the spray tank as it is being filled. To assure uniform mixture agitate the spray solution while mixing and during spraying.

Do not mix **TCIPN** with agricultural chemicals or fertilizers not specifically recommended on this label unless you have experience which has shown the mixture to be physically compatible, efficacious and non-phytotoxic (non-injurious).

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

SPRAY VOLUME PER ACRE FOR USE WITH TCIPN (gals)*

	Dilute	Concentrate	Aerial
Field Crops / Vegetables	100	5 to 20	5 to 10
Orchard Crops	200 to 400	20 to 150	10 to 20
Christmas Trees	100	20 to 50	
Forest Trees	NA**	NA	10 to 20
Nursery Beds	100	20 to 50	10 to 20

*General recommendations to be followed unless specifically directed otherwise on this label.

**NA - Not applicable.

CROP	DISEASES	RATE/ACRE	USE INSTRUCTIONS
Bean (Snap)	Rust	2 to 4 1/2 pts.	Initiate treatment during early bloom or when disease threatens. Treat on a weekly basis or as necessary to control disease. DO NOT apply within 7 days of harvest. DO NOT feed treated plant parts to livestock or graze treated areas.
	Botrytis Blight (gray mold)	4 1/2 pts.	
Beans, (Dry, Navy, Pinto, Kidney, Lima, Blackeye)	Rust, Anthracnose, Downy Mildew	2 1/2 pts.	Initiate treatment during early bloom and retreat on 7 to 10 day intervals. Use only on dry harvested beans where pods are removed. DO NOT apply within 6 weeks prior to harvest. DO NOT feed treated plant parts to livestock or graze treated areas.
	Cercospora Leaf Spot (Blackeye only)		
Cabbage, Cauliflower, Broccoli, Brussels Sprout	Alternaria Leaf Spot, Downy Mildew	2 1/2 pts.	Initiate treatment after transplanting or soon after emergence of direct seeded crops or as environmental conditions favor disease development. Retreat on 7 to 10 day intervals or as necessary to control disease.
	Ring Spot (California only)	2 1/2 pts.	Initiate treatment of Brussels Sprout at early sprout development or as environmental conditions favor disease development. Retreat on 7 to 10 day intervals or as necessary to control disease.
Carrot	Early Blight (Cercospora)	2 1/2 to 2 3/4 pts.	Initiate treatment as conditions favor disease development. Retreat on 7 to 10 day intervals or as necessary to control disease. TCIPN may be applied through center pivot irrigation equipment as directed below.
	Late Blight (Alternaria)		
Celery	Early Blight (Cercospora)	1 1/2 to 2 1/4 pts.	Initiate treatment when transplants are field set. Apply the lower rate (1 1/2 to 2 1/4 pts.) on a 3 to 5 day interval, the higher rate (3 to 4 1/2 pts.) on a 7 day interval. DO NOT treat within 7 days of harvest. TCIPN may be applied through center pivot irrigation equipment as directed below.
	Late Blight (Alternaria)		
	Basal Stem Rot (<i>Rhizoctonia solani</i>)	3 to 4 1/2 pts.	
	Pink Rot (Suppression)	4 1/2 pts.	
	Early Blight (Cercospora)	2 1/2 to 2 3/4 pts. per 100 gal.	SEEDBEDS - Treat twice a week or as necessary to control disease. Use sufficient water to assure thorough spray coverage. Apply the higher rate where disease pressure is severe.
	Late Blight (Alternaria)		

BEST AVAILABLE COPY

CROP	DISEASES	RATE / ACRE	USE INSTRUCTIONS
Corn (Sweet) for fresh market and Field grown for seed	Helminthosporium Leaf Blights Rust	1½ to 2½ pts.	Initiate treatment as conditions favor disease development. Retreat on 4 to 7 day intervals or as necessary to control diseases. Use the higher rate where disease pressure is severe. DO NOT treat sweet corn to be processed. DO NOT ensile, use as livestock feed or allow livestock to graze treated areas. DO NOT treat within 14 days of harvest.
Cucumber	Anthracnose Downy Mildew Target Spot (Florida only)	2½ to 2½ pts.	Initiate treatment when plants reach first true leaf or as conditions favor disease development. Retreat on 4 to 7 day intervals. Employ short interval and high rate for severe disease pressure. TCIPN may be applied through center pivot irrigation equipment as directed below.
	Gummy Stem Blight Leaf Blight Powdery Mildew (except South western states) Scab	2½ to 4½ pts.	
	Fruit Rot (<i>Rhizoctonia solani</i>) (suppression only)	12 pts.	Use sufficient water to assure runoff to soil surface. Apply once only when vines begin to form. TCIPN may be applied through center pivot irrigation equipment as directed below.
Cantaloupe, Honeydew Melon, Muskmelon, Pumpkin, Squash, Watermelon	Anthracnose Downy Mildew	1 to 2½ pts.	Initiate treatment when plants reach first true leaf or as conditions favor disease development.
	Cercospora Leaf Spot Gummy Stem Leaf Blight Powdery Mildew (except south western states) Scab	2½ to 4½ pts.	Retreat on 7 day intervals. Employ high rate for severe disease pressure. TCIPN may be applied through center pivot irrigation equipment as directed below. <ul style="list-style-type: none"> • DO NOT tank mix TCIPN with any other product on watermelons. This includes all crop protection products, foliar fertilizers, surfactants, spreaders, stickers, and oils. • DO NOT apply TCIPN to watermelon fruit within 21 days of harvest if periods of intense sunlight and high temperatures are expected. • DO NOT apply TCIPN to watermelon fruit within 21 days of harvest when vines do not provide proper shading of fruit or if vines are stressed due to drought.
Grasses (grown for seed)	Leaf Rust Selenophoma (eyespot) Stem Rust Stripe Rust	1½ to 2 pts.	Initiate treatment during stem elongation as conditions favor disease development. Retreat at flag leaf and head emergence. Where disease pressure is severe use the high rate and retreat on 14 day intervals. DO NOT feed or allow livestock to graze treated plants. DO NOT treat within 14 days of harvest.
Mint	Rust, Septoria Leaf Spot	2 pts.	Initiate treatment when plants are 4 to 8 inches tall. Retreat on 7 to 10 day intervals or as necessary to control disease. DO NOT treat more than three times per season or within 80 days of harvest. DO NOT feed treated plant parts, fresh or extracted, to livestock. TCIPN is restricted to use on mint in Indiana, Michigan and Wisconsin only.
Onion (dry bulb)	Botrytis Leaf Blight (Blast) Purple Blotch	2 to 3 pts.	Initiate treatment when first disease symptoms appear or when conditions favor disease development.
Onion (green bunching)	Botrytis Leaf Blight (Blast)	2 to 4 pts.	Retreat on 7 to 10 day intervals as necessary to control disease. During periods of heavy dew or continued rain use the shorter interval and higher rate. DO NOT treat within 7 days prior to harvest of onions (dry bulb) or garlic. DO NOT treat onions (green bunching), leeks or shallots more than 3 times per season or within 14 days of harvest. Where additional disease control is needed prior to harvest use another registered fungicide.
Onion (grown for seed) Garlic Leek Shallot	Downy Mildew (suppression) Purple Blotch	2 to 4 pts.	
Papaya	Alternaria Fruit Spot Stem End Rot Anthracnose	3 to 5½ pts.	Initiate treatment when conditions favor disease development. Thorough coverage of fruit and foliage is essential for control. Retreat on 14 day interval so long as conditions support disease development. Use the higher rate when disease pressure is severe. DO NOT feed treated plant parts, fresh or processed, to livestock. Treat with ground sprayers only.
Parasip	Alternaria Leaf Spot Anthracnose Botrytis Blight (Gray Mold) Bottom Rot (<i>Rhizoctonia</i>) Downy Mildew	2 to 3 pts.	Initiate treatment when conditions favor disease development. Retreat on a 7 to 10 day interval while conditions favor disease development. Use the high rate and short interval when disease pressure is severe. DO NOT treat more than 4 times per season or within 10 days prior to harvest. DO NOT feed treated plant parts to livestock.
Passion Fruit (Hawaii only)	Alternaria Fruit and Leaf Spot (Passion Fruit Brown Spot)	2½ pts.	Initiate treatment when conditions favor disease development (April to July) or when spots appear on fruit. Thorough coverage of fruit and foliage is essential for control. Retreat on 14 day intervals as needed for disease control. DO NOT feed treated plant parts to livestock grown for food.

8911

CROP	DISEASES	RATE / ACRE	USE INSTRUCTIONS
Peanut	Cercospora Leaf Spot (Early) Cercosporodinium Leaf Spot (Late)	1 1/2 to 2 1/2 pts.	Initiate treatment when conditions (leaf wetness) favor disease development often occurring 30 to 40 days after planting. Retreat on a 10 to 14 day interval. When conditions favor moderate to severe disease development, Late Leafspot, Rust or Web Blotch use the high rate and short interval. TCIPN at the high rate may be applied through center pivot irrigation equipment as directed below. DO NOT treat within 14 days of harvest. DO NOT feed treated plant parts to livestock.
	Rust Web Blotch	2 1/2 pts.	
Potato	Botrytis Vine Rot Early Blight Late Blight	1 1/2 to 2 1/2 pts.	Initiate treatment when plants are 6 to 8 inches tall or conditions favor disease development. Retreat on 7 to 10 day intervals or as necessary for disease control. When conditions favor severe disease development use the high rate and short interval. TCIPN may be applied through center pivot irrigation equipment as directed below. DO NOT exceed 10 day intervals between treatment when applying through center pivot.
	(Dryland Production) Early Blight Late Blight	1 1/2 to 2 1/2 pts.	
Soybean	Anthrachnose, Diaporthe Pod & Stem Blight, Frogeye Leaf Spot (Cercospora Soijna), Purple Seed Stain, Cercospora Leaf Blight (Cercospora Kikuchii), Septoria Brown Spot	1 1/2 to 2 1/2 pts. or 2 to 3 1/2 pts.	Where two treatments are planned apply 2 to 3 1/2 pts. per application. Where three treatments are planned apply 1 1/2 to 2 1/2 pts. per application. Three treatments are recommended for areas with a history of moderate to severe disease incidence. TCIPN may be applied through center pivot irrigation equipment as directed below. DETERMINATE SOYBEANS (Southern): TWO TREATMENTS 1st during early pod set (R3 stage, pods 1/8 to 3/8 inch long); 2nd at beginning of seed formation (R5 stage) about 14 days after 1st treatment. THREE TREATMENTS 1st at initiation of flowering (R1 stage); 2nd during early pod set (R3 stage); and 3rd at beginning of seed formation (R5 stage). INDETERMINATE SOYBEANS (Northern): TWO TREATMENTS 1st when largest pods are 1 to 1 1/2 inches long; 2nd 14 days later. THREE TREATMENTS - 1st 7 days after first bloom, 2nd and 3rd on a 14 day interval thereafter. TCIPN may be tank mixed with Benlate*50 WP for use on indeterminate soybeans. Treat with 1 1/2 pts. of TCIPN plus 8 oz. of Benlate 50 WP. Make 1st treatment when top pods are 1/2 to 1 inch long with the 2nd treatment 14 days later. DO NOT feed soybean hay or harvest trash or allow livestock to graze treated areas.
			*Benlate is a registered trademark of E.I. DuPont de Nemours and Company, Inc.
Tomato	Early Blight, Late Blight, Gray Leaf Spot, Gray Leaf Mold Septoria Leaf Spot	2 to 3 pts. (foliage)	Initiate treatment when conditions such as dew or rain favor disease development. For foliage, retreat on 7 to 10 day intervals; for fruit, retreat on 7 to 14 day intervals. When conditions favor severe disease incidence employ the high rates and short intervals. TCIPN may be tank mixed with KOCIDE 101, KOCIDE 606 or other copper base products registered for control of bacterial diseases. Review all label instructions and limitations before tank mixing. DO NOT tank mix with Copper - Count-N for concentrate sprays. TCIPN may be applied through center pivot irrigation equipment as directed below.
	Anthrachnose Alternaria Fruit Rot (Black Mold) Botrytis Gray Mold Late Blight Fruit Rot	3 to 4 pts. (fruit)	

BEST AVAILABLE COPY

USE AREA	DISEASES	RATE OUNCES /1000 sq. ft.	USE INSTRUCTIONS
Turfgrasses	Gray Snow Mold (Typhula spp)	8 to 16	Initiate treatment prior to snow cover in the fall. Apply the higher rate if turf remains frozen prior to snow cover. Where snow cover is intermittent or not present retreat with the low rate on a monthly basis so long as disease conditions persist.
	Pink Snow Mold (Gerlachia or Fusarium Patch)	8	
			For Pink Mold tank mix with Tersan* 1991 50 *WP at 2 ounces or Chipco** 26019 50 WP at 4 ounces per 1000 square feet. *Tersan is a registered trademark of E.I. Du Pont deNemours & Company, Inc. **Chipco is a registered trademark of Rhone - Poulenc, Inc.

9911

CROP	DISEASES	RATE PER ACRE	RATE PER 100 GAL ¹	USE INSTRUCTIONS
Apricot, Cherry, *Nectarine, Peach, Plum, Prune	Coryneum Blight (Shothole) Leaf Curl	4½ - 6 pts.	1½ - 2 pts.	Initiate treatment in the late autumn at leaf fall. When conditions favor severe disease development use the high rate applied one to two times in late winter prior to bud swell. Where leaf fall treatment cannot be made for control of leaf curl, a spring treatment prior to bud swell may be made. For control of Coryneum Blight (Shothole) also treat at budbreak and shuck split.
	Blossom Blight Brown Rot	4½ - 8 pts.	1½ - 2 pts.	Apply the high rate for trees taller than 20 ft. (6 to 8 pts. over 20 ft. and 4½ to 6 pts. under 20 ft.). Treat first at popcorn stage followed by a second treatment at full bloom. When conditions favor severe disease incidence make a third treatment at petal fall.
	Apricot Scab Cherry Leafspot Nectarine Scab Peach Scab	4½ - 6 pts.	1½ - 2 pts.	Treat first at full bloom; second treatment at petal fall; and third treatment at shuck split. To control cherry leafspot after harvest, treat foliage within 7 days after fruit removal. A second treatment at a 10 to 14 day interval is desirable where cherry leafspot has been severe. DO NOT apply TCIPN after shuck split and before harvest. *Use in conjunction with dilute spray volume recommendations on this label.
Conifers (Forests, Christmas trees and nursery beds)	Botrytis Seeding Blight Phoma Twig Blight	2 to 4 pts.	2 to 4 pts.	Initiate treatment when in nursery beds (seedlings 4 inches tall) and conditions favor disease development. Retreat on 7 to 14 day intervals as required for disease control. Use high rate and short interval when conditions favor severe disease incidence.
	Lophodermium Needlecast	2 to 4 pts.	2 to 4 pts.	Initiate treatment in North Central and Northeastern states in mid July to early August prior to infection occurring. Retreat on 3 to 4 week intervals so long as conditions favor disease development. Treat nurseries with 4 pts. of TCIPN on a 3 week schedule so long as conditions favor disease development.
	Swiss Needlecast	4 to 8 pts. (single treatment)	4 to 8 pts.	Treat Christmas trees or forests once in the spring when new growth is 1/2 to 2 inches long.
	Sclerotinia Canker (pines) Rhabdocline Needlecast (Douglas Fir) Swiss Needlecast	2 to 4 pts.	2 to 4 pts.	Treat in the spring when new growth is ½ to 2 inches long. Retreat on 3 to 4 week intervals so long as conditions favor disease development. In nursery beds use the high rate retreating at 3 week intervals so long as conditions favor disease development.
	Sirococcus Tip Blight	3 to 5 pts.	3 to 5 pts.	
	Scirrhia Brown Spot (pines) Rhizosphaera Needlecast (Spruces)	8 pts.	8 pts.	
Golf Course Fairways	Helminthosporium Leaf Spot	8 to 14 pts.		Initiate treatment when conditions favor disease development. Retreat on 7 to 21 day intervals depending on use rate and disease incidence. When conditions favor severe disease incidence use the high rate and short interval. Use 30 to 40 gallons of spray per acre Rhizoctonia Brown Patch should be retreated on 7 to 14 day intervals. DO NOT water or mow following treatment until spray deposits dry. Use in conjunction with high standards of turf management.
	Sclerotinia Dollar Spot	4 to 14 pts.		
	Rhizoctonia Brown Patch	8 to 14 pts.		

¹ Volumetric rates to be used only with full dilute spray volume specified on this label for tree and orchard crops.

USE AREA	DISEASES	RATE OUNCES /1000 sq. ft.		TREATMENT INTERVAL (days)	
		Prevent	Cure		
Golf Course Tees	Curvularia Leaf Spot	3 - 6	6 - 11	7 - 10	Initiate treatment when conditions favor disease development. Thorough spray coverage is essential to the performance of TCIPN. Use sufficient spray volume (usually 2 to 10 gallons per 1000 square feet) to insure thorough coverage. Preventative Rate: To prevent infection when disease conditions are light to moderate, use the lower rate and when disease conditions are severe use the high rate and short interval. Curative Rate: To control existing infections, apply the curative rate at a 7 day interval. Use the higher recommended rate under severe disease conditions.
Golf Green	Dollar Spot			7 - 14	
Ornamental	Gray Leaf Spot			7 - 10	
	Helminthosporium Leaf Spot and Melting Out			7 - 10	
Turf	Large Brown Patch			7 - 10	
	Red Thread	3 - 9	9 - 11	7 - 10	
	Copper Spot	6 - 9	9 - 11	7 - 10	
	Steam Rust of Bluegrass			7 - 14	

BEST AVAILABLE COPY

10911
7

USE AREA	DISEASES	RATE PINTS /100 gallons	USE INSTRUCTIONS
Ornamentals (See chart for species)	(See Chart)	2 pts.	Initiate treatment in accordance with the chart below. Retreat on a 7 to 14 day schedule while conditions favor disease development.

SPECIES BULBS & FLOWERING PLANTS	TARGET AND DISEASES	TREATMENT INITIATION
Carnation	Alternaria Leaf Spot / Branch Rot; Botrytis Flower Blight	Transplant of cuttings: Cool, moist conditions
Chrysanthemum, Daisy	Mycosphaerella Ray Blight, Septoria Leaf Spot Botrytis Flower Blight (Gray Mold)	Transplant of cuttings Pre-bloom
Geranium	Botrytis Blight, Rust	Cool, moist conditions
Gladiolus	Curvularia Leaf/Flower Spot, Botrytis Leaf/Flower Spot	Early propagation
Hollyhock	Rust	Early seedling stage
Hydrangea* (foliage only)	Cercospora and Septoria Leaf Spots, Rust	Early propagation
Iris	Botrytis Blossom Blight, Didymella Leaf Spot	Cool, moist conditions
Lily	Botrytis Gray Mold	Pre-bloom
Petunia*	Phytophthora Blight (foliar phase) Botrytis Blight	Pre-bloom
Rose (Use 1 1/2 pt. per 100 gallons)	Black Spot, Botrytis Blight	Spring bud break
Statico	Anthracnose, Cercospora, Alternaria, Botrytis Leaf Blights	Spring bud break
Zinnia	Powdery Mildew	First sign of disease

FOLIAGE PLANTS	TARGET AND DISEASES	TREATMENT INITIATION
Dracaena	Fusarium Leaf Spot	Pre-transplant
Pachysandra (Use 4 pt. per 100 gal)	Volutella Leaf Blight	Spring bud break
Leatherleaf fern	Ascochyta Blight, Cercospora Leaf Spot, Cylindrocadium Leaf Spot, Rhizoctonia Blight	Spring bud break
Parlor Palm (Chamaedorea)	Bipolaris (Helminthosporium) Leaf Spot	Cool, moist conditions
Prayer Plant (Maranta)	Helminthosporium Leaf Spot	Early propagation
Oyster Plant (Rhoeo)	Tan Leaf Spot	Early propagation
Syngonium	Cephalosporium Leaf Spot	Warm, moist conditions
Philodendron	Phytophthora Blight, Dactyaria Leaf Spot	Moist conditions

SHRUBS AND TREES	TARGET AND DISEASES	TREATMENT INITIATION
Ash (Fraxinus)	Cercospora, Cercosporidium, Cylindrosporium Leaf Spots	Spring bud break
Azalea*	Phytophthora die-back;	Near leaf emergence,
Rhododendron*	Ovulinia Flower Blight	Early bloom
Buckeye, Horsechestnut	Leaf Blotch, Anthracnose	Spring bud break
Cherry-Laurel	Cercospora Leaf Spot	Petal fall
Crabapple	Scab, Cedar-apple Rust, Sphaeropsis Leaf Spot	Spring bud break
Dogwood	Septoria Leaf Spot	Early bloom
Euonymus	Anthracnose	Spring bud break
Firethorn	Scab	Spring bud break
Flowering Almond, Quince, Sand Cherry	Monilinia Blossom / Branch Blight	Early bloom
Hawthorn	Rust, Febrées Leaf Spot	Pre-bloom
Holly	Rhizoctonia Web Blight	Warm, moist conditions
Mountain Laurel	Cercospora Leaf Spot	Spring bud break
Oak (red group only)	Taphrina Blister, Actinopelta Leaf Spot, Anthracnose	Before budawell
Oregon-Grape (Mahonia)	Rust	Spring bud break
Photinia	Febrées (Entomosporium) Leaf Spot	Spring bud break
Pieris (Andromeda)	Phytophthora die-back	New leaf emergence
Poplar	Marssonina Leaf Spot	Spring bud break
Privet	Cercospora Leaf Spot	Prolonged wet conditions
Sycamore, Planetree	Anthracnose	Spring bud break
Viburnum	Powdery Mildew	Mid-summer

*Discoloration of blooms has been seen on certain varieties when applications are made during flowering.

11 7 11

APPLICATION AND CALIBRATION TECHNIQUES THROUGH IRRIGATION EQUIPMENT

Apply this product only through center pivot, traveling gun, solid set and wheel mobile irrigation system(s). Do Not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do Not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. 'Public water system' means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days per year.

Controls for both irrigation water and pesticide injection systems must be functionally interlocked, so as to automatically terminate pesticide injection when the irrigation water pump motor stops. A person knowledgeable of the irrigation system and responsible for its operation shall be present so as to discontinue pesticide injection and make necessary adjustments should the need arise.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The system must also contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must be fitted with a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur. Do Not apply when wind speed favors drift beyond the area intended for treatment.

TCIPN may be used through two basic types of sprinkler irrigation systems as outlined in Sections 1 and 2 below. Determine which type of system is in place, then refer to the appropriate directions provided for each type.

I. Center Pivot or Traveling Gun

For injection of pesticides, these continuously moving systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. After determining the volume of water injected during a full operation circle, mix the recommended amount of TCIPN to cover the acreage involved with sufficient water to equal the volume injected during a full operation circle. Inject the TCIPN/water mixture into the system through one continuous circle. Mixture in the chemical supply tank must be continuously agitated during the injection run. Shut off injection device after one revolution or run, but continue to operate irrigation system until TCIPN has been flushed from all sprinkler heads.

II. Solid Set or Wheel Mobile

For injection of pesticides with stationary systems, a positive displacement injection pump (e.g., diaphragm pump) can be used. This pump must be effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Position irrigation equipment and determine area covered during a 30 to 60 minute set. Mix the recommended amount of TCIPN to cover the acreage involved with sufficient water to equal the volume injected during a 30 to 60 minute set. Operate the entire system at normal pressures recommended by the manufacturer of injection equipment used for amount of time established during calibration. No agitation should be required. Inject the TCIPN/water mixture into the irrigation system at the beginning or end of an irrigation set or as a separate application. Close the injection device after treatment is completed and continue to operate irrigation system until TCIPN has been flushed from all sprinkler heads.

III. Precautions Associated with Application of TCIPN

- A. DO NOT operate irrigation system without anti-back siphoning system.
- B. Injection equipment for TCIPN should be fitted to the discharge side of the irrigation pump or other pressurized devices.
- C. Irrigation water containing TCIPN should be contained to the treated area until absorbed by the soil.
- D. Irrigation equipment should always be operated in a manner as specified by its manufacturer.

WARRANTY STATEMENT

VETERANS-ILEX warrants that this product in its unopened package conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions on the plants and crops specified. THERE ARE NO OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING A WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE. This warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller and buyer assumes all risk of any such use.

BEST AVAILABLE COPY

110893