

61842-10

10/24/2012

1/15



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Mr. Ross Gilbert
Tessengerlo Kerley, Inc
C/O Pyxis Regulatory Consulting, Inc
4110 136th St. NW
Gig Harbor, WA 98332

NOTIFICATION

OCT 24 2012

Subject: Notification for Alternate Brand Name per PRN 98-10
Submission date: 9/10/2012
Product Name: FlagStick® Fungicide 70 DF
EPA Reg. No. 61842-10
Decision Number 469976

Dear Registrant:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10. The Registration Division (RD) has conducted a review of this request for applicability under PRN 98-10 and finds that the action(s) requested falls within the scope of PRN-98-10.

The Alternate Brand Name **FlagStick® Fungicide 70 DF** dated 9/10/2012 is "acceptable" and will be placed in the regulatory file.

If you have questions concerning this letter, please call Banza Djapao at 703-305-7269 or you may call me at 703-305-7740.

Sincerely,

A handwritten signature in cursive script that reads "William Cutchin".

William Cutchin
Acting Product Manager 20
Fungicide Branch
Registration Division (7504P)

2/15

Please read instructions on reverse before completing form.

Form Approved. U.S. No. 2070-0060. Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 61842-10	2. EPA Product Manager S. Joyner	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Tessengerlo Kerley, Inc. / FlagStick®	PM# 20	
5. Name and Address of Applicant (Include ZIP Code) Tessengerlo Kerley, Inc. c/o Pyxis Regulatory Consulting, Inc. 4110 136th St. NW Gig Harbor, WA 98332 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of an alternate brand name per PR Notice 98-10. The alternate brand name is "FlagStick® Fungicide 70DF". This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
* Certification must be submitted				<input checked="" type="checkbox"/> Other (Specify) poly lined foil bag	
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input checked="" type="checkbox"/> Container		4. Size(s) Retail Container 5.71 lbs.		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph Paper glued Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Ross Gilbert	Title Agent	Telephone No. (Include Area Code) 253-853-7369
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Agent	
4. Typed Name Ross Gilbert	5. Date 9/10/2012	

PYXIS REGULATORY CONSULTING, INC.

4110 136th St. NW
Gig Harbor, WA 98332

Phone: 253-853-7369
Fax: 253-853-5516
Email: www.PyxisRC.com

3/15
9/12

September 10, 2012

Shaja Joyner (PM 20)
Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 S. Crystal Drive
Arlington, VA 22202-4501

RE: Tessengerlo Kerley, Inc. – FlagStick® (EPA Reg. No. 61842-10)
Notification of an Alternate Brand Name per PRN 98-10

Dear Ms. Joyner,


On behalf of Tessengerlo Kerley, Inc., please find the enclosed notification of the addition of an alternate brand name per PRN 98-10 for FlagStick® (EPA Reg. No. 61842-10). The alternate brand name is "FlagStick® Fungicide 70DF".

In support of this notification, please find the following:

1. Completed Application for Registration (EPA Form 8570-1)
2. One (1) copy of the FlagStick® Fungicide 70DF labeling with changes tracked
3. One (1) copy of the FlagStick® Fungicide 70DF labeling with changes incorporated
4. Certification With Respect to Label Integrity
5. Once (1) copy of the FlagStick® Fungicide 70DF label on CD
6. Letter of Authorization

Please feel free to contact me (telephone 253-853-7369; email Ross@PyxisRC.com) if you have any questions or need any additional information.


Sincerely,



Ross Gilbert

Enclosures

cc: B. Thomassen; Tessengerlo Kerley, Inc.



FlagStick[®] Fungicide 70DF

GROUP 33 FUNGICIDE

ACTIVE INGREDIENT:	
Aluminum tris (O-ethyl phosphonate)	70.0%
OTHER INGREDIENTS:	30.0%
TOTAL:	100.0%

FlagStick[®] Fungicide 70DF contains aluminum tris (O-ethyl phosphonate), the same active ingredient found in Chipco[®] Signature[®].

KEEP OUT OF REACH OF CHILDREN WARNING – AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-877-1737 for emergency medical treatment information.	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

Causes substantial but temporary eye injury. Harmful if swallowed or absorbed through the skin or inhaled. Do not get in eyes or on clothing. Avoid contact with skin.

EPA Reg. No. 61842-10

Manufactured for:
 Tessengerlo Kerley, Inc.
 2255 North 44th Street, Suite 300
 Phoenix, AZ 85008-3279
 1-888-732-8246

EPA Est. No. 

Net Weight:

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistant category selection chart. Applicators and other handlers must wear long-sleeved shirt and long pants, chemical resistance gloves made of any waterproof material such as polyethylene or polyvinyl chloride, shoes plus socks, and protective eyewear.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic and estuarine invertebrates. Drift and runoff may be hazardous to aquatic and estuarine organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label before using this product. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

FlagStick® Fungicide 70DF is for use on ornamentals and bedding plants grown in field nurseries, greenhouses, and landscaping and conifer nurseries, phytophthora root rot prevention for conifers in nurseries, downy mildew control in roses, fire blight suppression in ornamental pear, pyracantha and hawthorn, bacterial blight suppression in English ivy, schefflera, anthurium, dieffenbachia, spathaphyllum, syngonium, and ficus to suppress *Xanthomonas campestris* pathovars *hederae*, *dieffenbachiae*, *syngonii*, and *fici* and on turf (golf courses, sod farms, and industrial or municipal turf areas).

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours.

Exception: If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks, and protective eyewear (such as goggles, face shield or safety mask).

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep unprotected persons out of the treated areas until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container and keep tightly closed. Store in a cool dry place.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CHEMIGATION

APPLICATION DIRECTIONS THROUGH SPRINKLER AND DRIP IRRIGATION SYSTEMS

Apply this product only through sprinkler irrigation systems including microjet, drip, solid set and center pivot. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

SET UP: Clean all tanks, hoses, fittings and other parts of the injector system. Be sure to remove residues of prior applications, salt residues or other foreign substances. Flush using clean water.

APPLICATION INSTRUCTIONS: First mix **FlagStick® Fungicide 70DF** in a tank filled with 1/2 to 3/4 the prescribed amount of water. Begin agitation. Add **FlagStick® Fungicide 70DF** to the tank, and then add the remaining volume of water. Set sprinkler to irrigate 0.1 to 0.3 inch of water per acre. Start sprinkler and inject the concentrated mixture of **FlagStick® Fungicide 70DF** into the irrigation system near the end of the irrigation cycle checking to be sure the proper rate per acre is being applied. The **FlagStick® Fungicide 70DF** mixture should be injected using a positive displacement pump into the main line in front of a right angle turn to achieve proper incorporation. Stop the injection equipment after treatment is completed and continue to operate the irrigation system until all of the **FlagStick® Fungicide 70DF** mixture has been cleared from the last sprinkler head.

If you should have any other questions about calibration, you should contact State Extension Service

specialists, equipment manufacturers or other experts.

NOTE: For Microjet and Drip Irrigation Systems: After treatment with **FlagStick® Fungicide 70DF** has been completed, prevent additional irrigation for 24 to 48 hours.

For Solid Set and Center Pivot Irrigation Systems: After treatment with **FlagStick® Fungicide 70DF** has been completed, prevent additional irrigation until foliage is dry to avoid washing **FlagStick® Fungicide 70DF** off the plants.

SPECIFIC REQUIREMENTS FOR SPRINKLER AND DRIP IRRIGATION SYSTEMS

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension.

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shutdown.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e. g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

SPECIFIC REQUIREMENTS FOR CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with

8/15

pesticides and capable of being fitted with a system interlock.

- 7. Do not apply when wind speed favors drift beyond the area intended for treatment.

ORNAMENTALS AND BEDDING PLANTS

FlagStick® Fungicide 70DF may be applied to ornamentals and bedding plants grown in field nurseries, greenhouses, and landscaping and conifer nurseries for the control of diseases caused by *Pythium* and *Phytophthora*. **FlagStick® Fungicide 70DF** may also be used to control downy mildew and fire blight and for the suppression of bacterial blight caused by certain pathovars of *Xanthomonas campestris*.

For best results application of **FlagStick® Fungicide 70DF** must be made before disease develops. Apply when conditions favor disease development and follow sound IPM and cultural management practices. When diseases pressure is severe, use higher rates. Plant injury may occur if specified rates are exceeded or applications are made at more frequent intervals than listed here.

PLANT TOLERANCE

- Plant tolerances to **FlagStick® Fungicide 70DF** have been found to be acceptable in the specific genera and species listed on this label. There are far too many species and varieties of ornamental plants to determine the plant tolerance of every one to **FlagStick® Fungicide 70DF**. If using **FlagStick® Fungicide 70DF** for the first time or plant tolerance is not known, test for possible phytotoxicity by treating a small area at the specified rate. Wait 3-7 days and inspect for visual damage. If damage is present, do not use.

ORNAMENTAL AND BEDDING PLANT TANK MIXTURES

ALWAYS READ AND FOLLOW THE LABEL DIRECTIONS OF ALL TANK MIX COMPONENTS.

Observe the most restrictive label limitations and precautions and do not exceed the label rates. Do not tank mix with products which prohibit tank mixing.

Additional Disease Control: **FlagStick® Fungicide 70DF** is compatible with most commonly used fungicides and thus may be tank mixed with fungicides labeled for control of additional diseases. For control of diseases such as Aerial Web Blight, Alternaria Leaf Blight, Botrytis Blight, Fusarium Leaf Spot, Helminthosporium Leaf Spot, and Ray Blight, **FlagStick® Fungicide 70DF** can be tank-mixed with other fungicides labeled for control of these diseases.

If a tank-mix with other fungicides is used, follow label directions for the use of that product and apply at the specified rate for control of the target disease organism. **FlagStick® Fungicide 70DF** is not physically compatible with some flowable formulations of pesticides such as chlorothalonil or mancozeb fungicides. These products must NOT be combined for application.

COMPATIBILITY

Do NOT tank-mix **FlagStick® Fungicide 70DF** with copper compounds. Tessengerlo Kerley Inc. has determined that the mixing of copper containing compounds with **FlagStick® Fungicide 70DF** may cause phytotoxicity. In addition, the application of **FlagStick® Fungicide 70DF** to foliage with copper residues may cause phytotoxicity from solubilized copper on the leaf surface. Allow at least a 14 day interval between applications of **FlagStick® Fungicide 70DF** and any copper based products.

FlagStick® Fungicide 70DF may not be compatible with foliar fertilizers.

- Some adjuvants such as stickers, wetting agents and those that enhance pesticide leaf penetration may not be compatible or may cause phytotoxicity when mixed with **FlagStick® Fungicide 70DF**.

To test the mixing compatibility of **FlagStick® Fungicide 70DF** with any tank mix partner, conduct a "jar test" by mixing a small amount of the tank mix components in a small container. Mix the ingredients with water in the concentration, and in the order of addition desired for the final spray application. Wait 15 minutes. Check for any indication of incompatibility such as separation or curdling. If any indication of incompatibility exists, do not use this application mixture.

ORNAMENTALS

FOLIAR APPLICATIONS to Aglaonema, Aphelandra, Azalea, Bougainvillea, Boxwood, *Cattelya skinneri*, Cissus, Dieffenbachia, Hibiscus, Juniper, Leather-leaf Fern, Pittosporum, Philodendron, Pothos, Rhododendron, Spathiphyllum and Taxus media.

Use Precautions:

- Do NOT apply more than 400 gallons of spray solution per acre.
- Do NOT make more than one application every 30 days.

Application Instructions: Mix 2.85 – 5.7 lbs. of **FlagStick® Fungicide 70DF** with 100 gallons of water and spray to wet. When using a spray solution of less than 100 gallons per acre, be sure to apply at least 2.5 - 5.0 lbs. of **FlagStick® Fungicide 70DF** per acre. Make repeat applications as necessary to control disease but do not apply more than one spray every 30 days.

DRENCH APPLICATIONS to Aphelandra, Azalea, Boxwood, Cissus, Dieffenbachia, Japanese Holly, Juniper, Monterey Pine, Philodendron, Pieris, Pittosporum, Rhododendron, Schefflera, Spathiphyllum, and Taxus media.

Use Precautions:

- Do NOT apply more than once every 30 days.

Application Instructions: Mix 0.46 – 0.91 lbs. (7.4 – 14.6 oz.) of **FlagStick® Fungicide 70DF** with 100 gallons of water and apply to 400 sq. ft. (2 pints of solution per sq. ft.). Repeat as necessary to control disease. Do not apply more than one spray every 30 days.

SOIL INCORPORATION to Rhododendron, Azalea, or Pieris for control of Phytophthora species **only**.

Use Precautions:

- Incorporate **FlagStick® Fungicide 70DF** in to soil media. Use for plants with established root systems only

Application Instructions: Immediately before potting, mix 0.57 – 0.91 lbs. (9.1 – 14.6 oz.) of **FlagStick® Fungicide 70DF** per cubic yard of soil mix. Further applications as a foliar spray or drench may be made as needed. Do NOT make more than one application every 30 days.

BEDDING PLANTS

FOLIAR APPLICATIONS to Begonia, Pansy, Vinca, Marigold, Zinnia, Petunia, Geranium, and Impatiens.

Use Precautions:

- Do NOT apply more than 400 gallons spray per acre.
- Do NOT make more than one application every 14 days.

Application Instructions: Mix 1.4 – 4.6 lbs. of **FlagStick® Fungicide 70DF** with 100 gallons of water and spray to wet. Repeat as necessary. Do NOT apply more than once every fourteen days.

PHYTOPHTHORA ROOT ROT PREVENTION FOR CONIFERS IN NURSERIES

NOTE: This type of application is NOT registered for use in California.

DIP TREATMENTS to conifers such as Douglas fir, spruce, and pines.

Use Precautions: Be sure to wear chemical/water resistant gloves, goggles or face shield, long pants (coveralls), long sleeved shirt, shoes, and socks when making dip applications.

Application Instructions: Thoroughly immerse plant, including roots by dipping transplants in a solution of 2.5 lbs. of **FlagStick® Fungicide 70DF** per 100 gallons of water. Dip immediately before transplanting.

FOLIAR APPLICATIONS to conifers such as Douglas fir, spruce, and pines.

Application Instructions: Mix 2.85 – 5.7 lbs. of **FlagStick® Fungicide 70DF** with 100 gallons of water and spray to wet. When using a spray solution of less than 100 gallons per acre be sure to apply at least 2.5 - 5.0

10/15

lbs. of **FlagStick® Fungicide 70DF** per acre. Make repeat applications as necessary to control disease but do not apply more than one spray every 30 days.

DOWNY MILDEW CONTROL IN ROSES

FOLIAR APPLICATIONS to field, container, landscape and mini varieties to control downy mildew (*Peronospora sparsa*). NOTE: Follow sound disease sanitation practices to reduce the spread of disease.

Use Precautions:

- Do NOT apply more than 400 gallons of spray solution per acre.
- Do NOT make more than one application every seven days.

Application Instructions: Mix 2.85 lbs. of **FlagStick® Fungicide 70DF** with 100 gallons of water and spray to wet. When using a spray solution of less than 100 gallons per acre, be sure to apply at least 2.5 - 5.0 lbs. of **FlagStick® Fungicide 70DF** per acre. Repeat as necessary but do NOT make more than one spray every seven days.

FIRE BLIGHT SUPPRESSION

FOLIAR APPLICATIONS to ornamental pear, pyracantha, and hawthorn. Apply before disease is present. **FlagStick® Fungicide 70DF** works only as a preventive treatment in this type of application. NOTE: Follow sound disease sanitation practices to reduce the spread of disease.

Use Precautions:

- Do NOT apply more than 400 gallons of spray solution per acre.
- Do NOT make more than one application every seven days.

Application Instructions: Mix 2.85 lbs. of **FlagStick® Fungicide 70DF** in 100 gallons of water and spray to wet. If applying less than 100 gallons of spray solution per acre, then apply 2.5 - 5.0 lbs. of **FlagStick® Fungicide 70DF** per acre. Start applications at the pre-bloom stage and reapply at 7-day intervals throughout the bloom period

If tank mixing **FlagStick® Fungicide 70DF** be sure to read the compatibility statements at the beginning of this section.

BACTERIAL BLIGHT SUPPRESSION

NOTE: This type of application is NOT registered for use in California.

FOLIAR APPLICATIONS to English ivy, schefflera, anthurium, dieffenbachia, spathaphyllum, syngonium, and ficus to suppress *Xanthomonas campestris* pathovars *hederae*, *dieffenbachiae*, *syngonii*, and *fici*. NOTE: Follow sound disease sanitation practices to reduce the spread of disease.

Use Precautions:

- Do NOT apply more than 400 gallons of spray solution per acre.
- Do NOT make more than one application every seven days.

Application Instructions: Mix 2.3 – 4.6 lbs. of **FlagStick® Fungicide 70DF** with 100 gallons of water and spray to wet. . When using a spray solution of less than 100 gallons per acre, be sure to apply at least 2.5 - 5.0 lbs. of **FlagStick® Fungicide 70DF** per acre. Make repeat applications as necessary but do NOT make more than one application every 7 days.

If tank mixing **FlagStick® Fungicide 70DF** be sure to read the compatibility statements at the beginning of this section.

TURF

NOTE: DO NOT apply to home or residential lawns.

11/15

FlagStick® Fungicide 70DF is a fungicide that works systemically. Use in an integrated pest management (IPM) program for the control of *Pythium* diseases (such as blight and root rot, and yellow tuft) on turf grass in golf courses, sod farms and other commercial turf areas. Because **FlagStick® Fungicide 70DF** contains fosetyl al, a known inducer of systemic acquired resistance (SAR) in plants, **FlagStick® Fungicide 70DF** assists in the development of thick, vigorous turf. Turf treated with **FlagStick® Fungicide 70DF** can better withstand plant stresses such as heat, humidity, drying winds, and excessive traffic, which can damage turf. **FlagStick® Fungicide 70DF** aids the turf's natural ability to resist disease. Plus, a thick healthy turf reduces the likelihood of other pests such as weeds and algae from developing.

Use Precautions:

- Do not graze animals on treated turf.
- Do not feed clippings from treated turf to livestock and poultry.
- Do not mow and/or water treated areas until foliage is completely dry.

Application Instructions: Begin using **FlagStick® Fungicide 70DF** when environmental conditions favor disease development and repeat applications as specified in the table below. Apply as a foliar spray at the rate specified below in 1 - 5 gallons of water per 1,000 sq. ft. Use an accurately calibrated sprayer and agitate while mixing and spraying.

Application Interval	Application Rate (Oz. per 1,000 Sq. Ft.)
14 days	4.6
21 days	9.1

COMPATIBILITY

NOTE: Do NOT mix **FlagStick® Fungicide 70DF** with any surfactant, sticker, adjuvant or wetting agent.

FlagStick® Fungicide 70DF may not be chemically compatible with foliar fertilizers or copper based fungicides. Mixing with these products may result in phytotoxicity. **FlagStick® Fungicide 70DF** is NOT physically compatible with flowable formulations of certain pesticides such as Daconil 2787® (chlorothalonil) or Fore® fungicides (mancozeb), Bayleton® 50W (triadimefon), Rubigan™ A.S. (fenarimol) and Spectro™ 90 WDG (chlorothalonil plus thiophanate-methyl). Do not use these products in tank mixes with **FlagStick® Fungicide 70DF**.

To determine the physical compatibility of **FlagStick® Fungicide 70DF** with another product, conduct a "jar test" by mixing a small amount of the tank mix components in a small container. Mix the ingredients with water in the concentration, and in the order of addition desired for the final spray application. Wait 15 minutes. Check for any indication of incompatibility such as separation or curdling. If any indication of incompatibility exists, do not use this application mixture. Be sure to read and follow all directions and precautions on this label and on the labels of any tank mix partner products.

TANK MIXTURES FOR TURF DISEASES

Follow the most restrictive of label limitations and precautions and do not exceed label dosage rates.

Do NOT mix **FlagStick® Fungicide 70DF** with any product which prohibits mixing on its label.

Summer Stress Complex / Summer Decline: **FlagStick® Fungicide 70DF** may be used in a tank mix with products labeled to control Summer Stress Complex / Summer Decline caused by a complex of *Pythium* and *Rhizoctonia* diseases on common turf grass on golf courses, sod farms, industrial or municipal turf areas.

Anthracnose: **FlagStick® Fungicide 70DF** may be used in a tank mix with products labeled for use on common turf grass on golf courses, sod farms, and industrial or municipal turf areas for control of anthracnose. For best results use a preventative program with repeated applications every 14 days. Begin application during conditions conducive for disease development and before symptoms appear.

12/15

Bentgrass Deadspot: FlagStick® Fungicide 70DF may be used in a tank mix with products used to control bentgrass deadspot in common turf grass on golf courses, sod farms, and industrial or municipal turf areas. For best results use a preventative program with repeated applications every 14 days. Begin application during conditions conducive for disease development and before symptoms appear.



Disease Treated	Tank Mix Partner	Application Rate (Oz. / 1,000 FT ²)	Application Interval
Bentgrass Deadspot (<i>Ophiosphaerella agrostis</i>)	FlagStick® Fungicide 70DF	4.6	
	+	+	
	Fore® 80 WP Rainshield™ (mancozeb)	8.0	
	or Daconil Ultrex® (chlorothalonil)	or 3.2	

‡ The following fungicides may be substituted for the companion products listed above for use with FlagStick® Fungicide 70DF in a Summer Stress Complex disease control program: Bayleton® 50† (triadimefon), Chlorostar (chlorothalonil), Cleary's 3336™ WP† (thiophanate-methyl), Cleary's 3336™ F† (thiophanate-methyl), Compass™ (trifloxystrobin), Equus 720 SST (chlorothalonil), Equus DF (chlorothalonil), Consyst WDG (chlorothalonil, thiophanate-methyl), Curalan® EG (vinclozolin), Daconil® Weather Stik®† (chlorothalonil), Daconil ZN® (chlorothalonil), Eagle 20EW Specialty Fungicide (myclobutanil), Echo® 90 DF (chlorothalonil), Echo® 720 Agricultural Fungicide (chlorothalonil), Endorse™ (polyoxin D zinc salt), Fungo® Flo (thiophanate-methyl), Fungo® 50 WSB (thiophanate-methyl), MANhandle™ (mancozeb), Manicure™ 6 Flowable (chlorothalonil), Manicure™ T/O Flowable (chlorothalonil), Medallion® (fludioxonil), Pentathlon® DF (mancozeb), SysStar® WDG (thiophanate-methyl), SysTec® 1998 WDG (thiophanate-methyl), Systhane® WSP (myclobutanil), Terrachlor® 75 WP (PCNB) and Terrachlor® Flowable Fungicide† (PCNB), Terraguard® 50W (triflumizole), Terrazole® 35% WP (etridiazole), Touche EG (vinclozolin), Turfcide® 4F (PCNB).

† NOTE: Disperse these products in to at least a 1/2 filled tank containing water first before adding FlagStick® Fungicide 70DF.

Tank-Mix Mixing Instructions

The order in which products are added to tank mixes can affect FlagStick® Fungicide 70DF ability to go in to solution. Unless otherwise indicated on this label, it is preferable to add FlagStick® Fungicide 70DF to a tank which has been partially filled with water and has no other tank mix components.

If tank-mixing with other pesticides, follow the label directions for the use of the tank-mix partner product and apply at the recommended rate for control of the target disease organism. Be sure to maintain agitation while mixing and spraying.

WARRANTY AND LIMITATION OF DAMAGES

Tessenderlo Kerley, Inc. warrants only that this product conforms to the product description on the label. Except as warranted by this label, Tessenderlo Kerley, Inc. makes no representation or warranty or guarantee, whether expressed or implied, of fitness for a particular purpose of merchantability, or of product performance. Tessenderlo Kerley, Inc. does not authorize any agent or representative to make any such representation, warranty or guarantee. To the extent consistent with applicable law, Tessenderlo Kerley, Inc.'s maximum liability for breach of its warranty or for use of this product, regardless of the form of action, shall be limited to the purchase price of this product. To the extent consistent with applicable law, buyer and user acknowledge and assume all risks and disposal liability resulting from handling, storage, use and disposal of this product. If buyer does not agree with or accept these warranty and liability limitations, buyer may return the unopened container to the place of purchase for full refund. Buyer's use of this product shall constitute conclusive evidence of buyer's acknowledgement and acceptance of the forgoing limitations. Some states do not allow the exclusion of implied warranties or the limitation of certain damages, so the above may not apply. The purchase, delivery, acceptance and use of this product by the buyer are subject to the terms and conditions of seller's sales invoice for this product.

Bayleton® and Prostar® are registered trademarks of Bayer.
 Chipco® and Signature® are registered trademarks of Bayer Crop Science LP
 Compass® is a registered trademark of Bayer.
 Endorse™ is a trademark of Avesta.
 Banner®, Maxx®, Heritage®, Daconil®, Daconil Ultrex®, Weather Stik®, Daconil Zn®, and Medallion® are registered trademarks of Syngenta.
 Dithane®, Systhane® and Fore® are registered trademarks of Dow AgroSciences Company.
 Manicure™, MANhandle™ and Lesco™ are trademarks of Lesco, Inc.
 Curalan® is a registered trademark of BASF Corporation.

3336™ and Spectro™ are trademarks of Cleary Chemical Company.
Systec® and Syster® are registered trademarks of Regal Chemical Company.
Echo® is a registered trademark of SIPCAM AGRO USA, Inc.
Fungo® is a registered trademark of the Scotts Company.
Terrachlor®, Terraguard®, Terrazole® and Turfcide® are registered trademarks of Uniroyal Chemical Company, Inc.
Pentathlon® is a registered trademark of Griffin L.L.C.
NovaSource® and **FlagStick® Fungicide 70DF** are registered trademarks of Tessenderlo Kerley, Inc.

FlagStick® Fungicide 70DF is not manufactured or distributed by Bayer Crop Science LP, seller of Chipco® Signature®.

EPA 20120910

NOTIFICATION
OCT 24 2012

