

## UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

Michael Kellogg Tessenderlo Kerley, Inc. c/o Pyxis Regulatory Consulting, Inc. 4110 136<sup>th</sup> Street, NW Gig Harbor, WA 98332

NOV 14 2011

Subject:

Label Amendment (Supplemental Label)

Sinbar WDG

EPA Reg. No. 61842-27

Application Dated: October 27, 2011

Dear Mr. Kellogg,

The supplemental labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable.

At your next label printing, or within eighteen (18) months of the date of this letter, whichever comes first, you must incorporate this supplemental labeling into the main product labeling. A stamped copy of your label is enclosed for your records.

If you have any questions, please contact Emily Hartman of my staff at (703) 347-0189 or hartman.emily@epa.gov.


Sincerely.

Kable Bo Davis, Product Manager 25

Herbicide Branch

Registration Division

Office of Pesticide Programs



GROUP 5 HERBICIDE

### SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR SINBAR WDG BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Sinbar WDG and this supplement.

### Sinbar® WDG

EPA Reg. No. 61842-27

**Expiration Date:** This supplemental label expires on November 9, 2014 and must not be used or distributed after this date.

## FOR USE OF SINBAR WDG ON WATERMELON TO CONTROL BROADLEAF WEEDS (EXCEPT CALIFORNIA)

# KEEP OUT OF REACH OF CHILDREN CAUTION

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Sinbar WDG should be used for control of broadleaf weeds (e.g. annual morningglory) in watermelons.

A maximum of one pre-emergence ground application of Sinbar WDG (active ingredient – terbacil) may be applied at a rate of 2 to 4 ounces product per acre (0.1 to 0.2 pound active ingredient per acre). Do not exceed 4 ounces of product per acre per year (0.2 pound active ingredient per acre per year). A 70-day preharvest interval (PHI) is required.

Sinbar WDG may be used for direct-seeded or transplant watermelons. Where seeded watermelons are used as cultural practice, make applications after planting, but before the crop emerges. Where transplants are used as a cultural practice, apply Sinbar WDG pre-transplant.

Where plastic mulch is used as a cultural practice, Sinbar WDG may be applied pre-emergence under plastic mulch or to row middles.

Sinbar WDG may also be applied at 2 to 4 ounces broadcast per acre over the plastic mulch prior to transplanting, or prior to punching holes into the plastic mulch for transplanting. Sinbar WDG must be washed off the surface of the plastic mulch with a minimum of 0.5 inches of rainfall or irrigation prior to transplanting watermelons.

Use the lower rate on coarse, textured soils lower in organic matter.

### **PRECAUTIONS**

Do not apply over the top or allow spray to contact crop as unexpected crop injury may result.

Do not apply within 70 days of harvest.

A reduction in residual weed control may occur under adverse environmental conditions such as heavy rainfall or when low rates are used on soil high in organic matter.

This labeling must be in the possession of the user at the time of pesticide application.

Read the "CONDITIONS OF SALE – LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES" in the label booklet for Sinbar WDG before using.

Sinbar is a registered trademark of Tessenderlo Kerley, Inc.

Tessenderlo Kerley, Inc. 2255 N. 44<sup>th</sup> Street, Suite 300 Phoenix, AZ 85008 USA 1-800-525-2803

[EPA approval date]