



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

August 23, 2021

Jolanta Ozatalay
Regulatory Affairs Project Manager
Fine Americas, Inc
c/o Fine Agrochemicals, LTD
1850 Mt. Diablo Blvd #405
Walnut Creek, California 94596

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment – Adding the referral statement to First Aid section and maximum rate for Mandarins to the Directions for use (DFU).
Product Name: Falgro® 2X LV
EPA Registration Number: 62097-32
Application Date: 07/30/2021
OPP Action Case Number: 00314419

Dear Mr. Ozatalay:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to your company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the U.S. Environmental Protection Agency (EPA). If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims

Page 2 of 2

EPA Reg. No. 62097-32

OPP Action Case No. 00314419

made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.


If you have any questions, please contact Menyon Adams by phone at (703) 347-8496 or via email at adams.menyon@epa.gov.

Sincerely,



Andrew Bryceland, Team Leader
Biochemical Pesticide Branch
Biopesticides and Pollution
Prevention Division (7511P)
Office of Pesticide Programs

Enclosure



MASTER LABEL

FALGRO® 2X LV

A plant growth regulator solution for use on food and non-food crops

ABN: FLORGIB 2X LV

Sublabel A: Agricultural Uses

Sublabel B: Ornamental Uses

Active Ingredient:

Gibberellic Acid (GA₃) 6.18%

Other Ingredients: 93.82%

Total: 100.00%

Falgro 2X LV contains 2.0 grams active ingredient per fluid ounce of formulated product.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

[See attached label booklet for first aid, precautionary, storage and disposal statements, and Directions for Use.]

FLAMMABLE – Keep away from heat and open flame

EPA Reg. No. 62097-32

EPA Est. No.

NET CONTENTS: 1 qt, ½ gal, 1 gal, 2.5 gal, 15 gal, 30 gal, 55 gal
Batch No.

Manufactured by:

© Fine Agrochemicals Limited

Hill End House

Whittington

Worcester, WR5 2RQ

United Kingdom

Falgro® is a registered trademark of Fine Agrochemicals Ltd.

SUBLABEL A: Agricultural Uses

FALGRO® 2X LV

A plant growth regulator solution for use on food and non-food crops

Active Ingredient:

Gibberellic Acid (GA₃) 6.18%

Other Ingredients: 93.82%

Total: 100.00%

Falgro 2X LV contains 2.0 grams active ingredient per fluid ounce of formulated product.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

[See attached label booklet for first aid, precautionary, storage and disposal statements, and Directions for Use.]

FLAMMABLE – Keep away from heat and open flame

EPA Reg. No. 62097-32

EPA Est. No.

NET CONTENTS: 1 qt, ½ gal, 1 gal, 2.5 gal, 15 gal, 30 gal, 55 gal

Batch No.

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.

IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information on this pesticide product (including general health concerns or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378, Monday through Friday, 8:00 AM to 12:00 PM Pacific Standard Time. In the event of a medical emergency, call your poison control center at 1-800-222-1222.</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Causes moderate eye irritation. Harmful if swallowed, absorbed through the skin, or inhaled. Avoid contact with eyes, skin, or clothing and breathing spray mist. Wear goggles or safety glasses. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical-resistance selection chart. Applicators and other handlers must wear:

- Protective eyewear
- Coveralls worn over short-sleeved shirt and short pants
- Shoes plus socks
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride or Viton

Follow manufacturer’s instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE items separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of the gloves before removing. As soon as possible, wash thoroughly and change into clean clothing

ENVIRONMENTAL HAZARDS

For terrestrial uses. Do not apply directly to water, to areas where surface water is

present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

PHYSICAL AND CHEMICAL HAZARDS

FLAMMABLE. Keep away from heat and open flame. FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

COMPATIBILITY

Conduct a compatibility test when you plan to mix Falgro 2X LV with other products. To determine the physical compatibility of Falgro 2X LV with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to approximately one quart of water with agitation. Add dry formulations first, then flowables, and then emulsifiable concentrates last. After thorough mixing, allow this mixture to stand for 5 minutes. If the combination remains mixed or can be readily remixed, it is physically compatible. Once compatibility has been proven, use the same procedure for adding products to the spray tank. Follow the more restrictive labeling requirements of any tank mix partner. Do not tank mix with products whose label prohibits tank mixing. Treat a small test plot if new combinations of products are being used for the first time.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your state or tribe, consult the state or tribal agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE) and restricted entry intervals. The requirements in this box only apply to uses of this product covered by the Worker Protection Standard.

Do not enter or allow entry into treated areas during the restricted entry interval level (REI) of 4 hours unless wearing appropriate PPE.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as, plants, soil or water is:

- Coveralls worn over short-sleeved shirt and short pants
- Shoes plus socks
- Protective eyewear
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride or Viton.

PRODUCT INFORMATION

Falgro 2X LV is a plant growth regulator solution for use on food and non-food crops to affect the growth characteristics of specified crops:

When used on table grapes, application of Falgro 2X LV for 'Stretch' produces looser cluster forms, allows better air circulation to aid in the control of bunch rot, increases light penetration to aid in sugar development and reduces the cost of thinning. Application of Falgro 2X LV for 'Thinning' (decreased berry set) reduces hand thinning costs and hastens maturity. Application of Falgro 2X LV for 'Sizing' produces larger berries and clusters when used in conjunction with girdling and thinning techniques. Falgro 2X LV may also be applied as a cluster dip or directed spray to berry clusters/bunches to increase berry size.

When used on grapefruit, application of Falgro 2X LV reduces water loss during storage and can be used to manage harvest timing. When applied before color change, use of Falgro 2X LV produces the greatest delay in rind aging and the firmest rind.

When used on lemons and limes, application of Falgro 2X LV decreases rind yellowing/aging and the amount of small tree ripe fruit, thus allowing a more desirable production pattern relative to market demand. Application of Falgro 2X LV is cumulative so that when applied for two years in succession, a larger difference in maturity and harvest pattern occurs.

On navel oranges, Falgro 2X LV may be used to reduce rind disorders (e.g., rind staining, water spotting, sticky surface, puffy rind and pressure rupture), aging and to manage harvest timing.

When used on tangerine hybrids, application of Falgro 2X LV increases fruit set and yields, and delays rind aging and increases peel strength in hybrids with pollination problems such as Minneola, Orlando, Robinson and Sunburst.

Falgro 2X LV can be used to increase fruit set and yield in Clementine Mandarin.

Falgro 2X LV may also be used for post-harvest applications on citrus. On lemons, Falgro 2X LV can be used to reduce sour rot pathogen (*Geotrichum candidum*) infection and to delay rind senescence and color changes to extend storage life.

Other uses of Falgro 2X LV include:

- Stimulates fruit growth on bananas when plant is stressed due to insect or disease pressure or adverse weather conditions.
- Improves fruit set on blueberries when natural fruit set is poor due to reduced honeybee activity as a result of adverse weather conditions.
- Maintains and extends high fruit bearing and reduces the occurrence of 'blind' nodes on red tart cherries.
- Produces larger, brighter colored, and firmer sweet cherries. This treatment may delay color development and harvest. This delay in color and maturity may be

- especially evident in blonde varieties such as Rainier, Royal Ann, Stardust, etc.
- Use of multiple applications of Falgro 2X LV on sweet cherries may result in a reduction in return bloom in the year following applications.
 - An application of Falgro 2X LV can be made to Tart cherry trees to maintain and extend high fruiting capacity by promoting spur formation and reducing the occurrence of "blind" nodes.
 - Reduces the flowering and fruiting of non-bearing sweet and tart cherries and other stone fruit, minimizing the competitive effect of fruiting on tree development.
 - Seasonal applications of Falgro 2X LV increase fruit firmness and improves fruit quality on stone fruit.
 - Application of Falgro 2X LV reduces internal browning, increases size and improves fruit quality on Italian prune.
 - Applications of Falgro 2X LV can be made to pecan trees to extend leaf retention and maintain green foliage.
 - On strawberries, application of Falgro 2X LV increases the production of runners by mother plants.
 - Application of Falgro 2X LV accelerates maturity, allowing early harvesting of artichoke.
 - On carrots, application of Falgro 2X LV delays leaf senescence and reduces the incidence of infection from *Alternaria dauci*.
 - On celery, application of Falgro 2X LV increases plant height and yield and increases plant ability to overcome stress due to cold weather conditions or saline soils, and obtain earlier maturity.
 - Application of Falgro 2X LV produces more uniform bolting and greater seed production on lettuce grown for seed.
 - On melons and cucumbers, application of Falgro 2X LV stimulates fruit set in extended periods of cool weather.
 - Application of Falgro 2X LV helps break dormancy and increase yield of forced rhubarb.
 - Application of Falgro 2X LV helps break dormancy and stimulates uniform sprouting on seed potato.
 - On spinach, Falgro 2X LV improves the quality, increases yield and aids harvest.
 - On watercress, applications of Falgro 2X LV helps enhance growth in adverse weather conditions, plants to resume growth following insect and disease infestations and helps to increase root free stem length during low light or short day conditions.
 - On cotton, application of Falgro 2X LV promotes early plant growth, increases seedling vigor and helps overcome stress induced by cool weather.
 - Application of Falgro 2X LV increases the yield and aids picking of seeded and seedless varieties of Fuggle hops and similar varieties.
 - Falgro 2X LV as a seed treatment promotes germination and emergence of semi-dwarf and tall rice varieties.
 - Post-emergence application of Falgro 2X LV on rice prior to permanent flooding promotes uniform and vigorous growth of semi-dwarfing varieties allowing earlier flooding with its associated agronomic benefits.
 - Falgro 2X LV can be applied to sugarcane to maintain yields in older plantings,

increase bio-mass and stimulate growth before harvest of cane in older production fields (>3 years).

Use only as directed. Read thoroughly and understand the label before making applications.

The term "grams of active ingredient per acre" is represented by "g ai/acre" throughout this product label. See Conversion Table at end of Directions for Use section.

Thoroughly spray all parts of the plant or crop.

Use a clean empty spray tank for preparing and mixing the solution.

The pH of water used to prepare the spray solution must be less than 8.5.

Prepare solution concentrations by mixing the required amount of product with water only.

Degradation of the active ingredient may occur if solution is held over an extended period of time. A reduction in effectiveness may occur if the spray solution is held over a 24 hour period. Dispose of any unused spray material at the end of the day. Refer to the Storage and Disposal section of this label for pesticide disposal instruction.

When a range of rates is indicated, consult your local experimental station, distributor, or agricultural extension agent for the best program suited to your local conditions.

Falgro 2X LV is best absorbed into the plant under slow drying conditions. Night-time applications will be more effective when day-time conditions cause rapid drying of applications. This allows for the spray solution to be completely absorbed. Re-apply Falgro 2X LV if significant rain occurs within 2 hours of application.

Falgro 2X LV has a 0-day pre harvest interval (PHI).

Do not apply using ULV application methods.

For aerial applications spray volumes must be greater than 2 gallons per acre (10 gallons per acre for tree crops).

Consult your local experimental station, distributor, or agricultural extension agent regarding the compatibility of gibberellic acid with other compounds.

SPRAYING GUIDE FOR GRAPES

Make applications to grapes by ground sprayer. Use a properly calibrated sprayer to ensure uniform and complete coverage of flower clusters and berries.

TABLE GRAPES

Table 1 – Application rates and timing for table grapes

NOTES:

¹ High doses can cause over-thinning or an excess of ‘shot’ berries, especially on young or vigorous vines.

² Conduct trials with small numbers of plants using recommended rates to determine the optimum rates for the situation before applying to the crop.

³ A rate of 40-50 ppm/ai is used when dipping **or** spraying the cluster directly.

Seedless Varieties	“Stretch”		“Thinning”		“Sizing”		
	Rate g ai/A	Timing	Rate g ai/A ¹	Timing	Rate g ai/A	Timing	Target Diameter
Thompson	8-24	1-3 sprays before bloom.	8-20	1-4 sprays during bloom. If bloom is extended apply 2 nd application 1-7 days after the 1 st application.	32-128	1-4 sprays starting at these berry sizes. All applications to be made within 14 days.	3-5 mm
Flame	8-24		3-16		20-128		6-9 mm
Perlette	8-24		²		32-128		4-5 mm
Raisin	8-24	Flower clusters 2-7 inches long.	3-12		4-20		3-5 mm
Others	²		0.5-12		8-60		3-14 mm
Black Corinth (Zante Currant)				²	1-12	One application 3-5 days after full bloom but before shatter begins.	
Seeded varieties²					“Sizing” “Reduced Berry Shivel” in Emperor		
Calmeria					20 ³	1 spray 14 days after shatter. Berry diameters of approx. 10-15 mm. May increase berry size.	12-16 mm
Christmas Rose							12-16 mm
Emperor							12-16 mm
Red Globe							12-18 mm
Rogue							12-16 mm
Queens							12-15 mm

BUMP SPRAY (Thompson Seedless)

To initiate the beginning of the berry growth period, apply 16-24 g ai/acre during the period between the last thinning spray and the first sizing spray.

BERRY SIZING CLUSTER DIP/DIRECTED SPRAY (SEEDED & SEEDLESS TABLE GRAPES) (All approved states except California)

Apply Falgro 2X LV at 20 to 50 ppm as a dip or directed spray to grape clusters when berries are 3-14 mm in size. To prepare the 20 to 50 ppm ai solution, mix 0.2 to 0.5 fl oz (5.6 to 14 ml) of Falgro 2X LV per 5 gallons of water. Due to varietal differences, perform testing on a small number of bunches for a variety that the grower does not have prior experience with using this application method or rates.

WINE GRAPES

To increase cluster length and improve air circulation and light penetration within the cluster, make a single spray application. Under certain conditions, i.e., moderate temperatures and high humidity, this application may help reduce the incidence of

bunch rot and sour rot.

The application of gibberellic acid on seeded wine grape cultivars will cause some reduction in yield. Yield reduction may result from an increase in shot berries in the year of application, and/or a reduction in fruitfulness in the first and second year following the application.

Table 2 – Application rates and timing for wine grapes

Variety	Rate g ai/A	Timing
Palomino Sauvignon Blanc Tinta Madeira	0.4-1	1 spray when average flower cluster length is 3-4 inches. Do not make application less than three weeks before full bloom. Use is based on 100 gallons of water per acre.
Aleatico Carignane Chardonnay Chenin Blanc French Colombard Pinot Noir Valdepenas	1-2	
Barbera Petite Sirah Zinfandel	2-4	
Green Hungarian	4-8	
Grenache Alicante	8	
Salvadore	8-16	

SPRAYING GUIDE FOR CITRUS

Use a properly calibrated sprayer to ensure uniform and complete coverage of all fruit. Do not exceed maximum application rates. Do not apply to trees of low vigor or suffering stress of any kind as this can cause severe fruit and/or leaf drop. Some drop of older leaves may occur after application of gibberellic acid. Never tank mix Falgro 2X LV with materials creating a high pH in the tank, e.g., white wash. To avoid significant leaf and/or fruit drop, do not use copper fungicides and/or oils within three weeks of a Falgro 2X LV application.

GRAPEFRUIT (All approved states except California)

All Grapefruit Varieties **Except** Star Ruby:

To delay rind aging and prevent pre-harvest drop of mature fruit and increases peel strength: Spray 16-56 g ai/acre as a dilute spray in a minimum of 250 gallons of water per acre. Ensure thorough wetting.

Early spray (Before color change)

Make one application 14 days before color break (usually August to September).

Late spray (After color break)

Make one application after marketable color has developed (usually October to December). Late application may cause some re-greening of the fruit.

Do not apply to areas that are destined to be harvested early as treatment will delay color change. Application after coloring will cause fruit to re-green if left on the tree for an extended period of time. Do not apply after December or after trees break dormancy in order to avoid adversely affecting the next crop. Efficacy of **grapefruit treatment varies from season to season, depending on environmental conditions.**

To increase fruit set and yield: Make one application December – January of 15-25 g ai/acre as a dilute spray of 125-175 gallons of water per acre with a pure organo-silicone surfactant at a concentration of 0.05% (6 fl. oz./100 gallons).

Grapefruit - Star Ruby Variety:

To reduce early fruit drop: Make one application of 25-35 g ai/acre as a dilute spray in a minimum of 250 gallons of water per acre during bloom.

Note: Efficacy of treatment varies from season to season, depending on environmental conditions. Use a fertilizer and watering program to maintain optimal tree health and vigor.

To enhance fruit set and yield: Make one application December – January of 15-25 g ai/acre as a dilute spray of 125-175 gallons of water per acre with a pure organo-silicone surfactant at a concentration of 0.05% (6 fl. oz./100 gallons).

LEMONS & LIMES

Make one application of 10-32 g ai/acre as a dilute spray or concentrate, ensuring thorough spray coverage, when the target crop is $\frac{1}{2}$ - $\frac{3}{4}$ full size and still green.

NAVEL ORANGES

Apply 16-48 g ai/acre as a dilute spray or concentrate ensuring thorough spray coverage. To avoid reduced yield in the following season, do not spray Navel oranges from January to July. Do not apply to areas that are destined to be harvested early as treatment will delay color change.

Early spray (before color change)

Make one application 14 days before color break (usually August to November).

Application at this timing produces the greatest delay in rind aging and the firmest rind possible.

Late spray (after color break)

Make one application after marketable color has developed (usually October to December). Late application may cause some re-greening of the fruit.

To enhance fruit set and yield (Florida use only): Make one application December – January of 15-25 g ai/acre as a dilute spray of 125-175 gallons of water per acre with a pure organo-silicone surfactant at a concentration of 0.05% (6 fl. oz./100 gallons).

TANGERINE HYBRIDS

To enhance fruit set and yield: Make one application of 8-30 g ai/acre as a dilute spray during full bloom ensuring all foliage is wetted. A reduction in fruit size, retardation of color development and a slight increase in the drop of mature leaves may occur with this type of treatment.

To reduce rind disorders: Make one application of 20-40 g ai/acre as a dilute spray 14 days before color break in sufficient water to ensure thorough spray coverage. Do not apply to areas that are destined to be harvested early as treatment will delay color change. To avoid pre-harvest staining, do not apply after coloring. Variations in rind color development can occur if applications are made during coloring.

VALENCIA ORANGES (All approved states) & OTHER ROUND ORANGES (All approved states except California)

To delay rind aging and softening: Make one application of 40-80 g ai/acre as a dilute spray or concentrate in August to October.

This application will delay color development and re-green any mature fruit present at application. After marketable color has been achieved, beneficial treatment affects will gradually lessen the longer the fruit remains on the tree.

To enhance fruit set and yield (Florida use only): Make one application December – January of 15-25 g ai/acre as a dilute spray of 125-175 gallons of water per acre with a pure organo-silicone surfactant at a concentration of 0.05% (6 fl. oz./100 gallons).

OTHER CITRUS

CLEMENTINE MANDARIN

Make EITHER 1-2 applications of 1-40 g ai/acre as a dilute spray from 50% petal fall up to 3 weeks after petal fall OR 1-4 applications from early bloom up to four weeks after petal fall. [Limit of 1-3 full applications in California]. Use a sufficient spray volume to ensure thorough coverage of tree canopy, typically, 1-8 g ai per 100 gallons of spray solution. Allow at least three days between sprays. Do not exceed 160 g ai/acre per season.

AMBERSWEET (Florida use only)

Make one application in January of 15-25 g ai/acre as a dilute spray of 125-175 gallons of water per acre with a pure organo-silicone surfactant at a concentration of 0.05% (6 fl. oz./100 gallons).

POSTHARVEST APPLICATIONS FOR CITRUS

LEMONS

Apply 50-100 ppm ai in 10 gallons of storage wax diluted as per wax label instructions.

YELLOW LEMONS AND OTHER MATURE CITRUS

Apply Falgro 2X LV postharvest at a rate of 50-100 ppm ai or add 1-2 fl. oz. of product to 10 gallons of storage wax diluted as per wax label instructions.

SPRAYING GUIDE FOR OTHER FRUIT CROPS

BANANAS

To treat fruit, make one application of 1-6 g ai/acre of Falgro 2X LV by air or ground every 30 to 90 days throughout the year. Monthly applications can be made up to 6 months prior in anticipated weather stress periods.

BLUEBERRIES (All approved states except California)

Highbush blueberry (varieties such as Berkley, Bluecrop, Blueray, Concord, Coville, Earliblue, Jersey, Stanley, Walcott, Weymouth, and 1316A):

Make EITHER a single application of 80 g ai/acre in 40-100 gallons of water, OR two applications of 40 g ai/acre in 40 to 100 gallons of water. Make the single application at full bloom (75% of the flowers are fully open). When making two applications, spray the first one at full bloom and the second one within 10 to 14 days. For Weymouth, application can be delayed up to two weeks after bloom to increase size of 'shot' berries.

Rabbiteye blueberry (varieties such as Aliceblue, Beckyblue, Bonita, Brightwell, Climax, Delite, Tiftblue, and Woodward):

Make EITHER a single application of 40-80 g ai/acre in 40-100 gallons of water when most of the flowers are elongated but not yet open (bloom stage 5), OR two to four applications of 20-40 g ai/acre in 50-300 gallons of water every 10-14 days starting at bloom stage 5. Ensure thorough spray coverage.

CHERRIES – RED TART (All approved states except California)

Make a single application 14-28 days after bloom when 3-5 terminal leaves are fully expanded or at least 1-3 inches of terminal growth have occurred. Apply Falgro 2X LV according to the table below as a dilute spray in 100-150 gallons of water/acre. Lower water volumes can be used, but plants must not be overdosed. Treatment results are not apparent until 2-3 years after initial application and are dependent on annually repeated applications.

Table 3 – Application rates for red tart cherries

Tree age (years)	Grams ai/acre
6 – 10	4 – 6
11 – 15	8 – 10
16 – 20	10 – 14
21 +	14 – 18

These application rates are based on tree vigor. Rates must be adjusted to the vigor expressed in each orchard. Use higher rates for trees of low vigor and the lower rates

for trees of high vigor. Treatment with Falgro 2X LV cannot overcome the effects of nutritional, water, pest, disease or other stress on trees. Excessive application will increase vegetative growth at the expense of fruit production the following year.

CHERRIES – SWEET

SINGLE APPLICATION

Make one application of 16-48 g ai/acre (do not exceed a total of 48 g ai/acre) as a dilute spray on large mature trees when the fruit is light green to straw colored in sufficient water to ensure thorough wetting.

SPLIT APPLICATIONS

Make the first application of 8-24 g ai/acre when fruit is light green to straw colored. Make the second application of 8-24 g ai/acre, 7 to 14 days later. Do not exceed a total of 48 g ai/acre.

MULTIPLE APPLICATIONS (All approved states except California)

Make the first application of 6-24 g ai/acre when the majority of fruit is light/translucent green. Make subsequent applications of 2-24 g ai/acre at 3 to 7 day intervals. Do not apply Falgro 2X LV later than 14-days before harvest. Do not exceed a total of 48 g ai/acre. Apply in sufficient water to ensure thorough wetting.

Treatment may delay color development and harvest.

CHERRIES - NON-BEARING SWEET AND TART (All approved states except California)

Make one application of 10-20 g ai/acre as a dilute spray in 25-50 gallons per acre 2-4 weeks after bloom. (This rate is based on a tree density of 100 trees/acre). In conditions of low vigor, 2 applications can be made allowing at least a 7-day interval between applications.

Do not treat trees in their first year. Treat in the second season to reduce fruiting in the third season and treat again in the third season if it is necessary to reduce fruiting in the fourth. Discontinue the use of Falgro 2X LV one year before commercial harvest is planned.

CRANBERRY (All approved states except California)

Apply Falgro 2X LV to decrease or eliminate fruit set in the year of application. Make one application of 10-50 g ai/acre at early bloom (2-5% scatter bloom) in sufficient water to ensure thorough coverage. To avoid applications having no effect or the opposite effect (increased fruit size), do not apply later than indicated.

PECAN (All approved states except California)

- Applications of Falgro 2X LV can be made to pecan trees to extend leaf retention and maintain green foliage. Make 1-4 applications of 10 g ai/Acre beginning in July. Applications may continue through October as needed. Complete spray coverage is essential. Use sufficient water spray volume to achieve thorough

coverage. In most cases 100 gallons per acre has been shown to be effective.

Falgro 2X LV may be tank mixed with pesticides and other materials as needed. As all possible combinations of products that may be applied have not been tested for compatibility, it is recommended that a jar test be performed with potential tank mix partners before mixing a tank for application. Test tank mixes for which there is no previous experience on a limited area to assure no damage or undesirable affect occurs from the particular tank mix.

Use Limitation:

- Do not make more than 1 application of Falgro 2X LV in July. Using more than 1 application in July may result in reduced return bloom.

PINEAPPLE (All approved states except California)

To improve fruit sizing: Make one application of 400 g ai/acre of Falgro 2X LV by air or ground 14-18 weeks after flowering.

To improve uniformity of fruit maturity and enhance harvest efficiency: Make the first application within a few days after planting once plants are established. Repeat applications may be made at 3 to 4 week intervals. The application rate per application is 12-24 g ai/acre.

OTHER STONE FRUIT

Make one application of 16-32 g ai/acre as a dilute spray 1-4 weeks prior to the beginning of harvest. Use sufficient water to ensure thorough spray coverage. Applications in May through July may cause a reduction in flower counts in the year following application.

ITALIAN PRUNE

Make one application of 16-48 g ai/acre in sufficient water to ensure thorough coverage. Apply 4-5 weeks before anticipated harvest.

NON-BEARING STONE FRUIT (All approved states except California)

Make one application of 20-80 g ai/acre during flower bud initiation for the following years' development. Use sufficient water to ensure thorough spray coverage of the tree canopy. Do not treat trees in their first year. Treat with Falgro 2X LV in the second season to reduce flowering and fruiting in the third season. Treat again in the third season if flower/fruit reduction is desired in the fourth season. Discontinue the use of Falgro 2X LV one year before commercial harvest is planned.

STRAWBERRIES (All approved states except California)

Make one application of 15-25 g ai/acre in 100 gallons of water/acre 10-30 days after planting, when plants have 1-6 leaves. Spray to the point of run-off. For best results, do not apply to plantings after mid-May. Do not apply to fruiting plants.

SUGARCANE (All approved states except California)

Falgro 2X LV can be applied to sugarcane to maintain yields in older plantings, increase

bio-mass and stimulate growth before harvest of cane in older production fields (>3 years). Make an application of 1.0-2.0 g ai/acre to sugarcane when at 1st to 5th internode stage in a minimum of 20 gal/acre spray volume. The addition of a non-ionic surfactant may increase activity.

SPRAYING GUIDE FOR VEGETABLE CROPS

ARTICHOKE

Make 1-3 applications of 10-20 g ai/acre to perennials at bud initiation. For annuals, make 1-4 applications at 2-week intervals, beginning at the fourth true leaf stage. Use sufficient water to ensure thorough wetting of the entire plant.

CARROTS (Fresh and processing)

Make first application of 1-6 g ai/acre 4-6 weeks after emergence using commercial ground or aerial equipment with spray concentrations of 20-30 ppm. A second application 14 days later may be required to achieve desired foliar recovery in severe disease situations or cool weather. Do not make more than two applications per crop cycle.

Dilutions of greater concentrations can increase the risk of excessive top growth, which can result in a delay in root development. This is especially true with a second application.

CELERY

Make one application of 2.5-10 g ai/acre, 1-4 weeks prior to harvest. Use 25-50 gallons of water/acre by ground application or 5-10 gallons of water/acre for aerial application (except in California). Use lower concentrations if applying 3-4 weeks before harvest and higher concentrations within 1-2 weeks before harvest. Do not apply earlier than 4 weeks before harvest as bolting can occur.

LETTUCE FOR SEED

Make 1-4 applications of 1-4 g ai/acre as a dilute spray at 2-week intervals beginning at the fourth true leaf stage. Use sufficient water to ensure thorough wetting.

MELONS AND CUCUMBERS (All approved states except California)

For cantaloupes and watermelons, make one application of 1-4 g ai/acre in sufficient water to ensure thorough wetting prior to bloom. Make 2 further applications at 10-14 day intervals. For cucumbers, 3-4 applications following fruit set may be required. For maximum benefit of these treatments to be achieved, vines must be in good condition.

PEPPERS (All approved states except California)

To promote plant growth: In areas with short growing seasons or where low temperatures cause slow plant growth, make 1-2 applications of 1-3 g ai/acre in 25-50 gallons of water/acre starting 2 weeks after planting. Repeat at 2-week intervals.

To promote plant growth and increase fruit set: Make 1-2 applications of 1-3 g ai/acre in 25-50 gallons of water/acre during flowering. Use the higher rate for areas and

varieties with pollination and fruit set problems.

To increase fruit size: Make one application of 1-3 g ai/acre in 25-50 gallons of water/acre at the beginning of picking. Use the higher rate for plants carrying high numbers of fruit.

RHUBARB

If the rest period is not broken, make a single application of 2 fl. oz. of a solution containing 20 g ai in 10 gallons of water to each cleaned crown. When the rest period is broken by cold weather, apply 2 fl. oz. of a solution containing 10 g ai in 10 gallons of water to each cleaned crown. Maintain forcing house temperatures at 40°F-50°F for 24 hours following application. To prevent lower yields and poor stalk color, keep temperatures below 50°F.

SEED POTATOES

Dip freshly dug seed potatoes in a solution of 0.2-0.4 g ai in 100 gallons of water before planting.

Potatoes treated with Falgro 2X LV must not be used for food or feed purposes. Do not treat rested seed and use the lower rate for dormant seed if soil temperatures are high.

SPINACH - FALL AND OVERWINTERED (All approved states except California)

Make one application of 6-10 g ai/acre 10-18 days before each anticipated harvest in 10-50 gallons of water if applied using ground equipment, or a minimum of 5-10 gallons of water/acre if sprayed by air. Ideally, spray in the early morning when dew is present on the crop and daytime temperatures are between 40°-70°F. Maximum benefit is seen from this treatment when low temperatures would limit the growth of untreated spinach.

Do not apply to spring sown spinach. Do not treat spinach after mid-winter or if temperatures are expected to exceed 75°F within several days of application as this can induce bolting.

SPINACH, MUSTARD GREENS, COLLARD GREEN AND TURNIP GREENS (All approved states except California)

Falgro 2X LV may be applied to certain leafy vegetables to facilitate harvest, increase yield and improve quality of fall and over-winter crops. Make one application of 4-10 g a.i./acre 10-18 days before each anticipated harvest. Make applications in 10-50 gallons of water per acre by ground sprayer or in a minimum of 5-10 gallons of water per acre by air.

Make applications when daytime temperatures are between 40° F – 70° F and during early morning hours when dew is present on crop. Maximum benefit is obtained when below normal temperatures prevail following application and growth would be otherwise slowed in untreated crops. When applied to promote growth of second cutting, wait until some regrowth has started before spraying.

Since the promotion of bolting has been known to occur, do not apply after the mid-winter period or if temperatures are expected to exceed 75° F within several days of application. Do not apply on spring plantings.

WATERCRESS (All approved states except California)

Make 1 to 2 applications of 15-25 g ai/acre/crop of Falgro 2X LV at 3 to 7 days prior to harvest. Apply in a spray volume of 50 to 100 gallons of water per acre.

SPRAYING GUIDE FOR OTHER CROPS

Do not exceed the application rates noted below or reduce the time interval between applications. Do not apply during extended warm periods if night temperatures exceed 65°F.

COTTON (All approved states except California)

Apply 1-6 g ai/acre by furrow application to the seed or as a foliar spray between the cotyledon and 5-leaf stage. Apply in 5-40 gallons of water/acre for ground applications or 3-10 gallons/acre by air. Repeat applications as necessary, but do not exceed 4 in total. Use higher rates when temperatures are likely to average 75°F or less during the 14 days after application. To avoid excessive growth, do not over treat.

HOPS (Northwestern US only)

Make one application of 4-6 g ai in 100-150 gallons of water/acre when vine growth is 5-8 feet long.

RICE SEED TREATMENT

Falgro 2X LV can only be applied to seed intended for drilling or dry broadcast. Do not apply to rice used in a 24-hour soak prior to broadcasting. Apply 1-2 g ai in 8-20 fl. oz. of water per 100 pounds of seed.

Falgro 2X LV can be applied using mist-treatment equipment, but best effect is achieved using the higher water volume. Half fill the seed treatment tank with water then add the required amount of Falgro 2X LV and mix thoroughly while adding any further seed treatment materials before making up to the desired final water volume. **An approved dye must be added to treated seed to prevent inadvertent use for food, feed or oil purposes.** Falgro 2X LV is compatible with most commonly used seed treatments, standard dyes and binding agents. However, users should conduct the following physical compatibility test and a mixing compatibility test before mixing and applying product to large areas: 1) Add pesticides in the correct proportions. 2) Mix thoroughly and let stand for at least 15 minutes. 3) Observe for any signs of incompatibility e.g., heat, separation, gelling.

RICE SEEDLING POST-EMERGENCE TREATMENT

Application of Falgro 2X LV prior to permanent flooding promotes uniform and vigorous growth of semi-dwarfing varieties allowing earlier flooding with its associated agronomic benefits. When permanent flooding takes place before tiller development, make one

application of Falgro 2X LV at 2-3 g ai/acre at the 1-2 leaf stage. When flooding will be made following initial tilling, apply 1-3 g ai/acre at the 3 to 4 leaf (4th leaf showing) stage. Use higher rates when temperatures are likely to average 75°F or less during the 14 days after application. Either application will allow the establishment of a permanent flood 7-10 days earlier.

Apply Falgro 2X LV by fixed wing aircraft equipped with spray systems capable of producing a uniform medium to fine spray droplet pattern in not less than 10 gallons of water/acre. Low-pressure ground sprayers equipped with boom and flat fan nozzles applying 10-15 gallons of water/acre can also be used.

Ensure fields have been drained of floodwater before application and avoid spray drift onto other crops. Do not apply to crops suffering stress. Do not add surfactants, oils or any type of adjuvants to the spray tank. These treatments may cause the crop to become a lighter shade of green. This is a temporary effect and is caused by the increased growth rate.

HYBRID RICE FOR SEED PRODUCTION (All approved states except California)

Applications of Falgro 2X LV will promote main culm and tiller panicle extension, resulting in increased pollination and improved harvest efficiency. Make one to five applications of Falgro 2X LV at 20-100 g ai/acre applied at regular intervals during the heading period.

CONVERSION TABLE

Falgro 2X LV is a 6.18% active ingredient liquid. For each gram of active ingredient required use 0.5 fl. oz. of Falgro 2X LV.

Grams ai per acre	To Falgro 2X LV per acre	Grams ai per acre	To Falgro 2X LV per acre
0.5	0.25 fl. oz.	20.0	10 fl. oz.
1.0	0.5 fl. oz.	32.0	16 fl. oz.
5.0	2.5 fl. oz.	40.0	20 fl. oz.
10.0	5 fl. oz.	48.0	24 fl. oz.
16.0	8 fl. oz.		

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Keep containers tightly closed when not in use.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into the application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap.

Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip.

Repeat this procedure two more times, or puncture or dispose of in a sanitary landfill, or by incineration. Then offer for recycling, or reconditioning, if available, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY

Fine Agrochemicals Limited ("FINE") warrants that this Product conforms to the specifications on this label. To the extent consistent with applicable law, FINE makes no other warranties and disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for a particular purpose. No agent of FINE or any other person is authorized to make any representation or warranty beyond those contained herein.

It is impossible to eliminate all risks associated with this Product. Plant injury, lack of performance, or other unintended consequences may result because of factors such as abnormal weather conditions, use of the Product other than in strict accordance with this label's instructions, presence of other materials, the manner of application or other factors, all of which are beyond the control of FINE or the seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

To the extent consistent with applicable law: 1) FINE disclaims any liability whatsoever for special, incidental or consequential damages resulting from the handling or use of this Product and 2) FINE's liability under this label shall be limited to the amount of the purchase price or, at the election of FINE, the free replacement of the Product.

Manufactured by:
© Fine Agrochemicals Limited
Hill End House
Whittington
Worcester, WR5 2RQ
United Kingdom

Falgro is a registered trademark of Fine Agrochemicals Ltd.

SUBLABEL B: Ornamental Uses

FALGRO 2X LV

A plant growth regulator solution for use on turf and ornamental crops

Active Ingredient:

Gibberellic Acid (GA₃) 6.18%

Other Ingredients: 93.82%

Total: 100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

[See attached label booklet for first aid, precautionary, storage and disposal statements, and Directions for Use.]

FLAMMABLE – Keep away from heat and open flame

EPA Reg. No. 62097-32

EPA Est. No.

NET CONTENTS: 1 qt, ½ gal, 1 gal, 2.5 gal, 15 gal, 30 gal, 55 gal
Batch No.

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.

IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
	<ul style="list-style-type: none"> • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including general health concerns or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378, Monday through Friday, 8:00 AM to 12:00 PM Pacific Standard Time. In the event of a medical emergency, call your poison control center at 1-800-222-1222</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Causes moderate eye irritation. Harmful if swallowed, absorbed through the skin, or inhaled. Avoid contact with eyes, skin, or clothing and breathing spray mist. Wear goggles or safety glasses. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical-resistance selection chart.

Applicators and other handlers must wear:

- Protective eyewear
- Coveralls worn over short-sleeved shirt and short pants
- Shoes plus socks
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride or Viton

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE items separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of the gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses. Do not apply directly to water, to areas where surface water is

present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

PHYSICAL AND CHEMICAL HAZARDS

FLAMMABLE. Keep away from heat and open flame. FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your state or tribe, consult the state or tribal agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE) and restricted entry intervals. The requirements in this box only apply to uses of this product covered by the Worker Protection Standard.

Do not enter or allow entry into treated areas during the restricted entry interval level (REI) of 4 hours unless wearing appropriate PPE.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as, plants, soil or water is:

- Coveralls worn over short-sleeved shirt and short pants
- Shoes plus socks
- Protective eyewear
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride or Viton.

PRODUCT INFORMATION

- Use only as directed. The label should be read thoroughly and understood before making applications. For best results, trials should be conducted to determine desired rates and timings for optimum responses under local growing conditions and with specific varieties.
- The term 'grams of active ingredient per acre' is represented by 'gai/acre' throughout this product label.
- All parts of the plant or crop should be covered by the spray to obtain the desired result.
- Prepare solution concentrations by mixing the required amount of product with water

only in a clean empty spray tank.

- Any unused spray material at the end of the day must be disposed of following local, State or Federal law.
- When a range of rates is indicated use the concentration and spray volume recommended locally. Consult your local experimental station or distributor for spray program best suited to your conditions.
- Falgro 2X LV works best when using water of neutral or slightly acidic pH. Water should always have a pH of less than 8.5.
- Absorption of Falgro 2X LV into the plant is greatest under slow drying conditions. Night-time applications will be more effective when day-time conditions cause rapid drying. Falgro 2X LV should be re-applied if significant rain occurs within 2 hours of application
- Data concerning the compatibility of Falgro 2X LV is not available.

SPRAYING GUIDE FOR ORNAMENTALS

Azalea

Applications of Falgro 2X LV can be used to help promote flowering by acting as a partial substitute for cold treatment. After plants have been exposed for 3-4 weeks of chilling and are at Stage 5 of floral development, three applications of 250-500 ppm each may be applied on a weekly schedule. Do not apply first application before Stage 5 (i.e., style elongated and open) as doing so may result in abnormal flower development. Do not apply to azaleas once flower buds begin to show color. Thorough and even coverage of foliage is necessary or plants may not flower evenly. For certain cultivars (Gloria, Prize and Redwing), a single application of 1000 ppm applied after 4 weeks of chilling has been shown to be effective to break dormancy. For best results, conduct trials to determine desired rates and timings for optimum responses under local growing conditions.

Application of Falgro 2X LV to the foliage can inhibit initiation of flower budding during vegetative growth. Apply 100-750 ppm to foliage 2-3 weeks after each pinch. One to two sequential applications can be made on a weekly basis following the initial application. Do not make more than 3 applications. For best results, conduct trials to determine desired rates and timings for optimum responses under local growing conditions.

Caladium sp.

Application will influence flowering. Soaking bulbs in solutions of between 250 ppm and 1000 ppm will produce a favorable result. For best results, conduct trials to determine desired rates and timings for optimum responses under local growing conditions and specific varieties.

Calla Lily

Application of Falgro 2X LV by soak, spray or field drench increases flowering, controls height, and shortens days to flowering. As a soak, use a 100-500 ppm solution. Spray applications are generally made at concentrations between 50 and 100 ppm to plants to stimulate flowering. Field drench is made over the planted bulbs at 100-500 ppm. For

best results, conduct trials to determine desired rates and timings for optimum responses under local growing conditions.

Camellia

Application of Falgro 2X LV can be used as a substitute when there has been a lack of required chilling for Camellia and has been shown to increase bloom size. Prepare a solution of Falgro 2X LV by mixing 0.5 fl. oz. of Falgro 2X LV with each 1.5 fl. oz. of water needed to make up the final mix size. The vegetative bud that is immediately adjacent to or below the floral bud will need to be removed before applying the solution. Then apply one drop of the prepared solution to the vegetative bud scar. To prevent run-off of the solution drop, a deposition aid may be added to the solution when mixing. Follow the instructions for mixing and use rates on the label for the deposition aid. For best results, conduct trials to determine desired rates and timings for optimum responses under local growing conditions and specific varieties.

Cyclamen sp.

Application to the foliage will favorably alter the vegetative and reproductive development of the plants. Foliar applications of between 25 ppm and 250 ppm will affect the vegetative and reproductive growth of the plant, thus reducing the days to flowering and enhancing production. For best results, conduct trials to determine desired rates and timings for optimum responses under local growing conditions.

Dieffenbachia sp.

Application to foliage will stimulate flowering for breeding and other purposes. Applications between 250 ppm and 1000 ppm stimulate the production of flowers, increasing the flexibility of breeding programs and frequency of flowers to be used to produce hybridized variants. For best results, conduct trials to determine desired rates and timings for optimum responses under local growing conditions.

Geranium (Pelargonium sp.)

Application of Falgro 2X LV to rooted cuttings will stimulate root growth and development when added directly to hydroponic nutrient solution. Applications of between 0.5 ppm and 50 ppm are recommended to increase productivity and produce other desirable plant characteristics. For best results, conduct trials to determine desired rates and timings for optimum responses under local growing conditions.

Poinsettia

Application to poinsettia is used to influence flowering. Foliar applications to plants of between 10 ppm and 50 ppm can be used. For best results, conduct trials to determine desired rates and timings for optimum responses under local growing conditions and specific varieties.

Pompom Chrysanthemum

Application of Falgro 2X LV elongates the peduncles. Make one application of 0.5-1 gal in 12 gallons of water 28-35 days after initiation of short day program. Apply this solution over 1000 sq. ft. of bed using overhead nozzles directed at the flower buds.

Apply at the correct timing and do not overdose as this can cause long, spindly, weak stems.

***Spathiphyllum* sp.**

Application to foliage will stimulate flowering for breeding and other purposes. Applications between 150 ppm and 250 ppm stimulate the production of flowers increasing the flexibility of production and the ability to sell plants in flower. For best results, conduct trials to determine desired rates and timings for optimum responses under local growing conditions.

Statice (*Limonium* sp.)

Application of Falgro 2X LV promotes early flowering and increases flower yield. Make one application of 40-50 g ai in 25 gallons of water when plants are more than 10 inches in diameter (approx. 90-110 days after planting). Apply 10 ml of spray as a drench to each plant.

Do not exceed the recommended dose or repeat treatment. Initiation of flowering is influenced by extended photo-period, nutrition and reduced night temperature.

Treatment with Falgro 2X LV lessens the need for reduced night temperature and long photo-period.

Tropical Foliage

For application to plants such as *Aglaonema commutatum*. Apply Falgro 2X LV to induce uniform flowering. Recommended dosages range from 100-400 ppm applied as a spray solution to runoff. Generally, treatment is used to induce flowering so breeding and cross pollinations can be carried out. For best results, conduct trials to determine desired rates and timings for optimum responses under local growing conditions.

SPRAYING GUIDE FOR TURFGRASS

Bermuda Golf Turf

Application of Falgro 2X LV can be used to initiate or maintain growth and prevent color change during periods of cold stress. Make one application of 10 g ai weekly or 25 g ai in 25-100 gallons of water/acre every 2 weeks. Do not exceed these recommended doses or reduce the time interval between applications. Do not apply during extended warm periods if night temperatures exceed 65°F. Continue good ground keeping practice and stop treatment if thinning of the grass occurs. Do not use on dormant turf.

For warm weather application, apply 1-3 g ai in 25-100 gallons of water/acre weekly.

CONVERSION TABLES

GRAMS OF AI PER ACRE

Falgro 2X LV is a 6.18% active ingredient liquid. For each gram of active ingredient required, use 0.5 fl. oz. of Falgro 2X LV.

Grams ai per acre	Falgro 2X LV per acre
0.5	0.25 fl. oz.
1.0	0.5 fl. oz.
5.0	2.5 fl. oz.
10.0	5 fl. oz.
16.0	8 fl. oz.
20.0	10 fl. oz.
32.0	16 fl. oz.
40.0	20 fl. oz.
48.0	24 fl. oz.

PARTS PER MILLION

Desired ppm	Cups of Falgro 2X LV per 100 gals	Quarts of Falgro 2X LV per 100 gals
50	1.25	-
100	2.5	0.6
200	5	1.3
300	7.5	1.9
400	10	2.5
500	12.5	3.1
1,000	25	6.3

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Keep containers tightly closed when not in use.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY

Fine Agrochemicals Limited ("FINE") warrants that this Product conforms to the specifications on this label. To the extent consistent with applicable law, FINE makes no other warranties and disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for a particular purpose. No agent of FINE or any other person is authorized to make any representation or warranty beyond those contained herein.

It is impossible to eliminate all risks associated with this Product. Plant injury, lack of performance, or other unintended consequences may result because of factors such as

abnormal weather conditions, use of the Product other than in strict accordance with this label's instructions, presence of other materials, the manner of application or other factors, all of which are beyond the control of FINE or the seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

To the extent consistent with applicable law: 1) FINE disclaims any liability whatsoever for special, incidental or consequential damages resulting from the handling or use of this Product and 2) FINE's liability under this label shall be limited to the amount of the purchase price or, at the election of FINE, the free replacement of the Product.

Manufactured by:
© Fine Agrochemicals Limited
Hill End House
Whittington
Worcester, WR5 2RQ
United Kingdom

Florgib is a registered trademark of Fine Agrochemicals Ltd.