



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Biopesticides and Pollution Prevention Division (7511C)
 1200 Pennsylvania Avenue NW
 Washington, DC 20460

EPA Reg. Number:

62097-33

Date of Issuance:

NOV 10 2014

Term of Issuance:

Unconditional

Name of Pesticide Product:

Exilis® 5 XL

NOTICE OF PESTICIDE:

Registration

Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Fine Agrochemicals Ltd.
 c/o SciReg, Inc.
 12733 Director's Loop
 Woodbridge, VA 22192

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA). Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on her motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This registration does not eliminate the need for continual reassessment of the pesticide. If the EPA determines at any time that additional data are required to maintain in effect an existing registration, the Agency will require submission of such data under section 3(c)(2)(B) of FIFRA.

This product is unconditionally registered in accordance with FIFRA Sec. 3(c)(5) and is subject to the following terms:

1. Submit and/or cite all data required for registration of your product under FIFRA section 3(c)(5) and section 4 when the Agency requires all registrants of similar products to submit such data.
2. A one year study is required to satisfy the Storage Stability and Corrosion Characteristics requirements (Guidelines: OCSPP 830.6317 and 830.6320). You have 18 months from the date of registration to provide these data.
3. Revise the EPA Registration Number to read, "EPA Reg. No. 62097-33."
4. Submit three (3) copies of the revised final printed labeling before you release the product for shipment.

A stamped copy of the label and an A-79 Enclosure are enclosed for your records.

Signature of Approving Official:

Linda A. Hollis, Biochemical Pesticide Branch Chief
 Biopesticides and Pollution Prevention Division (7511P)

Date:

11-10-14

2/15

Master Label

Exilis® 5 XL

Sub-label A: Agricultural/Commercial Use

Sub-label B: Ornamental Use

Active Ingredient: N-(phenylmethyl)-1H-purine-6-amine	9.51%
Other Ingredients:	<u>90.49%</u>
Total:	100.00%

KEEP OUT OF REACH OF CHILDREN

EPA Reg. No. 62097-

EPA Est. No.

ACCEPTED

NOV 10 2014

Under the Federal Insecticide, Fungicide,
 and Rodenticide Act, as amended, for
 the pesticide registered under
 EPA Reg. No. 62097-33

3/
15

Sub-label A

Exilis® 5 XL

For Agricultural/Commercial Use

ACCEPTED

4/
15

Exilis® 5 XL

Plant growth regulator solution for use on apples and pears

Active Ingredient: N-(phenylmethyl)-1H-purine-6-amine	9.51%
Other Ingredients:	90.49%
Total:	100.00%

KEEP OUT OF REACH OF CHILDREN
[See back panel for Precautionary Statements]

EPA Reg. No. 62097-

EPA Est. No.

NET CONTENTS:

Batch:

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear appropriate protective clothing. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical-resistance selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride or viton.
- Socks and shoes.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

[For 15 gal, 30 gal, and 55 gal containers] Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

PHYSICAL OR CHEMICAL HAZARDS

FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- * Coveralls
- * Shoes plus socks
- * Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride or viton.

PRODUCT INFORMATION

- EXILIS® 5 XL is a plant growth regulator for use on apples and pears.
- EXILIS® 5 XL is a post-bloom thinning agent.
- EXILIS® 5 XL may increase fruit weight, but this is dependent on fruit set and the physiological condition of the trees.
- Apply EXILIS® 5 XL using conventional spray application machinery or as otherwise directed in this label. Do not apply EXILIS® 5 XL using irrigation equipment.

- The effectiveness of EXILIS® 5 XL depends on the variety of target apple and pear trees.
- The thinning effectiveness of EXILIS® 5 XL is increased by:
 - High temperatures before and after application (70-75°F (21-24°C)).
 - Disease and physiological stress of target trees. Apply reduced rates to low vigor trees, or trees under stress from drought, winter injury, etc.
- The effectiveness of EXILIS® 5 XL is reduced by:
 - Low temperatures before and after application. Do not apply EXILIS® 5 XL if the temperature is below 60°F (16°C).
 - Rainfall immediately after application.
 - pH above 8. EXILIS® 5 XL performs best when the pH is 5-6.5. A buffer or pH enhancer may be used for this purpose.
- Direct 80% of the spray to the top 2/3rds of the tree structure.
- Use a properly calibrated sprayer to ensure uniform and complete coverage of all foliage and developing fruit without run-off. Spray volume should be adjusted for tree size and density.
- EXILIS® 5 XL works best if climatic conditions allow an extended drying period. Consider application in the morning or evening when temperatures are lower and humidity tends to be higher than during the day.
- Consult your farm advisor or extension specialist for advice about tank-mixing EXILIS® 5 XL with other compounds.
- DO NOT EXCEED 61.5 fl oz of product (182 grams of active ingredient) PER ACRE PER SEASON.
- DO NOT APPLY CLOSER THAN 86 DAYS BEFORE HARVEST.

SPRAYING GUIDE FOR FRUIT THINNING

When used to thin fruit, EXILIS® 5 XL is normally applied in a single spray program; if a second application is needed allow 7-10 days after the first to observe the effect and spray before the diameter of king bloom fruit exceeds 20mm.

Make 1-2 applications of 75-200 ppm when king fruitlets are 5-10 mm in diameter (see EXILIS® 5 XL Dilution Guide). The effectiveness of EXILIS® 5 XL depends on the variety of target apple and pear trees and weather conditions. Refer to your farm advisor or extension specialist for advice about rate of application. DO NOT EXCEED 61.5 fl oz of product (182 grams of active ingredient) PER ACRE PER SEASON.

SPRAYING GUIDE FOR FRUIT SIZING & RETURN BLOOM

Spray 2-4 applications of 10-50 ppm at petal fall and following at 3-10 day intervals. Generally use higher rates at longer intervals and lower number of applications. Refer to your farm advisor or extension specialist for advice about rate of application. DO NOT EXCEED 61.5 fl oz of product (182 grams of active ingredient) PER ACRE PER SEASON.

Applications for fruit sizing may cause thinning in some varieties or conditions of application.

7/15

EXILIS® 5 XL DILUTION GUIDE

Parts Per Million	10 ppm	25 ppm	50 ppm	75 ppm	100 ppm	125 ppm	150 ppm	175 ppm	200 ppm
Fluid Ounces of Product Per 100 Gallons of Water	1.2	3.2	6.4	9.6	12.8	16.0	19.2	22.4	25.6

[EXILIS® 5 XL contains 189.35 grams active ingredient per 64 fluid ounces (one-half gallon).]

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage: Keep container tightly closed when not in use. Store below 75°F.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

{for container sizes of 5 gallons and less} Container Handling: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, if available, or puncture or dispose of in a sanitary landfill, or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

{for container sizes greater than 5 gallons} Container Handling: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, if available, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY

Fine Agrochemicals Limited ("FINE") warrants that this Product conforms to the specifications on this label. To the extent consistent with applicable law, FINE makes no other warranties and disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for a particular purpose. No agent of FINE or any other person is authorized to make any representation or warranty beyond those contained herein.

It is impossible to eliminate all risks associated with this Product. Plant injury, lack of performance, or other unintended consequences may result because of factors such as

abnormal weather conditions, use of the Product other than in strict accordance with this label's instructions, presence of other materials, the manner of application or other factors, all of which are beyond the control of FINE or the seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

To the extent consistent with applicable law: 1) FINE disclaims any liability whatsoever for special, incidental or consequential damages resulting from the handling or use of this Product and 2) FINE's liability under this label shall be limited to the amount of the purchase price or, at the election of FINE, the free replacement of the Product.

© Fine Agrochemicals Limited
Hill End House
Whittington
Worcester, WR5 2RQ
United Kingdom

EXILIS® is a registered trademark of Fine Agrochemicals, Ltd.

Sub-label B

Exilis® 5 XL

For Ornamental Use

10/
15

EXILIS® 5 XL

Plant growth regulator solution for use on ornamentals

Active Ingredient: N-(phenylmethyl)-1H-purine-6-amine	9.51%
Other Ingredients:	90.49%
Total:	100.00%

KEEP OUT OF REACH OF CHILDREN
[See back panel for Precautionary Statements]

EPA Reg. No. 62097-

EPA Est. No.

NET CONTENTS:

Batch:

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear appropriate protective clothing. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical-resistance selection chart. Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride or viton.
- Socks and shoes.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

11/15

[For 15 gal, 30 gal, and 55 gal containers] Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

PHYSICAL OR CHEMICAL HAZARDS

FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE) and restricted entry intervals. The requirements in this box only apply to uses of this product covered by the Worker Protection Standard.

Do not enter or allow entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride or viton

GENERAL INFORMATION

Do not apply this product through any type of irrigation system.

FOR USE IN: COMMERCIAL GREENHOUSES, GLASSHOUSES (WHERE PLANTS ARE GROWN IN CONTAINERS), AND ON ACTIVELY GROWING STOCK PLANTS

EXILIS® 5 XL increases lateral or basal branching and promotes flowering in certain ornamental crops. In addition, applications of EXILIS® 5 XL reduce the overall height of the plant resulting in more compact and marketable plants.

12/
15

Plant response to EXILIS® 5 XL is strongly influenced by cultural and environmental variables. Growing media, water/fertilizer management, temperature, light, greenhouse composition, and other cultural practices may impact plant response. Plant size, container size and cultivar or variety can also influence treatment response. Although multiple cultivars have been tested for sensitivity to EXILIS® 5 XL and have demonstrated high levels of tolerance, it is impossible to insure an acceptable response in all cultivars. **For first time users of EXILIS® 5 XL, always conduct trials on a limited number of plants to determine crop response.**

APPLICATION INFORMATION

Apply EXILIS® 5 XL as a foliar spray using standard foliar application spray equipment. Make sure that sufficient volumes are used to thoroughly wet foliage. Spray uniformity is equally important. **Uniformly apply 1-2 quarts of finished spray solution to 100 sq. ft. of area.** A high quality wetting agent or spray adjuvant, approved for use on your crop, may be added to spray solutions according to the manufacturer's use instructions.

DIRECTIONS FOR USE

Christmas Cactus (*Schlumbergera* spp.):

EXILIS® 5 XL can be applied as a foliar spray to: 1) promote vegetative branching; or 2) increase the number of flower buds under reproductive conditions. Results vary with cultivar; therefore, trial a small group first.

To increase branching under vegetative conditions: Apply EXILIS® 5 XL at 100 ppm as a uniform foliar spray after planting when new vegetative growth begins.

To increase the number of flower buds under reproductive conditions: Apply EXILIS® 5 XL at 100 to 200 ppm as a uniform foliar spray after the start of short days following leveling, or when flower buds become visible. Below are specific guidelines based on lighted or natural season grown plants.

For floral initiation under short day conditions (i.e., lighting/black clothing): Wait 5 to 10 days after the start of short days to level by removing immature phylloclades. Make a single application of EXILIS® 5 XL immediately following or 1 day after leveling.

For floral initiation under natural environmental conditions: Make a single application of EXILIS® 5 XL after flower buds first become visible (i.e., pinpoint bud stage). First visible bud varies by cultivar. Avoid early applications which will result in phylloclade promotion and delayed flowering.

Plantain Lily (*Hosta* spp.):

Use of EXILIS® 5 XL on *Hosta* can promote lateral growth of finished plants by inducing the outgrowth of axillary and rhizomic buds. EXILIS® 5 XL can also be used to increase offsets during propagation.

For the promotion of lateral growth on finished plants: Apply EXILIS® 5 XL at rates ranging from 1000 – 3000 ppm in a uniform spray volume. EXILIS® 5 XL is most effective

13/
15

when plants are fully established prior to application (i.e., at least 3 to 4 weeks after potting), when there is evidence of surface root development, but before flower initiation.

For increased production of offsets (propagation): Apply EXILIS® 5 XL at rates ranging from 1000 – 3000 ppm in a uniform spray volume to fully established, actively growing stock plants. Repeat the application at 30-day intervals during the growing season. Offsets may be harvested at any time.

Treatment effects may vary by *Hosta* cultivar, and may respond differently to a given rate of EXILIS® 5 XL. Multiple applications at 30-day intervals using lower rates may be more effective than a single application at higher rates. Conduct trials on a small number of plants under actual use conditions to establish the proper use rates and timings.

Purple Coneflower (*Echinacea* spp.):

Foliar applications of EXILIS® 5 XL have been shown to increase the number of branches when applied at rates of 300 to 900 ppm during active growth.

Apply EXILIS® 5 XL as a foliar spray after plant establishment and resumption of growth (i.e., approximately 2 weeks after potting). Apply in a uniform spray volume of 2 quarts per 100 square feet of area. Application timing and rate may vary with cultivar. For first time users of EXILIS® 5 XL, determine the appropriate rate and application timing by conducting trials on a small number of plants on specific cultivars and under typical environmental growth conditions.

APPLICATION INSTRUCTIONS FOR USE ON CONTAINERIZED ANNUAL AND PERENNIAL FLOWERING AND FOLIAGE CROPS AND TROPICAL PLANTS

- Apply EXILIS® 5 XL as a foliar spray when plants are fully established during active growth i.e., at least 3 to 4 weeks after potting.
- Apply EXILIS® 5 XL at rates ranging from 50 to 500 ppm, depending on plant species and variety.
- Apply in a uniform spray volume of 2 quarts per 100 square feet of area. Cultivars within the same species may respond differently to a given rate of EXILIS® 5 XL. For first time users of EXILIS® 5 XL, **always treat a limited number of plants starting with 50 ppm on annuals and 100 ppm on perennials and tropical plants**; observe for plant responses and make rate adjustments accordingly.
- Multiple applications of EXILIS® 5 XL at 7-10 day spray intervals may be necessary to achieve optimum results, i.e. increased lateral growth.
- EXILIS® 5 XL may cause significant injury to *Exacum*. DO NOT use EXILIS® 5 XL on this species.
- EXILIS® 5 XL has been shown to cause a significant delay in flowering on *Gaillardia*. DO NOT apply EXILIS® 5 XL to *Gaillardia* more than once during the production cycle.

14/
15

EXILIS® 5 XL DILUTION GUIDE:

Desired Rate (ppm)	50	100	200	300	400	500	900	1000	2000	3000
fl oz EXILIS® 5 XL per gallon of water	0.064	0.128	0.256	0.384	0.512	0.640	1.15	1.28	2.56	3.84
ml EXILIS® 5 XL per gallon of water	1.9	3.8	7.6	11.4	15.2	19.0	34.2	38.0	76.0	114.0
ml EXILIS® 5 XL per liter of water	0.50	1.0	2.0	3.0	4.0	5.0	9.0	10.0	20.0	30.0

[EXILIS® 5 XL contains 189.35 grams active ingredient per 64 fluid ounces (one-half gallon).]

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage: Keep container tightly closed when not in use. Store below 75°F.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

{for container sizes of 5 gallons and less} Container Handling: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, if available, or puncture or dispose of in a sanitary landfill, or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

{for container sizes greater than 5 gallons} Container Handling: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, if available, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

15/
15

WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY

Fine Agrochemicals Limited ("FINE") warrants that this Product conforms to the specifications on this label. To the extent consistent with applicable law, FINE makes no other warranties and disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for a particular purpose. No agent of FINE or any other person is authorized to make any representation or warranty beyond those contained herein.

It is impossible to eliminate all risks associated with this Product. Plant injury, lack of performance, or other unintended consequences may result because of factors such as use of the Product other than in strict accordance with this label's instructions, presence of other materials, the manner of application or other factors, all of which are beyond the control of FINE or the seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

To the extent consistent with applicable law: 1) FINE disclaims any liability whatsoever for special, incidental or consequential damages resulting from the handling or use of this Product and 2) FINE's liability under this label shall be limited to the amount of the purchase price or, at the election of FINE, the free replacement of the Product.

© Fine Agrochemicals Limited
Hill End House
Whittington
Worcester, WR5 2RQ
United Kingdom

EXILIS® is a registered trademark of Fine Agrochemicals, Ltd.