



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

June 6, 2016

Frederick T. Smith
Senior Regulatory Specialist
Fine Agrochemicals Ltd.
c/o SciReg, Inc.
12733 Director's Loop
Woodbridge, VA 22192

Subject: Label Amendment – Adding me-too uses of strawberries, turf, watercress, and pears
Product Name: Kudos 27.5 WDG
EPA Registration Number: 62097-41
Application Date: 03/14/2016
Decision Number: 515168

Dear Mr. Smith:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Page 2 of 2
EPA Reg. No. 62097-41
Decision No. 515168

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Lisa Pahl by phone at (703) 347-0459, or via email at pahl.lisa@epa.gov.

Sincerely,

A handwritten signature in cursive script that reads "Heather Garvie".

Heather Garvie, Product Manager 24
Fungicide and Herbicide
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

MASTER LABEL

KUDOS® 27.5 WDG

Plant growth regulator for use on apples, grass grown for seed, peanuts, d'Anjou pears, strawberries, sweet cherries, turf, and watercress.

Sublabel A: Agricultural Uses


Sublabel B: Turf Uses

Active Ingredient: Prohexadione-calcium

[calcium 3-oxido-5-oxo-4-propionylcyclohex-3-enecarboxylate].....27.5%

Other Ingredients:72.5%

Total:100.0%



**KEEP OUT OF REACH OF CHILDREN
CAUTION**

[See inside panel for First Aid Statements]

EPA Reg. No. 62097-41

EPA Est. No.

Net Contents:

© Fine Agrochemicals Limited, Hill End House, Whittington, Worcester, WR5 2RQ,
United Kingdom

Kudos® 27.5 WDG is a registered trademark of Fine Agrochemicals, Ltd.

SUBLABEL A: Agricultural Uses

KUDOS® 27.5 WDG

Plant growth regulator for use on apples, grass grown for seed, peanuts, d'Anjou pears, strawberries, sweet cherries, turf, and watercress.

Active Ingredient: Prohexadione-calcium

[calcium 3-oxido-5-oxo-4-propionylcyclohex-3-enecarboxylate].....27.5%

Other Ingredients:72.5%

Total:100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

[See inside panel for First Aid Statements]

EPA Reg. No. 62097-41

EPA Est. No.

Net Contents:

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Poison Control Hotline at 1-800-222-1222 for emergency medical treatment information 24 hours a day, seven days a week.	

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE items separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE) and restricted entry intervals. The requirements in this box only apply to uses of this product covered by the Worker Protection Standard.

Do not enter or allow entry into treated areas during the restricted entry interval level (REI) of 12 hours unless wearing appropriate PPE.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as, plants, soil or water is:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical resistant gloves made of any waterproof material

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your state or tribe, consult the state or tribal agency responsible for pesticide regulation.

Kudos 27.5 WDG is a plant growth regulator which inhibits the biosynthesis of gibberellins. Inhibition of gibberellin reduces vegetative growth.

- Thorough spray coverage of plant foliage is necessary for good uptake.
- The performance of Kudos 27.5 WDG can be affected by many factors including: crop growth stage, environmental conditions and crop vigor due to moisture availability, fertility level, etc.
- Correct timing of application is critical. Apply Kudos 27.5 WDG to actively growing trees at rates and stages of growth listed in this label.
- Add one pound of a spray grade ammonium sulfate (AMS) for every pound of Kudos 27.5 WDG.
- Use a standard spray adjuvant, preferably a non-ionic surfactant, to improve leaf coverage and performance consistency. Follow the manufacturer's rate recommendations.
- Rainfall within 1 hour of application on grass grown for seed and within 8 hours for apples, peanuts, and sweet cherries may reduce the efficacy of Kudos 27.5 WDG.
- Clean spray equipment thoroughly using a strong detergent or commercial sprayer cleaner according to the manufacturer's directions before and after applying Kudos 27.5 WDG.

Restrictions

- **DO NOT** apply to crops under stress due to lack of moisture, hail damage, flooding, mechanical injury, or injury caused by other prior products, i.e., phytotoxicity.
- **DO NOT** tank mix Kudos 27.5 WDG with calcium (including gypsum) or boron sprays, as a reduction in growth control may result.
- **DO NOT** apply this product through any type of irrigation system.
- This product cannot be used to formulate or reformulate any other pesticide product.

General Tank Mixing Information:

Testing has shown that Kudos 27.5 WDG when used in combination with AMS and a wide range of adjuvants does not result in phytotoxicity. Kudos 27.5 WDG has shown to be compatible with many commonly used fungicides and insecticides. However, not all crop varieties and cultivars have been tested with possible tank-mix combinations. Since local conditions can influence crop tolerance, test any tank-mix combination on a small portion of the crop to be treated to ensure crop safety. Read and follow the applicable Directions For Use on all products involved in tank-mixing. Always refer to the most restrictive labeling.

Mixing Order:

1. Fill spray tank half full with clean water and begin agitation.
2. Place any product contained in a water-soluble PVA bag into the tank and make sure the bag(s) have fully dissolved before continuing.
3. Add Ammonium Sulfate (AMS).
4. Add Kudos 27.5 WDG.
5. Add any water dispersible formulations, i.e., Water Dispersible Granules, Wettable Powders, Suspension Concentrates, etc.
6. Add adjuvants.
7. Add any water soluble products, i.e., Soluble Powders, Soluble Liquids, Emulsifiable Concentrates, etc.
8. Add remaining quantity of water and continue agitation through application.

Maintain constant agitation during application.

APPLICATION INSTRUCTIONS

APPLES:

Kudos 27.5 WDG can be used as a production management tool on apples to suppress vegetative growth, i.e., shoot growth reduction, resulting in a balance between canopy development and fruit production. Growth suppression with Kudos 27.5 WDG generally lasts for 2-5 weeks per application. Kudos 27.5 WDG does not affect vegetative growth the year following application.

Benefits resulting from the use of Kudos 27.5 WDG include:

- Reduction in vegetative shoot growth
- Lessens the need or frequency for summer and/or dormant pruning
- Improved fruit color in red or bi-color apple varieties due to better light penetration into the tree canopy
- Reduced incidence and severity of fire blight of shoots

During years when both Kudos 27.5 WDG and products containing gibberellins, e.g., Novagib[®] 10L are used, reductions in efficacy may occur in the Kudos 27.5 WDG and/or gibberellin treatments.

The use of Kudos 27.5 WDG may result in an increase in fruit retention. Therefore, thinning programs may need to be adjusted when using Kudos 27.5 WDG.

When used as recommended, Kudos 27.5 WDG will reduce the incidence and severity of fire blight infection of shoots and leaves, i.e., shoot blight, by reducing vegetative growth, thus decreasing host susceptibility. This decrease in susceptibility will not become effective until approximately 10 days after application. Kudos 27.5 WDG does not have direct antibiotic activity against the fire blight bacteria (*Erwinia amylovora*) and is not effective for the suppression of blossom blight. For maximum reduction in fire blight susceptibility, apply Kudos 27.5 WDG at least 10 days before the occurrence of weather conditions favorable for shoot and leaf infections, and used as one component of a comprehensive control strategy against fire blight.

Application Timings: For vegetative growth control, make the first application of Kudos 27.5 WDG when shoots have 1-3" of new growth. For best results, make subsequent applications at 1-4 week intervals, before or immediately following signs of shoot regrowth.

Number of Applications: The number of applications will vary depending on the timing of the first application, tree vigor, fruit load, pruning, variety, rootstock, and/or the management history of the orchard. For apple orchards in locations with a long growing season or on high vigor trees and/or trees with light fruit load, 3 to 5 applications per year may be required.

Tree Vigor: Adjust the Kudos 27.5 WDG rate according to the vegetative vigor of the trees (see Table 1). Vegetative vigor can be influenced by many factors, including variety, rootstock, fruit load, cultural management and location.

Tree Size: Calculate the rate per acre of Kudos 27.5 WDG based on tree size. Base the application rate on the volume of water needed to spray the trees to drip i.e. dilute spray or Tree Row Volume (TRV). Consult your local extension agent or consultant for a recommendation on spray volume.

Application Rates: Recommended rates will be based on desired treatment effect, i.e., vegetative growth control, shoot blight suppression, etc. For vegetative growth control, base Kudos 27.5 WDG rates on level of tree vigor and/or length of growing season. Refer to Table 1 for recommended rates per application.

For consistent performance on apples, add one pound of a spray grade ammonium sulfate (AMS) for every pound of Kudos 27.5 WDG. Use a standard spray adjuvant, preferably a non-ionic surfactant, to improve leaf coverage and performance consistency. Follow the manufacturer's rate recommendations.

Table 1. Recommended Application Rates on Apples

Desired Treatment Effect	Rate per Acre¹	Application Directions
Growth Reduction on Medium to High Vigor Trees	18-36 oz (5-9.9 g ai)	<ul style="list-style-type: none"> • Apply at 1-3" of new shoot growth. • Make subsequent applications at 1-4 week intervals and before or immediately after shoot regrowth occurs.
Growth Reduction on Low Vigor Trees	9-24 oz (2.5-6.6 g ai)	<ul style="list-style-type: none"> • Apply at 1-3" of new shoot growth. • Make subsequent applications at 1-4 week intervals and before or immediately after shoot regrowth occurs.
Growth Reduction During Long Growing Season	9-24 oz (2.5-6.6 g ai)	<ul style="list-style-type: none"> • Apply at 1-3" of new shoot growth. • Make a second and third application at 7-14 day intervals. • Make subsequent applications as needed at 10-14 day intervals.
To Decrease June Drop on Trees With Light Bloom	30-36 oz (8.3-9.9 g ai)	<ul style="list-style-type: none"> • Apply at 1-3" of new shoot growth.
To Shape Tree Canopy	See footnote 2	<ul style="list-style-type: none"> • Apply at 1-3" of new shoot growth. • Direct the spray to the portion of the tree where growth control is desired.
To Reduce Fire Blight Infections of Shoot by Decreasing Vegetative Growth	18-36 oz (5-9.9 g ai)	<ul style="list-style-type: none"> • Apply at 1-3" of new shoot growth. • Make a second application if new shoot growth occurs.

¹ Based on 300 gallons of dilute spray per acre.

²Apply 6-12 oz of Kudos 27.5 WDG (1.7-3.3 g ai) per 100 gallons of dilute spray until run-off.

Ground Application: Apply Kudos 27.5 WDG to actively growing trees using calibrated ground equipment. Kudos 27.5 WDG may be applied using either dilute or concentrate spray equipment as long as sufficient spray coverage is achieved. Spray should be directed to the portion of the tree where growth control is desired. To achieve good coverage, apply Kudos 27.5 WDG in a sufficient spray volume per acre utilizing proper spray pressure, nozzles, nozzle spacing and tractor speed.

Aerial Application (all approved states except California): Apply Kudos 27.5 WDG in a minimum of 10 gallons of spray solution per broadcast acre. Aerial applications generally only provide spray coverage in the top part of the tree canopy. Vegetative growth control will be limited to those areas of the canopy that receive spray coverage.

Special Directions for Vegetative Growth Control on Apples Grown in Idaho, Oregon and Washington: Apply Kudos 27.5 WDG to actively growing trees according to the tree size, rates and application timings listed in Table 2. It is important to take into consideration the size and vigor of the trees when determining the spray volume and application frequency, timing and rate required to achieve vegetative growth control. Spray volumes are based on the amount of solution required to thoroughly wet the tree foliage to the point of runoff. Consult your local extension agent or consultant for a recommendation on spray volume.

Table 2. Recommended Application Rates for Vegetative Growth Control of Apples in Idaho, Oregon or Washington.

Apple Tree Size	Rate per Acre ¹	Application Directions
Small Trees (8-10 feet tall on dwarf rootstocks)	6-12 oz (1.7-3.3 g ai)	Apply at 1-3" of new shoot growth. Make subsequent applications at 1-4 week intervals when shoots show signs of regrowth. High vigor trees may require more frequent applications through the growing season.
Medium Trees (10-14 feet tall on semi-dwarf rootstocks)	6-18 oz (1.7-5 g ai)	
Large Trees (greater than 14 feet tall on standard non-dwarf rootstocks)	18-24 oz (5-6.6 g ai)	

¹Spray volumes must be a minimum of 100 gallons per acre and increase as necessary to achieve thorough canopy coverage.

Limitations on Apples:

- Kudos 27.5 WDG is rainfast 8 hours after application.
- The active ingredient in Kudos 27.5 WDG has been shown to increase fruit cracking on apple varieties such as Empire and Stayman, which are known to be prone to cracking.

Restrictions on Apples:

- **DO NOT** apply more than a total of 99 oz of Kudos 27.5 WDG (27.2 g ai) per acre, per year.
- **DO NOT** apply more than a total of 48 oz of Kudos 27.5 WDG (13.2 g ai) within any 21-day interval.
- **DO NOT** apply within 45 days before harvest.
- **DO NOT** apply to crops that show injury (leaf phytotoxicity) produced by any other prior pesticide applications because this injury can be enhanced or prolonged.
- Restricted Entry Interval (REI): 12 hours.

GRASS GROWN FOR SEED (all approved states except California):

Kudos 27.5 WDG can be used as a production management tool on grass grown for seed production to suppress vegetative growth i.e. shorten internode length, resulting in a lower potential for lodging. Reducing lodging can result in improved pollination, increased seed set and improved yields. Kudos 27.5 WDG must be absorbed by foliage to be effective. Always use sufficient spray volume to thoroughly wet the leaves without runoff. The growth regulator effects of Kudos 27.5 WDG do not occur by soil uptake. Kudos 27.5 WDG does not affect vegetative growth the following year.

For consistent performance on grass grown for seed, add one pound of a spray grade ammonium sulfate (AMS) for every pound of Kudos 27.5 WDG. Use a standard spray adjuvant, preferably a non-ionic surfactant, to improve leaf coverage and performance consistency. Follow the manufacturer’s rate recommendations.

Refer to Table 3 for application rates and timings of Kudos 27.5 WDG to reduce vegetative growth in grass grown for seed.

Table 3. Application Rates and Timings for Vegetative Growth Control in Grass Grown for Seed.

Application Timing	Application Rate per Acre
Single application: <ul style="list-style-type: none">• Apply from flag leaf emergence up to early heading growth stage.	14-29 oz (3.9-8 g ai)
Split applications: <ul style="list-style-type: none">• Apply from flag leaf emergence up to early heading growth stage.• Make a second application 7-10 days later when new growth occurs.	7-14 oz (1.9-3.9 g ai)

Ground Application: Use a minimum of 10 gallons of spray solution per broadcast acre.

Aerial Application: Use a minimum of 10 gallons of spray solution per acre.

Suppression of Annual Bluegrass in Washington, Oregon, Idaho and Utah:

Kudos 27.5 WDG can be used to for the suppression of annual bluegrass in grass grown for seed production. For maximum suppression, annual bluegrass must be sprayed when in the flowering stage, and must receive thorough coverage. Kudos 27.5 WDG may not be effective against some annual bluegrass biotypes.

Limitation on Grass Grown for Seed:

- Kudos 27.5 WDG is rainfast 1 hour after application.

Restrictions on Grass Grown for Seed:

- **DO NOT** apply more than a total of 29 ounces of Kudos 27.5 WDG (8 g ai) per acre, per year.
- **DO NOT** apply within 35 days before harvest.
- **DO NOT** graze livestock for 49 days following application.
- **DO NOT** cut forage or hay for livestock feed for 49 days following application.
- Plant back/rotation restriction: If replanting or crop rotation is necessary in fields treated with Kudos 27.5 WDG, **DO NOT** plant any crop other than grass grown for seed for 30 days following application.
- Restricted Entry Interval (REI): 12 hours.

PEANUTS (all approved states except California):

Kudos 27.5 WDG can be used as a production management tool on peanuts to suppress vegetative growth i.e. shorten internode length, resulting in better harvest efficiency. Kudos 27.5 WDG does not affect vegetative growth the year following application.

For consistent performance on peanuts, add one pound of a spray grade ammonium sulfate (AMS) for every pound of Kudos 27.5 WDG. Use a standard spray adjuvant, preferably a non-ionic surfactant, to improve leaf coverage and performance consistency. If Kudos 27.5 WDG is being applied without a tank-mix partner, a non-phytotoxic crop oil concentrate (COC) can be used in place of a non-ionic surfactant. Always follow the manufacturer’s rate recommendations.

Apply Kudos 27.5 WDG to actively growing peanut plants according to the rates recommended in Table 4. Initially apply 7.25 ounces (2 g ai) per acre when 50% of the peanut stems are touching the row middle i.e. row closure. Make a second application at 100% row closure, as needed. Under conditions that are favorable to extremely rank growth, a third application may be required. **DO NOT** make more than two applications of Kudos 27.5 WDG in less than 6 weeks. Make Kudos 27.5 WDG applications prior to loss of visual row pattern. Crops under stress due to lack of moisture, hail damage, flooding, mechanical injury, etc. will show little response to Kudos 27.5 WDG.

Table 4. Application Rates and Timings for Vegetative Growth Control in Peanuts.

Application Timing	Application Rate per Acre
First application: <ul style="list-style-type: none">• Apply when 50% of the peanut stems are touching in row middle (row closure).	7.25 oz (2 g ai)
Second application: <ul style="list-style-type: none">• Make a second application at 100% row closure as needed.	3.6-7.25 oz (1-2 g ai)

Ground Application: Use a minimum of 20 gallons of spray solution per broadcast acre.

Limitation on Peanuts:

- Kudos 27.5 WDG is rainfast 8 hours after application.

Restrictions on Peanuts:

- **DO NOT** apply more than a total of 21.75 oz of Kudos 27.5 WDG (6 g ai) per acre, per year.
- **DO NOT** make more than 2 applications of Kudos 27.5 WDG in less than 6 weeks.
- **DO NOT** within 25 days before harvest.
- **DO NOT** graze or feed livestock treated crops.
- **DO NOT** apply Kudos 27.5 WDG on peanuts by air.
- **DO NOT** apply to crops under stress due to lack of moisture, hail damage, flooding, herbicide injury, or mechanical injury as reduced activity may result.
- **DO NOT** apply to crops that show injury (leaf phytotoxicity or plant stunting) produced by any other prior product applications because this injury can be enhanced or prolonged.
- Plant back/rotation restriction: If replanting or crop rotation is necessary in fields treated with Kudos 27.5 WDG, **DO NOT** plant any crop other than peanuts for 30 days following the last application.
- Restricted Entry Interval (REI): 12 hours.

PEARS - d'Anjou:

Kudos 27.5 WDG can be used as a production management tool on d'Anjou pears to suppress vegetative growth, i.e., shoot growth reduction, resulting in a balance between canopy development and fruit production. Growth suppression with Kudos 27.5 WDG generally lasts for 2-5 weeks per application. Kudos 27.5 WDG does not affect vegetative growth the season following application.

Benefits resulting from the use of Kudos 27.5 WDG include:

- Reduction in vegetative shoot growth
- Lessens the need or frequency for summer and/or dormant pruning
- Improved light penetration into the tree canopy
- Reduced incidence and severity of fire blight of shoots

During seasons when both Kudos 27.5 WDG and products containing gibberellins, e.g. Novagib 10L, are used, reductions in efficacy may occur in the Kudos 27.5 WDG and/or gibberellin treatments.

The use of Kudos 27.5 WDG may result in an increase in fruit retention. Therefore, thinning programs may need to be adjusted when using Kudos 27.5 WDG.

When used as recommended, Kudos 27.5 WDG can reduce the incidence and severity of fire blight infection in d'Anjou pears in two ways: 1) The active ingredient in Kudos 27.5 WDG can reduce latent bloom in d'Anjou pear. During bloom pear trees are more susceptible to infection from fire blight. By reducing the bloom period and thusly reducing the time frame for infection to occur, the chance of fire blight infection may be decreased. 2) Applications of Kudos 27.5 WDG reduce vegetative growth in d'Anjou

pears. This reduction in vegetative growth helps lessen host susceptibility. This decrease in susceptibility will not become effective until approximately 10 days after application. Kudos 27.5 WDG does not have direct antibiotic activity against fire blight. For maximum reduction in fire blight susceptibility, apply Kudos 27.5 WDG at least 10 days before the occurrence of weather conditions favorable for shoot and leaf infections. Use Kudos 27.5 WDG as just one component of a comprehensive control strategy against fire blight.

Application Timings: For vegetative growth control, make the first application of Kudos 27.5 WDG when shoots have 1-3" of new shoot growth. For best results, make subsequent applications at 1-4 week intervals, before or immediately following signs of shoot regrowth.

Number of Applications: The number of applications will vary depending on the timing of the first application, tree vigor, fruit load, pruning, variety, rootstock, and/or the management history of the orchard. For pear orchards in locations with a long growing season or on high vigor trees and/or trees with light fruit load, 3 to 5 applications per season may be required.

Tree Vigor: Adjust the Kudos 27.5 WDG rate according to the vegetative vigor of the trees (see Table 5). Vegetative vigor can be influenced by many factors, including variety, rootstock, fruit load, cultural management and location.

Tree Size: Calculate the rate per acre of Kudos 27.5 WDG based on tree size. The application rate should be based on the volume of water needed to spray the trees to drip i.e. dilute spray or Tree Row Volume (TRV). Consult your local extension agent or consultant for a recommendation on spray volume.

Application Rates: Recommended rates will be based on desired treatment effect, i.e., vegetative growth control, shoot blight suppression, etc. For vegetative growth control, Kudos 27.5 WDG rates should be based on level of tree vigor, size and/or length of growing season. Refer to Table 1 for recommended rates per application.

For consistent performance on pears, add one pound of a spray grade ammonium sulfate (AMS) for every pound of Kudos 27.5 WDG. Use a standard spray adjuvant, preferably a non-ionic surfactant, to improve leaf coverage and performance consistency. Follow the manufacturer's rate recommendations.

Table 5. Recommended Application Rates on d’Anjou Pears

Application Timing	Rate per Acre¹	Rate per 100 Gallons of Dilute Spray²	Restrictions
For Vegetative Growth Control: First application: <ul style="list-style-type: none"> • Apply at 1-3” of new shoot growth Subsequent applications: <ul style="list-style-type: none"> • Make any subsequent applications at 1-4 week intervals and before or immediately after shoot regrowth occurs. 	18 oz	6 oz	See “General Restrictions & Limitations on [d’Anjou] Pears” section below”
For Vegetative Growth Control and Reduced Latent Bloom (Fire Blight Management): <ul style="list-style-type: none"> • Apply at 1-3” inches of new shoot growth. • Make a second application after 21 days. 	30-36 oz	10-12 oz	See “General Restrictions & Limitations on [d’Anjou] Pears” section below.

¹ Based on 300 gallons of dilute spray per acre.

² Apply 6-12 oz of Kudos 27.5 WDG per 100 gallons of dilute spray.

Ground Application: Apply Kudos 27.5 WDG to actively growing trees using calibrated ground equipment. Kudos 27.5 WDG may be applied using either dilute or concentrate spray equipment as long as sufficient spray coverage is achieved. Direct spray to the portion of the tree where growth control is desired. To achieve good coverage, apply Kudos 27.5 WDG in a sufficient spray volume per acre utilizing proper spray pressure, nozzles, nozzle spacing and tractor speed.

Aerial Application (All approved states except California): Apply Kudos 27.5 WDG in a minimum of 10 gallons of spray solution per broadcast acre. Aerial applications generally only provide spray coverage in the top part of the tree canopy. Vegetative growth control will be limited to those areas of the canopy that receive spray coverage.

Limitation on d’Anjou Pears:

- Kudos 27.5 WDG is rainfast 8 hours after application.

Restrictions on d’Anjou Pears:

- **DO NOT** apply more than a total of 99 oz (6.2 lbs) of Kudos 27.5 WDG per acre, per year.
- **DO NOT** apply more than a total of 48 oz (3 lbs) of Kudos 27.5 WDG within any 21-day interval.
- **DO NOT** apply within 45 days before harvest.
- Restricted Entry Interval (REI): 12 hours.

- **DO NOT** apply to crops that show injury (leaf phytotoxicity) produced by any other prior pesticide applications because this injury can be enhanced or prolonged.
- **DO NOT** apply Kudos 27.5 WDG to d'Anjou pears through any type of irrigation system.

STRAWBERRIES (all approved states except California):

Kudos 27.5 WDG can be used as a production management tool on strawberries to suppress vegetative growth i.e. shoot growth reduction. The growth suppression effects of Kudos 27.5 WDG on strawberries will normally last for 2-5 weeks per application during the current growing season. Kudos 27.5 WDG does not affect vegetative growth the year following application.

For consistent performance on strawberries, add one pound of a spray grade ammonium sulfate (AMS) for every pound of Kudos 27.5 WDG. Use a standard spray adjuvant, preferably a non-ionic surfactant, to improve leaf coverage and performance consistency. Follow the manufacturer's rate recommendations.

Apply Kudos 27.5 WDG to strawberries at a rate of 2.0 oz/acre per application. Make the first application of Kudos 27.5 WDG in the fall once a minimum of 80% of the strawberry plants have 5 fully expanded leaves and before any runner formation has begun. If additional vegetative growth control is needed, make subsequent applications at 14-21 day intervals.

Ground Application: Apply Kudos 27.5 WDG to the foliage of actively growing strawberries using calibrated ground application equipment. Kudos 27.5 WDG is taken up through the leaves of the plants so good spray coverage of the plant foliage is essential to get good uptake and consistent growth suppression. To achieve good coverage, apply Kudos 27.5 WDG in a sufficient spray volume per acre utilizing proper spray pressure, nozzles, nozzle spacing and tractor speed. It is recommended to apply in a minimum spray volume of 10 gallons/acre and not more than 30 gallons/acre.

Limitation on Strawberries:

- Kudos 27.5 WDG is rain-fast 8 hours after application.

Restrictions on Strawberries:

- **DO NOT** apply within 21 days before harvest.
- **DO NOT** use Kudos 27.5 WDG on strawberries grown in greenhouses.
- **DO NOT** apply this product to strawberries using air-blast or aerial spray equipment.
- **DO NOT** apply Kudos 27.5 WDG to strawberries using any type of irrigation system.
- **DO NOT** apply to crops that show injury (leaf phytotoxicity) produced by any other prior pesticide applications as this injury can be enhanced or prolonged. Refer to General Tank Mixing Information for additional tank mixing instructions and precautions.
- **DO NOT** tank mix or apply Kudos 27.5 WDG with calcium or boron products as the addition of these products to the spray mix can reduce product effectiveness.
- **DO NOT** apply more than 6 oz of Kudos 27.5 WDG per year.

- Do not make more than 3 applications of Kudos 27.5 WDG to strawberries per year.
- Restricted Entry Interval (REI) is 12 hours.

SWEET CHERRIES (All approved states except California):

Kudos 27.5 WDG can be used as a production management tool on sweet cherries to suppress vegetative growth i.e. shoot growth reduction, which can reduce or delay the need for pruning. Kudos 27.5 WDG does not affect vegetative growth the year following application.

For consistent performance on sweet cherries, add one pound of a spray grade ammonium sulfate (AMS) for every pound of Kudos 27.5 WDG. Use a standard spray adjuvant, preferably a non-ionic surfactant, to improve leaf coverage and performance consistency. Follow the manufacturer’s rate recommendations.

Apply Kudos 27.5 WDG to sweet cherries according to the rates recommended in **Table 6**. Make the first application of Kudos 27.5 WDG in the spring when shoots have 1-3” of new growth. If additional vegetative growth control is needed, make subsequent applications at 1 to 4 week intervals, before or immediately following signs of shoot regrowth.

Number of Applications: The number of applications will vary depending on the timing of the first application, tree vigor, fruit load, pruning, variety, rootstock, and/or the management history of the orchard. For sweet cherry orchards in locations with a long growing season or on high vigor trees and/or trees with light fruit load, 3-5 applications per season may be required.

Table 6. Application Rates for Vegetative Growth Control in Sweet Cherries

Tree Vigor	Application Rate per Acre	Restrictions
High Vigor Trees	8-20 oz (2.2-5.5 g ai)	<ul style="list-style-type: none"> • DO NOT apply more than 20 oz/A within any 14-day interval. • DO NOT apply more than 40 oz/A per year.
Medium Vigor Trees	8-12 oz (CA only) (2.2-3.3 g ai) 6-12 oz (all other states) (1.7-3.3 g ai)	<ul style="list-style-type: none"> • DO NOT apply more than 12 oz/A within any 14-day interval. • DO NOT apply more than 30 oz/A per year.
Low Vigor Trees	Not applicable	<ul style="list-style-type: none"> • DO NOT apply Kudos 27.5 WDG to low vigor trees.

Ground Application: Apply Kudos 27.5 WDG to actively growing trees using calibrated ground equipment. Kudos 27.5 WDG may be applied using either dilute or concentrate spray equipment as long as sufficient spray coverage is achieved. To achieve good coverage, apply Kudos 27.5 WDG in a sufficient spray volume per acre utilizing proper spray pressure, nozzles, nozzle spacing and tractor speed.

Aerial Application (All approved states except California): Apply Kudos 27.5 WDG in a minimum of 10 gallons of spray solution per broadcast acre. Aerial applications generally only provide spray coverage in the top part of the tree canopy. Vegetative growth control will be limited to those areas of the canopy that receive spray coverage.

Limitation on Sweet Cherries:

- Kudos 27.5 WDG is rainfast 8 hours after application.

Restrictions on Sweet Cherries:

- **DO NOT** apply more than a total amount of Kudos 27.5 WDG per acre, per year as outlined in the **Application Rates for Vegetative Growth Control in Sweet Cherries** table in this label.
- **DO NOT** apply more than a total of 20 ounces of Kudos 27.5 WDG (5.5 g ai) in any 14-day interval.
- **DO NOT** apply within 20 days before harvest.
- Restricted Entry Interval (REI): 12 hours.
- **DO NOT** use Kudos 27.5 WDG on tart cherries.
- **DO NOT** apply to crops that show injury (leaf phytotoxicity) produced by any other prior pesticide applications because this injury can be enhanced or prolonged. Refer to General Tank Mixing Information for additional tank mixing instructions and precautions.
- **DO NOT** apply Kudos 27.5 WDG on low vigor trees.

TURF:

Kudos 27.5 WDG can be used as a production management tool on turf to reduce vegetative growth. The active ingredient in Kudos 27.5 WDG, prohexadione-ca, reduces the growth of turf by inhibiting the biosynthesis of gibberellin, a natural plant hormone that regulates cell elongation. The inhibition of gibberellin slows the growth of turf which will result in reduced frequency of mowing and amounts of grass clippings in treated areas. Following an application of Kudos 27.5 WDG to turf, a reduction in growth and grass clippings will typically be observed by three days.

Kudos 27.5 WDG can be applied to turf that is green and actively growing. Under conditions of stress, delay applications. Apply lower rates if the turf is entering stressful growing conditions such as pest pressure, high temperatures, low moisture conditions or certain cultural practices such as, but not limited to, aerification or verticutting. Under certain environmental conditions, such as those that may occur early in the season, transient bronzing may be observed. For information on application rates and timings on turf for Kudos 27.5 WDG see Application Rates & Timings for Kudos 27.5 WDG in Turf (Table7) below.

For consistent performance on turf, add one pound of a spray grade ammonium sulfate (AMS) for every pound of Kudos 27.5 WDG applied. Use a standard spray adjuvant, preferably a non-ionic surfactant, to improve leaf coverage and performance consistency. Follow the manufacturer's rate recommendations.

Ground Application: Apply Kudos 27.5 WDG to actively growing turf using calibrated application equipment. Kudos 27.5 WDG is taken up through the leaves of the plants. So, good spray coverage of the plant foliage is essential to get good uptake and consistent growth suppression. To achieve good coverage, apply Kudos 27.5 WDG in a sufficient spray volume utilizing proper spray pressure, nozzles, nozzle spacing and applicator speed. Apply Kudos 27.5 WDG in a spray volume of 1 to 2 gallons water/1000 sq. ft. but it can be applied in a spray volume of 0.5 to 5.0 gallons water/1000 sq. ft. Avoid applying Kudos 27.5 WDG during warm, low humidity or windy conditions. The absorption of the spray mix is improved with longer drying times.

Maintain constant agitation during mixing of the spray solution and application. Do not allow the spray tank to sit for prolonged periods without any agitation. Do not allow the spray mix to sit in the tank overnight.

Mowing & Irrigation: Mowing and irrigation events can affect the efficacy of the Kudos 27.5 WDG applications. After mowing, do not apply Kudos 27.5 WDG for at least one hour. Following an application of Kudos 27.5 WDG, do not mow treated turf for at least one day and do **not** apply irrigation for 4 hours.

Turf Maintenance: Following applications of Kudos 27.5 WDG, maintain the areas of turf that were treated as normal with proper levels of irrigation and fertility. Implement a good pest control plan as necessary to maintain the turf in a healthy condition. If tank mixing with other product(s) precaution is advised. Test tank mixes that the user is unfamiliar with on a small areas before the treatment is applied to larger areas. Test tank mixes for compatibility before mixing a larger quantity for an application. User assumes all risk for tank mixes as it is not possible for manufacturer to test all possible tank mix combinations for compatibility, safety and effect on efficacy of the products.

Table 7. Applications for Growth Management, Color and Quality Enhancement in Turf Grown for Sod

Turf Type	oz/acre	Application Interval	Notes
Warm Season Turf such as, but not limited to, bermudagrass	14.5 – 43.6	2-4 week interval	Since turfgrass response can be influenced by a variety of factors, limit the initial evaluation of Kudos 27.5 WDG to a small area.
Cool Season Turf such as, but not limited to, Kentucky bluegrass, perennial ryegrass & fescues	7.25 – 29.1	2-4 week interval	Determine the acceptability of the turfgrass response using a lower use rate and shorter interval before gradually moving to higher rates and longer intervals.

Limitation on Turf:

- Kudos 27.5 WDG is rainfast 1 hour after application.

Restrictions on Turf:

- **DO NOT** apply more than the total amount of Kudos 27.5 WDG per acre per year as stated above.
- **DO NOT** apply more than 26 oz per 1000 square feet of Kudos 27.5 WDG per year.
- **DO NOT** tank mix or apply Kudos 27.5 WDG with calcium or boron products as the addition of these products to the spray mix can reduce product effectiveness.
- **DO NOT** apply Kudos 27.5 WDG to turf that has been sodded or sprigged until it has knitted down and rooted firmly.
- **DO NOT** apply to turf growing in conditions that are not ideal for good growth. Also, **DO NOT** apply to turf that shows injury (leaf phytotoxicity) produced by any other prior pesticide applications as this type of injury can be enhanced or prolonged. Refer to General Tank Mixing Information for additional tank mixing instructions and precautions.
- **DO NOT** graze treated areas or feed clippings previously treated with Kudos 27.5 WDG to livestock.
- **DO NOT** apply Kudos 27.5 WDG to turf using any type of irrigation/chemigation system.
- Following an application of Kudos 27.5 WDG, **DO NOT** mow treated turf for at least one day and **DO NOT** apply irrigation for 4 hours.
- Restricted Entry Interval (REI): 12 hours.

WATERCRESS (all approved states except California):

Kudos 27.5 WDG can be used as a production management tool on watercress to suppress vegetative growth and increase stem diameter in watercress. Following an application of Kudos 27.5 WDG to watercress a reduction in growth will typically be observed after 4 days. This reduction in growth may last up to 14 days depending on growing conditions.

DO NOT apply Kudos 27.5 WDG to watercress with any additive, adjuvant, fertilizer, nutrient or any other pesticide product added to the spray mix. Only mix Kudos 27.5 WDG with water when making applications to watercress.

Apply Kudos 27.5 WDG to watercress which has regrown from the plant stubble remaining following the previous harvest. Depending on the growing conditions the first application may be made from 5 to 10 days. If deemed necessary a second application of Kudos 27.5 WDG may be applied. Application rates and volumes are noted in **Table 11. Kudos 27.5 WDG Application Rates and Timings for Vegetative Growth Control of Watercress.**

Kudos 27.5 WDG may be applied through ground or sprinkler application equipment. Application equipment must be properly designed and calibrated before making applications of Kudos 27.5 WDG to watercress. To achieve desired results, apply Kudos 27.5 WDG in a sufficient spray volume per acre that will provide good coverage based on the size and density of the watercress foliage. Adequate coverage of the plant foliage must be achieved for good activity but avoid over application that results in runoff.

Ground Application: Apply Kudos 27.5 WDG to the foliage of actively growing watercress using properly calibrated ground application equipment. Ensure adequate spray coverage is applied to achieve good uptake of the spray solution.

Sprinkler Irrigation Application: Chemigation applications of Kudos 27.5 WDG to watercress are to **only** be made through sprinkler type irrigation systems such as center pivot, lateral move, solid set or hand move irrigation systems.

The chemical tank and injector system must be thoroughly cleaned before an application of Kudos 27.5 WDG is made. Flush system with clean water prior to beginning the Kudos 27.5 WDG application.

Application Instructions: Apply Kudos 27.5 WDG at the rates and timings stated in this label. Chemigation applications must be made as concentrated as possible. To achieve best results apply at 0.1 inch up to 0.15 inch (2,716 to 4,073 gallons) of water per acre.

Sprinkler Irrigation Application Use Directions:

- Apply Kudos 27.5 WDG only through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. **DO NOT** apply this product through any other type of irrigation system.
- Add Kudos 27.5 WDG to the pesticide supply tank containing sufficient water to maintain a continuous flow by the injection equipment. In continuous moving systems, inject this product/water mixture continuously, to apply the proper labeled rate per acre for that crop. **DO NOT** exceed 1/2 inch water (13,577 gallons) per acre. In stationary or non-continuous moving systems, inject the product/water mixture in the last 15 to 30 minutes of each set allowing sufficient time for all of the required pesticide to be applied by all the sprinkler heads and applying the labeled rate per acre for that crop. **DO NOT** apply if wind speed favors drift beyond the intended area of the treatment. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. Thorough coverage of foliage is required for good control. Good agitation must be maintained during the entire application period.
- Contact state extension service specialists, equipment manufacturers or other experts if you have questions or concerns about calibration of the system.
- The system must contain a functional check valve, vacuum-relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from back-flow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

- Functional interlocking controls must be contained in the system to automatically shut off the pesticide-injection pump when the water pump motor stops.
- In order to prevent a water pressure decrease to the point where pesticide distribution is adversely affected, the irrigation line or water pump must include a functional pressure switch that will stop the motor.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- The irrigation water should not be turned off before allowing sufficient time for the pesticide to be flushed through all lines and all nozzles. A person knowledgeable of the chemigation system and responsible for its operation, or under supervision of the responsible person, should shut down the system and make necessary adjustments should the need arise.
- **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

Specific Instructions for Public Water Systems:

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Table 8. Kudos 27.5 WDG Application Rates and Timings for Vegetative Growth Control of Watercress

Kudos 27.5 WDG Rate (oz/acre/application)	Application Timing
8	Apply to watercress which has regrown leaves/foliage from the plant stubble remaining following the previous harvest (5 to 10 days).
8	If deemed necessary a second application may be made 7 days after the first application. This second application may or may not be needed depending on growing conditions.

Limitation on Watercress:

- Kudos 27.5 WDG is rain-fast 8 hours after application.

Restrictions on Watercress:

- **DO NOT** apply more than the total amount of Kudos 27.5 WDG per acre per year outlined above.
- **DO NOT** apply more than 16 oz of Kudos 27.5 WDG per acre per year.
- **DO NOT** apply Kudos 27.5 WDG within 3 days before harvest.
- **DO NOT** use Kudos 27.5 WDG on watercress grown in greenhouses.
- **DO NOT** apply to watercress crops that show injury (leaf phytotoxicity) produced by any other prior pesticide applications as this type of injury can be enhanced or prolonged. Refer to General Tank Mixing Information for additional tank mixing instructions and precautions.
- **DO NOT** apply Kudos 27.5 WDG to watercress with any additive, adjuvant, fertilizer, nutrient or any other pesticide product added to the spray mix. Only mix Kudos 27.5 WDG with water when making applications to watercress.
- Restricted Entry Interval (REI): 12 days.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container only. After partial use, close the container tightly. Store in a secure place that is cool and dry. Use spray and stock solutions within 24 hours. Immediate use is required if another component is added to the spray solution.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY

Fine Agrochemicals Limited ("FINE") warrants that this Product conforms to the specifications on this label. To the extent consistent with applicable law, FINE makes no other warranties and disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for a particular purpose. No agent of FINE or any other person is authorized to make any representation or warranty beyond those contained herein.

It is impossible to eliminate all risks associated with this Product. Plant injury, lack of performance, or other unintended consequences may result because of factors such as abnormal weather conditions, use of the Product other than in strict accordance with this label's instructions, presence of other materials, the manner of application or other factors, all of which are beyond the control of FINE or the seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

To the extent consistent with applicable law: 1) FINE disclaims any liability whatsoever for special, incidental or consequential damages resulting from the handling or use of this Product and 2) FINE's liability under this label shall be limited to the amount of the purchase price or, at the election of FINE, the free replacement of the Product.

© Fine Agrochemicals Limited, Hill End House, Whittington, Worcester, WR5 2RQ, United Kingdom

Kudos® 27.5 WDG is a registered trademark of Fine Agrochemicals, Ltd.
Novagib® is a registered trademark of Fine Agrochemicals Ltd.

SUBLABEL B: Turf uses

KUDOS® 27.5 WDG

Plant growth regulator for use on turf.

Active Ingredient: Prohexadione-calcium

[calcium 3-oxido-5-oxo-4-propionylcyclohex-3-enecarboxylate].....27.5%

Other Ingredients:72.5%

Total:100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

[See inside panel for First Aid Statements]

EPA Reg. No. 62097-41

EPA Est. No.

Net Contents:

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Poison Control Hotline at 1-800-222-1222 for emergency medical treatment information 24 hours a day, seven days a week.	

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION. Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE items separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your state or tribe, consult the state or tribal agency responsible for pesticide regulation.

TURF:

Kudos 27.5 WDG can be used as a production management tool on turf to reduce vegetative growth. The active ingredient in Kudos 27.5 WDG, prohexadione-ca, reduces the growth of turf by inhibiting the biosynthesis of gibberellin, a natural plant hormone that regulates cell elongation. The inhibition of gibberellin slows the growth of turf which will result in reduced frequency of mowing and amounts of grass clippings in treated areas. Following an application of Kudos 27.5 WDG to turf, a reduction in growth and grass clippings will typically be observed by three days.

Kudos 27.5 WDG can be applied to turf that is green and actively growing. Under conditions of stress, applications should be delayed. Apply lower rates if the turf is entering stressful growing conditions such as pest pressure, high temperatures, low moisture conditions or certain cultural practices such as, but not limited to, aerification or verticutting. Under certain environmental conditions, such as those that may occur early in the season, transient bronzing may be observed. For information on application rates and timings on turf for Kudos 27.5 WDG see Application Rates & Timings for Kudos 27.5 WDG in Turf (Table 1) below.

For consistent performance on turf, add one pound of a spray grade ammonium sulfate (AMS) for every pound of Kudos 27.5 WDG applied. Use a standard spray adjuvant, preferably a non-ionic surfactant, to improve leaf coverage and performance consistency. Follow the manufacturer's rate recommendations.

Ground Application: Apply Kudos 27.5 WDG to actively growing turf using calibrated application equipment. Kudos 27.5 WDG is taken up through the leaves of the plants. So, good spray coverage of the plant foliage is essential to get good uptake and consistent growth suppression. To achieve good coverage, apply Kudos 27.5 WDG in a sufficient spray volume utilizing proper spray pressure, nozzles, nozzle spacing and applicator speed. Apply Kudos 27.5 WDG in a spray volume of 1 to 2 gallons water/1000 sq. ft. but it can be applied in a spray volume of 0.5 to 5.0 gallons water/1000 sq. ft. Avoid applying Kudos 27.5 WDG during warm, low humidity or windy conditions. The absorption of the spray mix is improved with longer drying times.

Maintain constant agitation during mixing of the spray solution and application. Do not allow the spray tank to sit for prolonged periods without any agitation. Do not allow the spray mix to sit in the tank overnight.

Mowing & Irrigation: Mowing and irrigation events can affect the efficacy of the Kudos 27.5 WDG applications. After mowing, do not make an application of Kudos 27.5 WDG for at least one hour. Following an application of Kudos 27.5 WDG, do not mow the treated turf for at least one day and do **not apply** irrigation for 4 hours.

Turf Maintenance: Following applications of Kudos 27.5 WDG, maintain the areas of turf that were treated as normal with proper levels of irrigation and fertility. Implement a good pest control plan as necessary to maintain the turf in a healthy condition. If tank mixing with other product(s) precaution is advised. The user should test tank mixes that are unfamiliar on a small area before the treatment is applied to larger areas. Test tank mixes for compatibility before mixing a larger quantity for an application. User assumes all risk for tank mixes as it is not possible for manufacturer to test all possible tank mix combinations for compatibility, safety and effect on efficacy of the products.

Table 1. Application Rates & Timings

	Kudos 27.5 WDG Applications to Slow Vertical Growth, Reduce Mowing, Improve Color & Quality			
	Fairway & Rough Areas (Application Intervals: 2-4 Weeks)		Green & Tee Areas (Application Intervals: 1-2 Weeks)	
	oz/acre	oz/1000 sq. ft.	oz/acre	oz/1000 sq. ft.
Warm Season Turf*				
Hybrid Bermudagrass including Tifway 419, TifSport & Riviera	29.1 – 43.6	0.67 – 1.0	7.25 – 14.5	0.17 – 0.34
Cool Season Turf				
Bentgrass: L-93, Penn A4, Penn G2, Pennlinks, Penneagle, Southshore	7.25 – 14.5	0.17 – 0.34	1.8 – 7.25	0.04 – 0.17
Kentucky bluegrass: Midnight, Odyssey	14.5 – 21.8	0.34 – 0.50		
Perennial ryegrass: Manhattan IV	21.8 – 29.1	0.5 – 0.67		

*Lower rates may be applied for color and quality enhancement. When using shorter application intervals, performance should first be evaluated using the lower recommended rates.

Table 2. Applications for Overseeding Enhancement in Bermudagrass.

Turf Type	oz/acre	Application Interval	Notes
Hybrid Bermudagrass (including Tifway 419, TifSport & Riviera)	0.90 – 1.75	Apply one application of Kudos 27.5 WDG 3-5 days prior to seeding. Any preparation of the site such as verticutting, spiking or scalping should be done 1-2 days after the application.	Kudos 27.5 WDG is a foliar-active product. Therefore germination and seedling growth of the over-seeding are not affected when used as directed. Use good cultural practices, such as fertilization and

			irrigation, in conjunction with Kudos 27.5 WDG to optimize results. Some temporary yellowing of the bermudagrass turf may be observed, however, this will not affect the over-seeding.
--	--	--	--

Table 3. Applications for Reduction of *Poa annua* in Bermudagrass Overseeding.

Turf Type	oz/acre	Application Interval
Hybrid Bermudagrass (including Tifway 419, TifSport & Riviera)	0.90 – 1.75	Reapply Kudos 27.5 WDG at 3-4 week intervals during periods of active <i>Poa annua</i> growth.

Limitation on Turf:

- Kudos 27.5 WDG is rainfast 1 hour after application.

Restrictions on Turf:

- **DO NOT** apply more than the total amount of Kudos 27.5 WDG per acre per year as stated above.
- **DO NOT** apply more than 26 oz per 1000 square feet of Kudos 27.5 WDG per year.
- **DO NOT** tank mix or apply Kudos 27.5 WDG with calcium or boron products as the addition of these products to the spray mix can reduce product effectiveness.
- **DO NOT** apply Kudos 27.5 WDG to turf that has been sodded or sprigged until it has knitted down and rooted firmly.
- **DO NOT** apply to turf growing in conditions that are not ideal for good growth. Also, **DO NOT** apply to turf that shows injury (leaf phytotoxicity) produced by any other prior pesticide applications as this type of injury can be enhanced or prolonged. Refer to General Tank Mixing Information for additional tank mixing instructions and precautions.
- **DO NOT** graze treated areas or feed clippings previously treated with Kudos 27.5 WDG to livestock.
- **DO NOT** apply Kudos 27.5 WDG to turf using any type of irrigation/chemigation system.
- Restricted Entry Interval (REI): 12 hours.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container only. After partial use, close the container tightly. Store in a secure place that is cool and dry. Use spray and stock solutions within 24 hours. Immediate use is required if another component is added to the spray solution.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY

Fine Agrochemicals Limited ("FINE") warrants that this Product conforms to the specifications on this label. To the extent consistent with applicable law, FINE makes no other warranties and disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for a particular purpose. No agent of FINE or any other person is authorized to make any representation or warranty beyond those contained herein.

It is impossible to eliminate all risks associated with this Product. Plant injury, lack of performance, or other unintended consequences may result because of factors such as abnormal weather conditions, use of the Product other than in strict accordance with this label's instructions, presence of other materials, the manner of application or other factors, all of which are beyond the control of FINE or the seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

To the extent consistent with applicable law: 1) FINE disclaims any liability whatsoever for special, incidental or consequential damages resulting from the handling or use of this Product and 2) FINE's liability under this label shall be limited to the amount of the purchase price or, at the election of FINE, the free replacement of the Product.

© Fine Agrochemicals Limited, Hill End House, Whittington, Worcester, WR5 2RQ, United Kingdom

Kudos® 27.5 WDG is a registered trademark of Fine Agrochemicals, Ltd.