



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

October 1, 2021

Georgia Anastasiou
Agent for Quimetal Industrial S.A.
2641 South Clark Street, Suite 710
Arlington, VA 22202

Subject: Label Amendment – Adding language to resistance management and crop and application rates sections per PRN 2003-1; other minor label changes
Product Name: Sulfur Dust
EPA Registration Number: 62562-11
Application Date: 11/30/2018
Decision Number: 546970

Dear Ms. Miller:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 62562-11
Decision No. 546970

with FIFRA section 6. If you have any questions, please contact Jasmin Jackson by phone at 703-347-0732, or via email at jackson.jasmin@epa.gov.

Sincerely,

A handwritten signature in cursive script that reads "Paul Di Salvo".

Paul Di Salvo, MPS, AWB®
Special Assistant / Wildlife Biologist
Registration Division (7505P)
Office of Pesticide Programs

Enclosure: Stamped Label

SULFUR	GROUP	UN	INSECTICIDE
SULFUR	GROUP	M02	FUNGICIDE

ACCEPTED
10/01/2021
Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 62562-11

SULFUR DUST

For Organic Production



Active Ingredient:	
Sulfur.....	99.99%
Other Ingredients.....	0.01%
TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Side Panel for Additional Precautionary Statements

FIRST AID	
If on skin	<ul style="list-style-type: none"> •Take off contaminated clothing. •Rinse skin immediately with plenty of water for 15-20 minutes. •Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> •Hold eye open and rinse slowly and gently with water for 15-20 minutes. •Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. •Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If swallowed	<ul style="list-style-type: none"> •Call poison control center or doctor immediately for treatment advice. •Have person sip a glass of water if able to swallow. •Do not induce vomiting unless told to do so by the poison control center or doctor. •Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	

EPA File Symbol 62562-11

EPA Est. No. _____

Manufactured by: *QUIMETAL INDUSTRIAL S.A.*
Los Yacimientos 1301 • Maipu • Santiago • Chile
Phone: (56-2) 2381 7000
PRODUCT OF CHILE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear: A) Long-sleeved shirt and long pants; B) Chemical-resistant gloves made of any waterproof material including polyethylene or polyvinyl chloride ≥ 14 mils; C) Shoes plus socks; D) Goggles or faceshield.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

User should: • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate. Do not apply when weather conditions favor drift from treated areas.

PHYSICAL AND CHEMICAL HAZARDS

Sulfur dust suspended in air ignites easily. Keep away from heat, sparks or flame. Do not smoke while applying this product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours. PPE required for early entry to treated areas that is permitted under the worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is: A) Coveralls; B) Chemical-resistant gloves made of any waterproof material including polyethylene or polyvinyl chloride ≥ 14 mils; C) Shoes plus socks; D) Goggles or faceshield.

USE PRECAUTIONS

IMPORTANT: Sulfur may burn foliage and fruit of sensitive varieties if applied when temperatures exceed 90°F or during periods when the temperatures within 3 days after application are expected to exceed 90°F in the treated area.

USE RESTRICTIONS

DO NOT make applications within 2 weeks of an oil spray unless specified otherwise, for the individual crop, in the CROPS AND APPLICATION RATES section (see Citrus).

CALIFORNIA DRIFT RESTRICTIONS

The following conditions must be followed when using this product in the State of California for both ground and air applications.

1. The operator of the property and the applicator must establish a buffer zone of enough distance to prevent drift onto non-target areas including hospitals, clinics, schools, residential areas and any other area designated by the county agricultural commissioner.
2. The operator of the property and the applicator must evaluate the method and equipment for each site to ensure proper and safe use. Evaluations shall include, but not be limited to the appropriateness of ground or air application.
3. Sulfur -must not be applied when the wind velocity exceeds 10 miles per hour. Applicators must be aware that in some areas, "dead calm" conditions are often associated with an inversion situation. In these areas, applying sulfur when there is a minimum air movement of 2 miles per hour will ensure that an inversion situation does not exist.
4. When applying sulfur dust, best application practices must be utilized. Contact your local pest control advisor or county cooperative extension advisor to obtain information regarding best application practices.

RESISTANCE MANAGEMENT

Fungicide Resistance Management

For resistance management, Sulfur Dust contains a Group M02 fungicide. Any fungal population may contain individuals naturally resistant to Sulfur Dust and other Group M02 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of Sulfur Dust or other Group M02 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact QUIMETAL INDUSTRIAL S.A. at 56-2-2381-7000.. You can also contact your pesticide distributor or university extension specialist to report resistance.

Insecticide Resistance Management

For resistance management, Sulfur Dust contains a Group UN acaricide. Any mite population may contain individuals naturally resistant to Sulfur Dust and other Group UN acaricides. The resistant individuals may dominate the mite population if this group of acaricides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay acaricide resistance, take the following steps:

- Rotate the use of Sulfur Dust or other Group UN acaricides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with acaricides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues for the targeted pests between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual acaricides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for mite resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).

- Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
- The mite resistance management benefits of an acaricide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an mite resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for acaricide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact QUIMETAL INDUSTRIAL S.A. at 56-2-2381-7000. You can also contact your pesticide distributor or university extension specialist to report resistance.

INFORMATION

This material may be applied with ground equipment or aircraft depending upon local conditions. When growing crops for processing, consult the processor before applying sulfur.

For control of Mites, use higher rates shown under conditions of high pest pressure. For control of Fungi, use higher rates shown when weather conditions indicate a potential for increased pest pressure.

CROPS AND APPLICATION RATES

ALFALFA:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Powdery Mildew, Two Spotted Mite, Pacific Mite and Strawberry Mite (Atlantic Mite)	30 to 50	Apply when disease first appears and repeat at 14 day intervals or as necessary.

ALMONDS, CHERRIES, NECTARINES, PEACHES, PLUMS AND PRUNES:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Brown Rot Blossom Blight, Peach Canker, Shothole, Silver Mite, Stem Blight, Cherry Leaf Spot, Prune Rust and Powdery Mildew	35 to 60	Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals.

APPLES and PEARS:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Scab and Powdery Mildew	40 to 50	Apply in pre-bloom, bloom and post-bloom periods.
Use Restriction: On pears DO NOT apply after the fruit begins to turn down.		

ARTICHOKEs (GLOBE, JERUSALEM, CHINESE):

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Leaf Spot	25 to 35	Begin when disease first appears and repeat at 7 to 10 day intervals through season.

ASPARAGUS:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Rust and Asparagus Mite	40 to 50	Dust after cutting stops, irrigate, cultivate, then apply the sulfur. Repeat at 7 to 10 day intervals through season.

AVOCADOS:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Mite Suppression	40 to 50	Apply when mites first appear and repeat as necessary

BARLEY AND OATS:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Rust	25 to 40	Apply at first sign of disease. Repeat at 7 to 10 day intervals.

BEANS (DRY, GREEN) AND PEAS (FIELD, GREEN, AND SUGAR SNAP):

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Powdery Mildew, Leaf Spot Rust, Spider Mites and Thrips	15 to 35	Apply when disease first appears and repeat as necessary. Certain varieties may be injured by applications of sulfur. For these crops, it is advised that a test be conducted on a small number of plants to check for sulfur sensitivity prior to application to the entire field.

BEETS:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Rust	20	Apply at first signs of disease. Repeat at 7 to 10 day intervals.

BLACKBERRIES, BLUEBERRIES, CURRANTS, DEWBERRIES, GOOSEBERRIES, HUCKLEBERRIES, LOGANBERRIES:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Powdery Mildew	20 to 30	Begin when new shoots are 6 inches long and before blossoms open. Repeat at 10 day intervals or as necessary.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, COLLARDS, KALE, RUTABAGAS, TURNIPS:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Powdery Mildew	20 to 25	Begin at first sign of disease. Repeat at 7 to 10 day intervals or as necessary.

CARROTS:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Petrobia Mite, Powdery Mildew and Rust	20 to 50	Begin at first sign of disease. Repeat as necessary.

CELERY:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Early and Late blight	6 to 12	Begin as soon as plants emerge. Repeat at 7 to 10 day intervals or as necessary.

CITRUS (CITRON, GRAPEFRUIT, KUMQUAT, LEMON, ORANGE, PUMMELO, TANGELO, TANGERINE, LIME):

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Citrus Thrips, Citrus Flat Mite, Yuma Spider Mite, Six-Spotted Mite and Citrus Rust Mite	100 to 125	Apply when pests first appear and repeat as necessary.
Use Restriction: DO NOT use within 21 days of an oil spray treatment.		

CLOVER:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Powdery Mildew, Rust, Spider Mites	15 to 25	Begin when crop is in bud stage or at first sign of disease or mites. Repeat at 7 to 10 day intervals or as necessary.

COTTON:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Strawberry (Atlantic) Mite	30 to 40	Use when mites first appear and repeat as necessary.

COWPEAS:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Rust	5 to 25	Begin soon after seedlings emerge. Repeat at 7 to 10 day intervals through season.

CUCUMBERS, PUMPKINS, AND SQUASH:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Powdery Mildew and Scab	10 to 35	Begin when first true leaves appear and repeat at weekly intervals. Important: Certain varieties may be injured by applications of sulfur. For these crops, it is advised that a test be conducted on a small number of plants to check for sulfur sensitivity prior to application to the entire field.

DATES:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Banks Grass Mites	50 to 125	Apply by ground or air. Use the higher rates by air. Thorough coverage is essential for good control. Begin at first sign of disease. Repeat as necessary.

EGGPLANTS:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Powdery Mildew	20 to 35	Begin when first true leaves appear. Repeat at weekly intervals.

FIGS:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Fig Rust Mite	40 to 50	Dust when damage to foliage (russeting) appears. A second application may be necessary.

FLAX:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Powdery Mildew	20 to 30	Begin at first sign of disease. Repeat at 7 to 10 day intervals or as necessary.

GARLIC:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Downy and Powdery Mildew	20 to 35	Begin at first sign of disease. Repeat as necessary.
Petrobia Mite	25 to 30	

GRAPES:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Powdery Mildew and Rust	10 to 40	Make first application when shoots are 6 to 8 inches long. Make a second application when shoots are 12 to 15 inches long and third about 14 days later. Repeat applications as necessary to protect new growth. Important: Concord or Labrusca type grapes may be injured by sulfur.

GRASS SEED CROPS:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Timothy Mites	30	Apply when mites first appear and repeat as necessary.

HOPS:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Spider Mite	30 to 50	Apply when mites first appear and repeat as necessary.

LETTUCE:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Powdery Mildew	10 to 15	Begin at first sign of disease. Repeat as necessary.

MELONS (CANTALOUPE, MUSKMELON, WATERMELON):

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Powdery Mildew and Scab	15 to 50	Treat when disease first appears.
Spider Mites.	30 to 50	Important: Certain varieties may be injured by applications of sulfur. For these crops, it is advised that a test be conducted on a small number of plants to check for sulfur sensitivity prior to application to the entire field.

OLIVES:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Olive Mite	50 to 70	Begin at first sign of disease. Repeat as necessary.
Use Restriction: DO NOT use sulfur in hot weather as damage may result to crop and foliage.		

ONIONS:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Petrobia Mites, Botrytis Blight and Powdery Mildew	20 to 50	This product may be applied by air or ground equipment. Apply when disease first appears and repeat at 14 day intervals or as necessary.

PEANUTS:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Leaf Spot and Powdery Mildew	20 to 25	Begin when plants are 40 days old or when first spots appear on lower leaves. Repeat at 10 to 14 day intervals and after rains.
Rust	20 to 30	Apply when disease threatens. Repeat at 7 to 10 day intervals.

PEPPERS (BELL, CHILI, AND SWEET):

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Powdery Mildew	20 to 35	Begin before disease appears. Repeat at weekly intervals.

PISTACHIOS:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Mites	50	Make by ground or air application to bearing or non-bearing trees up to the day of harvest. Begin at first sign of disease. Repeat as necessary.

POMEGRANATES:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Mites	50	Apply by ground or air when mites first appear and repeat as necessary to maintain control. There is no time

		limitation between applications and harvest.
--	--	--

POTATOES:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Powdery Mildew	5 to 15	Apply before disease appears. Repeat at 7 to 10 day intervals or as necessary.

QUINCE:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Brown Rot, Powdery Mildew, Scab	35 to 60	Begin before diseases are expected to appear. Repeat at 7 to 10 day intervals or as necessary.

SOYBEANS:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Leaf Spot and Powdery Mildew	8 to 15	Apply at first sign of disease. Repeat at 7 to 10 day intervals.

SPINACH:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Powdery Mildew and Rust	8 to 30	Apply when disease first appears. Repeat at 7 to 10 day intervals.

STRAWBERRIES, RASPBERRIES and BOYSENBERRIES:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Powdery Mildew	3 to 50	Apply when disease first appears and repeat as necessary.

SUGAR BEETS:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Powdery Mildew, Two-Spotted Mite, Pacific Mite and Strawberry Mite (Atlantic Mite)	35 to 50	Apply when disease first appears and repeat at 14 day intervals or as necessary.

TOMATOES:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Tomato Russet Mite and Powdery Mildew	20 to 50	Begin at first sign of disease. Repeat as necessary.

WALNUTS:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Spider Mites	40 to 50	Apply throughout the season as needed. Repeat if necessary. Check tolerance prior to using. Consult local Agricultural Authorities as to the effectiveness of sulfur on Spider Mites.

WHEAT:

Disease/Pest	Application Rate (lbs. a.i./acre)	Use Instructions
Powdery Mildew	35 to 50	Apply when disease first appears and repeat at 14 day intervals or as necessary.

ORNAMENTALS

Use 20 to 40 lbs. a.i. per acre. Dust for complete coverage on both sides of leaves, applying when disease first appears and repeat at 5 to 10 day intervals and after rain, or as necessary. Keep new growth protected. AVOID APPLICATION TO SULFUR SENSITIVE PLANTS.

Ornamentals	Disease/Pest
Asters, Carnations:	Powdery Mildew and Rust
Calendulas, Cedars, Cherry Laurels, Chrysanthemums, Hollyhock, Juniper, Smilax, Spruce:	Leaf Spot and Powdery Mildew
Clematis, Columbine, English Ivy, Laurel, Petunia, Sage:	Leaf Spot.
Cosmos, Crepe Myrtle, Dahlia, Delphinium, Dogwood, Euonymus, Gladiolus, Golden Fleece, Hibiscus, Holly, Lady's Mantle, Ligustrum, Lilac, Poplars, Silver Vine, Spirea, Sunflower, Sweet Pea, Verbena, Violets, Willow, Zinnias:	Powdery Mildew
Roses:	Black Spot, Powdery Mildew, Rust

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store unused product in a cool, dry, secure area in the original container only.

Pesticide Disposal: Pesticide spray mixture or rinse water that cannot be used in accordance with label instructions must be disposed of according to applicable Federal, state or local procedures. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Non-refillable bag. Do not reuse or refill this bag. Offer for recycling, if available. If recycling is not available, dispose of bags in a sanitary landfill or by incineration if allowed by state and local authorities.

WARRANTY STATEMENT

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on this label when used in accordance with directions under normal conditions of use; but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions not reasonably foreseeable to seller; to the extent consistent with applicable law, the buyer assumes the risk of any such use.

Batch Code / Lot Number: _____
Net Weight: _____ lbs