STRATION DIVISION (75-767) NASHINGTON: DC 20460 NOTICE OF PESTICIDE: (Under the Federal Insecticide, Fungicide, BMP 123 (32 LC) and Rodenticide Act, as amended) NAME AND ADDRESS OF REGISTRANT (Include ZIP code) Becker Microbial Products, Inc. 9464 N.W. 11th Street Plantation, FL 33322 NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number. On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered unde the Federal Insecticide, Fungicide, and Rodenticide Act. A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith; Registration is in no way to be construed as an indersement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others. This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(C) provided that you: 1. Submit the studies listed below by December 21, 1994. Guideline Reference BEST AVAILABLE COPY Title of Study Number 154A-16 Avian Oral 30 Day Freshwater Fish Toxicity/ 154A-19 Pathogenicity Testing Freshwater Aquatic Invertebrate 154A-20 Toxicity/Pathogenicity Testing Honey Bee Toxicity/Pathogenicity Test 154A-24 Analysis of Samples<sup>1</sup> 154A-13 The Avian Oral and Freshwater Fish study requirements can also be satisfied by submitting data waiver requests which are reviewed and found satisfactory by the Agency on or before 12/21/94. <sup>1</sup>See page 40 of the <u>Guidance for the Reregistration of</u> Pesticide Products Containing Bacillus thuringiensis as the Active Ingredient. You must analyze for lepidopteran active toxin as per ATERSTERNIE APPLICABLE SIGNATURE OF APPROVING OFFICIAL

- 2. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.
- 3. Submit production information (pounds or gallons produced) for this product for the fiscal years in which this product is conditionally registered in accordance with FIFRA section 29. The fiscal year begins October 1 and ends September 30. The production information will be submitted to the Agency no later than November 15, following the end of the preceding fiscal year. This information should be submitted to:
  - U.S. Environmental Protection Agency Office of Pesticide Programs (H7504C) Document Processing Desk (RSB) 401 M Street SW., CM #2 Washington, DC 20460
- 4. Make the following label changes before you release the BMP 123 (32 LC) product for shipment.
- a) In the ingredients statement, delete "Lepidopteran active toxins" since you base your active ingredient percentage on bioassay rather than  $\delta$ -endotoxin percentage. Also, add "strain HD-263" after "kurstaki."
- b) Add the heading "Hazards to Humans and Domestic Animals" above the block of text that starts "CAUTION: Harmful if inhaled..." and add the signal word CAUTION below the KEEP OUT OF REACH OF CHILDREN statement. All point sizes must comply with 40 CFR 156.10.
- c) Comply with PR Notice 87-1 by adding "Do not apply this product through any type of irrigation system." or per section XI.(C)(1) of the Notice add the exact language specified in sections IV. through IX. of the Notice. Chemigation instructions beyond these must be submitted via an amendment application.
- d) Delete "highly selective" and the "BENEFICIAL INSECTS:" heading and associated text since nontarget data substantiating these claims have not yet been submitted. Claims for safety to nontarget organisms must be supported by test data for specific organisms and life stages listed on the label.

- e) Place a STORAGE AND DISPOSAL heading above your storage and disposal block and add the statement "Do not contaminate water, food, or feed by storage or disposal." above the storage line of text.
  - f) List your full establishment number on the label.
  - g) Add "EPA REGISTRATION NUMBER: 62637-4" to the label.
- h) Per PR Notice 82-1, replace the statement "Read and, follow all precautions and restrictions on ovicide labels." with "Use of the resulting tank mix must be in accordance with the most restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing."
- 5. Submit a revised Confidential Statement of Formula that modifies the active ingredient portion by deleting "Lepidopteran active toxin" since you base your active ingredient percentage on bioassay rather than  $\delta$ -endotoxin percentage. The active ingredient percentage of 1.6% based on (8000 IU/mg) / (500,000 IU/mg) x 100% should remain the same.
- 6. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

Note that this acceptance of your product does not relieve you of your obligation to comply with the Worker Protection Standard. If any of your products are covered by this standard, you are required to submit, and receive the Agency's approval by April 21, 1994, a revised label reflecting the required label statements of 40 CFR 156, published in the FEDERAL REGISTER on August 21, 1992 (57FR 38102). Further guidance will be issued. According to 40 CFR 156, subpart K: "No product to which this subpart applies shall be distributed or sold without amended labeling by any registrant after April 21, 1994. Please call 1-800-777-2185 if you have any labeling questions regarding the Worker Protection Standard.

The Agency will require a change in the labeling regarding Endangered Lepidopteran Species when similar products are required to do so.

Per your representative Christine Dively's letter of

BEST ATTACK

5/20/93, which certified that Becker Microbial Products, Inc. is willing to voluntarily cancel this registration on 5/21/96, this registration will expire on 5/21/96.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA \$ 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records

Sincerely,

. Phil Hutton

Product Manager 18

Insecticide-Rodenticide Branch

\*\*Registration Division (H7505C)

### Aqueous Suspension

ACTIVE INGREDIENT: Bacillus thuringiensis subspecies kurstaki, 8,000 International Toxic Units (ITU) per milligram (Equivalent to 32 Billion ITU/gallon; 8.45 Billion ITU/liter) Lepidopteran active toxins 1.60% Potency units should not be used to adjust use rates beyond those specified in the Directions for Use
Section. INERT INGREDIENTS:

### KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes or clothing Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remo : contaminated clothing and wash before reuse.

### STATEMENTS OF PRACTICAL TREATMENT:

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF INHALED: Remove victim to fresh air.

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS: Do not contaminate water when disposing equipment washwaters.

BMP 123 (32 LC) is a highly selective microbial larvicide effective against lepidoptera in a variety of habitats.

BENEFICIAL INSECTS: Honey bees which feed in areas treated with BMP 123 (32 LC) are not harmed by the product. BMP 123 (32 LC) does not affect the activities of parasitic and predaceous arthropods in integrated pest management programs.

DIRECTIONS FOR USE: It is a violation of federal law to apply this product in a manner inconsistent with its labeling.

BMP 123 (32 LC) is a highly selective, microbial larvicide for use against the listed lepidopterous insects. Close scouting and early attention to insect infestations are recommended. BMP 123 (32 LC) must be ingested to be effective. The following directions should always be followed:

- 1. Larvae must be actively feeding on exposed plant parts.
- 2. Applications should be made when larvae are in the early instars before extensive crop damage occurs. For control of mixed insect populations or overlapping generations, use BMP 123 (32 LC) in combination with another approved insecticide.
- 3. Thorough spray coverage with BMP 123 (32 LC) is required for consistent control. Spray nozzles and volume applied will vary depending on crop and pest. Under heavy pest pressure, higher rates, increased gallonage, and shorter spray intervals will improve crop coverage and pest control.

AUG 05 1993

Under the Fed ral Investicide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

- 4. Repeat applications at 3-14 day intervals to maintain control. This will depend on weather, plant growth rate, and insect activity. Single spray applications should be made when egg hatch is essentially complete and before extensive crop damage occurs.
- 5. In applications on hard to wet crops such as cole crops, a spreader/sticker approved for use on growing and harvested crops, should be added to improve coverage and weathering properties.
- 6. Unless otherwise indicated use at least two (2) gallons of water per acre except in Californai where 5-10 gallons is typically used.
- 7. BMP 123 (32 LC) may be applied as a pre and post harvest treatment to raw agricultural commodities.
- BMP 123 (32 LC) may be applied in conventional aerial and ground equipment with sufficient water to provide thorough coverage.
   Approved spreader/stickers should be added to the mix tank after

   BMP 123 (32 LC) has been mixed with the appropriate amount of water.


#### APPLICATION RATE

Crop	Pest	Ounces/Acre		
Vegetables and Commercial Crops of the following groups:	Loopers Hornworms	16 to 64 16 to 32		
Root and Tuber Leaves of Root and Tuber Bulb	Imported Cabbageworms Diamondback	16 to 32		
Leafy Brassica (Cole Leafy)	Moth Green Cloverworm	32 to 64 16 to 32		
Legume Foliage of Legume	Webworm Armyworms*	16 to 32 32 to 64		
Fruiting Cucurbit	Saltmarsh Caterpillar	32 to 64 32 to 64		
(Melon, squash, cucumber, etc.)	Cutworms Heliothis Spp.** Pickleworm	32 to 64 16 to 32		
	Melonworm Rindworm	16 to 32		
	Complex***	32 to 64		

\*BMP 123 (32 LC) may be used to suppress 1st and 2nd instar armyworm larvae when populations are light and full coverage, ground applications are used.

\*\* Tobacco Budworm, Iomato Fruitworm, Corn Earworm. On tomatoes apply weekly in a preventative program.

\*\*\* Apply weekly in a preventative program.



Crop	Pest	Ounces/Acre
Sunflowers	Loopers	32 to 64
Peanuts	Green Cloverworms	16 to 32
Safflower	<b>Velvetbean</b>	
Mint '	Caterpillar	16 to 32
Canola	Podworm*	32 to 64
Rape	Armyworms	32 to 64
Cereal Grains	Diamondback	
Amaranth	Moth	32 to 64
Jojoba	Saltmarsh	
Okra	Caterpillar	32 to 64
Pawpaw	Hornworms	16 to 32
Lentils .	Heliothis spp.***	32

\* RMP 123 (32 LC) can be used to suppress podworm in an integrated pest management program.

\*\* Suppression Only.

Crop	Pest	Ounces/Acre
Artichokes	Artichoke Plume	
	Moth	32 to 80
	Armyworms	32 to 96
	Loopers	16 to 80
Cotton	Tobacco Budworm*	32 to 64
	Cotton Bollworm*	32 to 64
	Loopers	32 to 64
	Cotton Leaf	
	Perforator	32 to 64
	Cotton Leafworm	32 to 64
	Saltmarsh	
	Caterpillar	32 to 64
	Armyworms	32 to 96

\*Use in conjunction with regular scouting in an integrated pest management program when beneficial insects are present. Applications should be made against light to moderate populations of 1st and 2nd instar larvae. Repeat applications at 3 to 5 day intervals as required and as long as population suppression is satisfactory. For added control of budworms and bollworms, tankmixing of BMP 123 (32 LC) with an approved ovicide is recommended. Read and follow all precautions and restrictions on ovicide labels.

# BEST AVAILABLE COPY

Crop	Pest	Ounces/Acre
Tobacco	Tobacco Budworm Hornworms	32 to 64 16 to 32
•	Loopers	32 to 64

BMP 123 (32 LC) may be applied in conjunction with sucker control products at the recommended label rates. Observe label precautions for all products used. Apply BMP 123 (32 LC) as necessary to maintain control.

	Crop	Pest	Ounces/Acre
	Greenhouse		,
$\mathcal{C}$	Flowers	Loopers	16 to/32
	Bedding Plants	Tobacco Budworm	16 to 32
	Ornamentals	Omnivorous Looper	16 to 32
)	(Ground Only)	Omnivorous	20 00 02
,	(Ground Only)	Leafroller	16 to 32
		Diamondback Moth	32 to 64
		Armyworms	32 to 128
		Ello Moth	52 to 120
(		(Hornworm)	16 to 32
ν.		Io Moth	16 to 32
		Oleander Moth	16 to 32
		Azalea Caterpillar	· · · · · · · ·
		Azarea Caterprirar	10 00 32
			•
	Crop	Pest	Ounces/Acre
<u></u>	<b>↑</b> Pastures	Loopers	16 to 64
	Cereal Grain	Alfalfa Caterpillar	· · · · · · · · · · · · · · · · · · ·
	Forage	European Skipper	8 to 16
•	Fodder	Armyworms	64 to 96
,	Straw	,	
	Grass and nongrass		3. 7.
	commodities.	,	**
	Rangeland	Range Caterpillar*	16 to 32
·	*Apply in a minimum of 1/2	to 1 gallon of water per	acre.
	Land of		
	A.Turf	Sod Webworm	64 to 128

bisi mailing capy

Crop	Pest	Ounces	/Acre	
		Ground *	Air**	
Forest	Gypsy Moth	32 to 128	32 to 128	
Shade Trees	Eastern Tent			
Schrubs	Caterpillars	16 to 32	16 to 32	
Ornamentals	Spruce Budworm	32 to 96	32 to 96	
Sugar Maple Trees	Douglas Fir			
-	Tussock Moth	16 to 64	16 to 64	
•	Western Tussock			
	Moth	16 to 32	16 to 32	
•	Browntail Moth	16 to 32	16 to 32	
	Bagworm	8 to 16	8 to 16	
	Spring and Fall	•	•	
	Cankerworms	16 to 32	16 to 32	
	Elm Spanworm	16 to 32	16 to 32	
	Fall Webworm	16 to 32	16 to 32	
	California			
	Oakworm	16 to 32	16 to 32	
	Redhumped			
	Caterpillar	16 to 32	16 to 32	
	Pine Butterfly	32 to 48	32 to 48	
	Blackheaded			
	Budworm	32 to 48	32 to 48	
	Jack Pine Budwo		32 to 64	
	Mimosa Webworm	16 to 32	-	
	Saddle Prominen	=		
	Caterpillar	16 to 32	16 to 32	
	Saddleback	10 +- 20	10 +- 20	
	Caterpillar	16 to 32	16 to 32	
	Fruittree	10 4- 20		
	Leafroller	16 to 32	-	
	Forest Tent	10 4- 00	10 4- 00	
	Caterpillar	16 to 32	16 to 32	

\*Ground applications are made at the rate of 100 gallons per acre for hydraulic sprayers. For mist blowers mix the applicable amount in ten gallons of water.

\*\*Use 1/2 to ten gallons of water depending on the type and density of tree.



# Fruits and Nuts

Crop	Pest	Ounces/Acre		
Grapes				
Small Fruits	Loopers	32 to 64		
Berries	Grape Berry Moth*	32 to 64		
Hops	Grapeleaf Skeletonizer*	32 to 64		
	Grape Leaffolder	32 to 64		
Small Fruits	Saltmarsh Caterpillar*	32 to 64		
	Achema Sphinx Moth*	16 to 32		
•	Omnivorous Leafroller*	32 to 64		
	Blueberry Leafroller	16 to 64,		
	Cherry Fruitworm	16 to 32 -		
	Fruittree Leafroller	16 to 64		
	Gypsy moth	32 to 64		
	Green Fruitworm	16 to 32 🕶		
	Orange Tortrix	32 to 64		
	Grape Leafroller	16 to 64::		
	Spotted Cutworm	32 to 64		

\*Application by ground only.

Crop	Pest	Ounces/Acre	• • •
		Ground	A!r
Stone Fruits	Tent		
Nut Trees	Caterpillars	32 to 64	32 to 64
Pomegranates	Cankerworms	16 to 32	16 to 32
_	Fall Webworm	32 to 64	32 to 64
•	Walnut		,
	Caterpillar	16 to 64	16 to 64
•	Redhumped		(4)
• ·	Caterpillar	32 to 64	32 to 64
	Omnivorous		
	Leafrolier	32 to 64	
	Citrus Cutworm	32 to 64	32 to 64
	Filbert		•
 &	Leafroller	32 to 64	32 to 64
	Filbert Webworm	32 to 64	32 to 64

BEST AVAILABLE COPY

Crop	Pest	Ounces/Acre	9
		Ground	Air
Pome Fruits*	Gypsy Moth	32 to 96	3: to 96
	Cankerworms Tent	16 to 32	16 to 32
	Caterpillars Tufted Apple	32 to 64	32 to 64
	Budmoth Variegated	32 to 64	
•	Leafroller Redbanded	32 to 64	
	Leafroller	32 to 64	
Citrus**	Fruittree		
	Leafroller	32 to 64	'
	Citrus Cutworm	32 to 64	32 to 64
	Orangedog	16 to 32	
Avocado	Loopers	32 to 128	32 to 128
	Orange Tortrix Omnivorous	32 to 64	32 to 64
	Leafroller	32 to 64	32 to 64
	Amorbia	64	


	Crop	Pest				Ounces/Acre	
			G	rou	nd		Air
	Tropical				- <del></del>		<del></del>
	Fruits***	Hornworm	32	to	128		<b>-</b> →
)	Papayas	Leafrollers	32	to	128		
	Mangoes	Omnivorous Looper	32	to	128		
	Kiwi	Loopers	32	to	128		
•	Persimmons	_					
	Pomegranate						
	Bananas***	Banana Skipper	24	to	64	,	

\*Apply BMP 123 (32 LC) in 450-450 gallons/acre for pest suppression in pest management programs.

\*\*Apply in 100 to 400 gallons per care.

\*\*\*Apply at least 200 gallons per acre.

\*\*\*\*Hawaii Only, Ground Equipment Only, Apply to the Point of Runoff.



Storage: Store in a cool dry place.

<u>Pesticide Disposal</u>: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent), then puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned stay out of smoke. Do not reuse container.

NOTICE TO USER: Seller makes no warranty, express or implied, of merchantability, fitness or otherwise concerning the use of this product other than as indicated on the label. User assumes all risks of use, storage or handling not in strict accordance with label instructions.

BECKER MICROBIAL PRODUCTS, INC. 9464 N.W. 11th Street Plantation, Florida 33322

EPA REGISTRATION NUMBER: 62637-