SEPA Environmental Protection Washington, DC 204	on Agency	Registration Amendment Other	OPP Identifier Number
1. Company/Product Number	2. EPA Product Manager		oposed Classification
62637-4	Janet Andersen	j	· —
4. Company/Product (Name) BMP 123 (32 LC)	PM# Acting Division		None Restricts
5. Name and Address of Applicant (Include ZIP Code) Becker Microbial Products, Inc. 9464 N.W. 11th Street Plantation, FL 33322 Check if this is a new address	(b)(i), my product is si to: EPA Reg. No	milar or identical in co	
	Product Name Section - II		
Amendment - Explain below. Resubmission in response to Agency letter lated X Notification - Explain below.	Final printed la Agency letter of "Me Too" Appl	ication.	
Explanation: Use additional page(s) if necessary. (For section Notification - PR Notice 95-3; Reduc		Is for low risk	pesticides
	Section - III		
Material This Product Will Be Packaged In:	·- 	 	
Child-Resistant Packaging Yes* No No No No No No. per Unit Packaging Ves No. per Unit Packaging wgt. container	Water Soluble Packaging Yes No If "Yes" Package wgt No. per Container	2. Type of Container Metal Plastic Glass Paper Other (5	Specify)
3. Location of Net Contents Information 4. Size(s) Re Label Container	tail Container 5.	Location of Label Direction On Label On Labeling accom	
6. Manner in Which Label is Affixed to Product Lithor Paper Stend	graph Other _ glued illed		
	Section - IV		
1. Contact Point (Complete items directly below for identification)			***************************************
Name Pamela Noyes, Jellinek, Schwartz & Connolly, Inc.	Regulatory Specialis	st (703)	ne Nq. (Include Area Code 312–8559
Certific. I certify that the statements I have made on this form and lacknowledge that any knowingly false or misleading stated both under applicable law.	d all attachments thereto are true, a		6. Date Application . Rankined , (Stamped)
2. Signature	3. Title Authorized Repre Becker Microbial		
4. Typed Name / () Pamela Noyes	5. Date 8-10-95		

BMP 123 (32 LC) BIOLOGICAL LARVICIDE AQUEOUS SUSPENSION

ACTIVE INGREDIENT: Bacillus thuringiensis subspecies kurstaki strain BMP 123....1.60% POTENCY: 8,000 International Units (IU) per milligram (Equivalent to 32.0 Billion IU

per gallon: 8.45 Billion IU / Liter).

Potency units should not be used to adjust rates beyond those specified in the Directions for Use Section.

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENTS OF PRACTICAL TREATMENT

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation

persists.

IF INHALED: Remove victim to fresh air.

IF IN EYES: Flush eyes with plenty of water. Get medical

attention if irritation persists.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist.

Personal Protective Equipment (PPE): Applicators and other handlers must wear:

- * Long-sleeved shirt and long pants
- Waterproof gloves
- * Shoes and socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS:

Users should wash hands before eating, drinking, chewing gum, using tobacco or using toilet.

ENVIRONMENTAL HAZARDS STATEMENT: Do not contaminate water when disposing equipment wash waters.

DIRECTIONS FOR USE:

It is a violation of federal law to apply this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticidé regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.



Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

BMP 123 (32 LC) is a microbial larvicide for use against the listed lepidopterous insects. Close scouting and early attention to insect infestations are recommended. BMP 123 (32 LC) must be ingested to be effective. The following directions should always be followed:

- 1. Larvae must be actively feeding on exposed plant parts.
- 2. Applications must be made when larvae are in the early instars before extensive crop damage occurs. For control of mixed insect populations or overlapping generations, use BMP (32 LC) in combination with another approved insecticide.
- Thorough spray coverage with BMP 123 (32 LC) is required for consistent control.
 Spray nozzles and volume applied will vary depending on crop and pest. Under heavy pressure, higher rates, increased gallonage, and shorter spray intervals will improve crop coverage and pest control.
- 4. Repeat applications at 3-14 day intervals to maintain control. This will depend on weather, plant growth rate, and insect activity.
- In applications on hard to wet crops such as cole crops, a spreader sticker approved for use on growing and harvested crops, should be added to improve coverage and weathering properties.
- 6. Unless otherwise indicated use at least two (2) gallons of water per acre except in California where five (5) to ten (10) gallons are typically used.

- 7. BMP 123 (32 LC) may be applied as a pre and post harvest treatment to raw agricultural commodities.
- 8. Do not apply this product through any kind of irrigation system.

BMP 123 (32 LC) may be applied in conventional aerial and ground equipment with sufficient water to provide thorough coverage. Approved spreader/stickers should be added to the mix tank after BMP 123 (32 LC) has been mixed with the appropriate amount of water.

APPLICATION RATE

Crop	Pest	Ounces / Acre
Vegetables and commercial	Loopers	32 to 64
crops of the fellowing groups:	Hornworms	32 to 64
Root and Tuber	Imported Cabbageworms	32 to 64
Leaves of Root and Tuber	Diamondback moth	32 to 96
Bulb	Green Cloverworm	32 to 64
Leafy	Webworm	32 to 64
Brassica (Cole Leafy)	Armyworms*	32 to 128
Legume	Saltmarsh Caterpillar	32 to 64
Foliage of Legume	Heliothis (Helicoverpa) Spp.**	32 to 128
Fruiting	Cutworms	32 to 64
Cucurbit (Melons, Squash,	Pickleworm	32 to 64
Cucumbers, etc.)	Melonworm	32 to 64
• •	Rindworm Complex***	32 to 96

- * BMP 123 (32 LC) may be used to suppress 1st and 2nd instar armyworm larvae when populations are light and full coverage ground applications are used.
- ** Tobacco Budworm, Tomato Fruitworm, Com vorm. On tomatoes apply weekly in a preventative program.
- *** Apply weekly in a preventative program.

Crop	Pest	Ounces / Acre
Sunflowers	Loopers	32 to 64
Peanuts	Green Cloverworms	32 to 64
Safflower	Velvetbean Caterpillar	32 to 64
Mint	Podworm*	32 to 64
Canola	Armyworms*	32 to 128
Rape	Diamondback Moth	32 to 96
Cereal Grains	Saltmarsh Caterpillar	32 to 64
Amaranth	Hornworms	16 to 32
Jojoba	Heliothis (Helicoverpa) spp.	32 to 128
Okra	, , , , , ,	
Pawpaw		

- * BMP 123 (32 LC) can be used to suppress podworm and armyworms in an integrated pest management program.
- ** Suppression only.

Lentils

Crop	Pest	Ounces / Acre
Artichokes	Armyworms	32 to 128
	Loopers	16 to 80
Cotton	Tobacco Budworm*	32 to 128
	Cotton Bollworm*	32 to 128
	Loopers	32 to 64
	Cotton Leaf Perforator	32 to 64
	Cotton Leafworm	32 to 64
	Saltmarsh Caterpillar	32 to 64
	Armyworms	32 to 128

^{*} Use in conjunction with regular scouting in an integrated pest management program when beneficial insects are present. Applications should be made against light to moderate populations of 1st and 2nd instar larvae. Repeat applications at 3 to 5 day intervals as required and as long as population suppression is satisfactory. For added control of budworms and bollworms tankmixing of BMP 123 (32 LC) with an approved ovicide is recommended. Use of the resulting tank mix must be in accordance with the most restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product can not be mixed with any product containing a label prohibition against such mixing.

Crop	Pest	Ounces / Acre
Tobacco	Tobacco Budworm Homworms	32 to 128 16 to 32
	Loopers	32 to 64

BMP 123 (32 LC) may be applied in conjunction with sucker control products at the recommended label rates. Observe label precautions for all products used. Apply BMP 123 (32 LC) as necessary to maintain control.

Crop	Pest	Ounces / Acre	
Greenhouse	Loopers	16 to 32	
Flowers	Tobacco Budworm	16 to 32	
Ornamentals	Omnivorous Looper	16 to 32	
Bedding Plants	Omnivorous Leafroller	16 to 32	
(Ground Only)	Diamondback Moth	32 to 64	
•	Armyworms	32 to 128	
	Ello Moth (Hornworm)	16 to 32	
	lo Moth	16 to 32	
	Oleander Moth	16 to 32	
	Azalea Caterpillar	16 to 32	
Crop	Pest	Ounces / Acre	•
Pastures, Cereal Grain,	Loopers	32 to 64	,
Forage, Fodder, Straw,	Alfalfa Caterpillar	32 to 32	
Grass and nongrass commodities	European Skipper	8 to 16	,
-	Armyworms	64 to 128	•
Rangeland	Rangeland Caterpillars*	16 to 64	

^{*} Apply in a minimum of 1/2 to 1.0 gallon per acre.

Crop	Pest	Ounces / Acre
Turf	Sod Webworm	64 to 128

Crop	<u>Pest</u>	<u> </u>	/ Acre
		Ground*	Air^*
Forest	Gypsy Moth	32 to 128	32 to 128
Shade Trees	Eastern Tent Caterpillars	16 to 32	16 to 32
Shrubs	Spruce Budworm	32 to 96	32 to 96
Ornamentals	Douglas Fir Tussock Moth	16 to 64	16 to 64
Sugar Maple Trees	Western Tussock Moth	16 to 32	16 to 32
• ,	Browntail Moth	16 to 32	16 to 32
	Bagworm	8 to 16	8 to 16
	Spring and Fall Cankerworms	16 to 32	16 to 32
	Elm Spanworm	16 to 32	16 to 32
	Fall Webworm	16 to 32	16 to 32
	California Oakworm	16 to 32	16 to 32
	Redhumped Caterpillar	16 to 32	16 to 32
	Pine Butterfly	32 to 48	32 to 48
	Blackheaded Budworm	32 to 48	32 to 48
	Jack Pine Budworm	32 to 64	32 to 64
	Mimosa Webworm	16 to 32	
	Saddle Prominent Caterpillar	16 to 32	16 to 32
	Saddleback Caterpillar	16 to 32	16 to 32
	Fruittree Leafroller	16 to 32	
	Forest Tent Caterpillars	16 to 32	16 to 32

^{*} Ground applications are made at the rate of 100 gallons per acre for hydraulic sprayers. For mist blowers mix the applicable amount in ten (10) gallons of water.

^{**} Use 1/2 to ten (10) gallons of water depending on the type and density of tree.

Crop	Pest	Ounces / Acre	
Grapes	Loopers	32 to 64	
Small Fruits	Grape Berry Moth*	32 to 64	
Berries	Grapeleaf Skeletonizer*	32 to 64	
Hops	Grape Leaffolder	32 to 64	
	Saltmarsh Caterpillar*	32 to 64	
	Achema Sphinx Moth*	16 to 64	
	Omnivorous Leafroller*	32 to 64	
	Blueberry Leafroller	16 to 64	
	Cherry Fruitworm	16 to 32	
	Fruittree Leafroller	16 to 64	•
	Gypsy Moth	32 to 64	•
	Green Fruitworm	16 to 64	•
	Orange Tortrix	32 to 128	
	Grape Leafroller	16 to €4	
	Spotted Cutworm	32 to €4 ·	٠
	Armyworms**	32 to 128	
	Tobacco budworm	32 to 128	

^{*} Application by ground only.

^{**} BMP 123 (32 LC) may be used to suppress 1st and 2nd instar armyworm larvae when populations are light and full coverage applications are used.

Crop	Pest	Ounces / Acre	
		Ground	Air
Otana Farita	Tank Oaks william	004-04	22 to 64
Stone Fruits	Tent Caterpillars	32 to 64	32 to 64
Nut Trees	Cankerworms	16 to 32	16 to 32
Pomegranates		32 to 64	32 to 64
	Walnut Caterpillar	16 to 64	16 to 64
	Redhumped Caterpillar		32 to 64
	Omnivorous Leafroller	32 to 64	
	Citrus Cutworm	32 to 64	32 to 64
	Filbert Leafroller	32 to 64	32 to 64
	Filbert Webworm	32 to 64	32 to 64
		Ounces / Acre	
Crop	Pest	Ground	Air
Pome Fruits*	Gypsy Moth	32 to 96	32 to 96
	Cankerworms	16 to 32	16 to 32
	Tent Caterpillars	32 to 64	32 to 64
	Tufted Apple Budmoth		OL 10 04
	Variegated Leafroller	32 to 64	
	Redbanded Leafroller	32 to 64	
Citrus**	Fruittree Leafroller	32 to 64	
	Citrus Cutworm	32 to 64	32 to 64
	Orangedog	16 to 32	
Avocado	Loopers	32 to 128	32 to 128
	Orange Tortrix	32 to 64	32 to 64
	Omnivorous Leafroller	32 to 64	32 to 64
	Amorbia	64	
Crop	Pest	Ounces / Acre	
		Ground	Air
Tropical			
Fruits***	Hornworm	32 to 128	
Papayas	Leafrollers	32 to 128	
Mangoes	Omnivorous Looper	32 to 128	
Kiwi	Loopers	32 to 128	
Persimmons Pomegranate	Loopers	32 10 120	
Bananas****	Banana Skipper	24 to 64	

Apply BMP 123 (32 LC) in 50-500 gallons / acre for pest suppression in pest management programs.

^{**} Apply in 100 to 400 gallons per acre.

^{***} Apply at least 200 gallons per acre.

^{****} Hawaii Only, Ground Equipment Only, Apply to the point of runoff.

STORAGE AND DISPOSAL Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool dry place.

Pesticide Disposal: Wastes resulting from use of this product may be disposed of on

site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or Equivalent), then puncture and dispose of in a sanitary

landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned stay out of smoke. Do not reuse container.

NOTICE TO USER: Seller makes no warranty, express or implied, of merchantability, fitness

or otherwise concerning the use of this product other than as indicated on the label. User assumes all risks of use, storage or handling not in strict

accordance with label instructions.

BECKER MICROBIAL PRODUCTS, INC. 9464 N. W. 11th Street Plantation, Florida 33322

EPA Registration Number: 62637-4

EPA Establishment Number: 68148-SC-001

Net Contents:

Lot Number: