Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

RESTRICTED USE PESTICIDE

May Injure (Phytotoxic) Susceptible, Non-Target Plants. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Commercial certified applicators must also ensure that all persons involved in these activities are informed of the precautionary statements.

Tordon* 22K

EPA Reg. No. 62719-6

(For use in areas west of the Mississippi River)

Directions for Wick or Carpet Roller Application

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Tordon 22K before applying. Carefully follow all precautionary statements and applicable use directions.
- In additions to limitations imposed by this supplemental labeling, use of Tordon 22K is subject to all use
 precautions and limitations imposed by the label affixed to the container for Tordon 22K.

Directions for Use

Tordon* 22K herbicide can be applied using wick or carpet roller equipment where drift presents a hazard to susceptible crops, surface waters and other sensitive areas. Mix 1 part of Tordon 22K with 2 parts of water to prepare a 33% solution.

Wick Application

Apply when weeds are actively growing and are above most desirable plants. For ironweed and goldenrod, best results are obtained with applications made prior to early bud stage. Wick applicator should be drained and cleaned after each use. Ropes should be changed when flow is reduced from wear, extended use, poor cleaning or intermittent use.

Carpet Roller Application

Apply to previously untreated plants less than 6 feet tall, and short enough to pass beneath the tractor without breaking off at the ground. Applications made during periods of extended drought conditions will not provide acceptable control. Do not mow, burn or otherwise disturb the treated plants during the remainder of the growing season. Operate carpeted rollers as close to the ground as possible without breaking the stems, but above the tallest grasses. Grasses growing adjacent to treated plants may exhibit temporary injury. Maximize herbicide deposition on foliage and stems and minimize drippage losses by rotating the carpeted roller at 30 to 40 rpm with the lower edge moving in the same direction as the • • direction of travel. Maintain the carpet sufficiently wetted to apply up to 1 gal/acre of herbicide-water • mixture to stands of average density (100 to 200 plants/acre), and up to 2 gal/acre in dense stands (390 to 400 plants/acre). Rewet rollers at regular intervals.

ACCEPTED

JUL 1 4 1999

Under the Federal Insecticide. Fungicide, and Rodenticide Act. as amended, for the pesticide registered under EPA Republic Learning

Carpet Roller - Use Recommendations

Brush Species	Amount of Tordon 22K	Specific Use Directions
mesquite regrowth	1 gal alone or with 2 qt Reclaim* herbicide to make 8 gal spray mixture	Include 1 ounce of a recommended agricultural surfactant per gallon of herbicide-water mixture (0.75% vol/vol). Apply from May through August, but preferably in May and June, when moisture availability is sufficient to allow normal plant growth. See "Broadcast Application" section for more details regarding optimum conditions for mesquite control.
huisache/blackbrush	f gal alone or with 2 to 3 qt of Reclaim to make 8 gal of spray mixture	Include 1 ounce of a recommended agricultural surfactant per gallon of herbicide-water mixture (0.75% vol/vol). Apply in the fall.

Maximum Use Rates: For control of other broadleaf weeds and woody plants, Tordon 22K may be applied broadcast at up to 1 quart per acre per annual growing season. Spot treatments may be applied at an equivalent broadcast rate of up to 2 quarts per acre per annual growing season, but not more than 50% of an acre may be treated. In noxious weed control programs, Tordon 22K may be applied at up to 2 quarts per acre per annual growing season as a broadcast treatment.

*Trademark of Dow AgroSciences LLC

M1A / Tordon 22K / Amend-Suppl / 11-20-98 D06-111-001 EPA-accepted __/_/_ Initial printing.

