


PM 19 62719-23

1041

U.S. Crop Production

# Supplemental Labeling



## DowElanco

DowElanco Quad IV, 9002 Purdue Road P O Box 681428 Indianapolis, Indiana 46268-1189 USA

### Lorsban\* 4E Insecticide (EPA Reg. No. 62719-23)

#### Application With Dry Bulk Fertilizers To Control Red Imported Fire Ant And Other Ant Species On Citrus Orchard Floors (For Distribution and Use Only In the State of Florida)

- ATTENTION**
- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
  - This labeling must be in the possession of the user at the time of application.
  - Read the label affixed to the container for Lorsban 4E insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
  - Use of Lorsban 4E according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Lorsban 4E.

#### Directions for Use

To control foraging ants and suppress mounds, apply Lorsban 4E to the orchard floor at the rate of 1 1/2 to 2 pints per acre. Retreat as needed. For best insect control, uniform coverage of the orchard floor is necessary. Dry fertilizers or fertilizer blends impregnated with Lorsban 4E may be used for treatment of citrus orchard floors. Uniform application of fertilizer is necessary to insure optimum results.

For impregnating Lorsban 4E on dry fertilizers, use a closed rotary drum mixer equipped with suitable spraying equipment. The spray nozzle should be positioned inside the mixer to provide uniform spray coverage of the tumbling fertilizer. Apply Lorsban 4E at the rate of 1 1/2 to 2 pints per acre to control ants in citrus orchard floors. The maximum concentration of Lorsban 4E to be added is 2 pints per 200 pounds of fertilizer and the minimum concentration is 2 pints per 500 pounds of fertilizer. At the higher concentration of Lorsban 4E, the fertilizer may not readily absorb all of the liquid. For a suitable free-flowing mixture, an absorptive powder such as Micro-Cel E should be added separately and uniformly to the fertilizer blend following addition of Lorsban 4E. Bulk fertilizers impregnated with Lorsban 4E should be applied immediately, not stored. All bulk containers should be tightly covered while the products are being transported and applied to reduce the chance of loss of Lorsban 4E via volatilization.


Compliance with any and all federal and state laws and regulations relating to the Lorsban 4E and fertilizer mixture is the responsibility of the person offering such mixture for sale or distribution.

Refer to label for Lorsban 4E for additional application methods for orchard floors.

Foliar applications of Lorsban 4E may be made in addition to the orchard floor treatments.

#### Restrictions:

- Do not apply last treatment within 28 days before harvest.
- Do not allow livestock to graze in treated areas.
- Do not apply more than 3 quarts per acre per season.



\*Trademark of DowElanco

Label Code) Approved \_\_\_/\_\_\_/\_\_\_  
Initial printing.

#### Amendments:

- 1) Added application with dry bulk fertilizer for control of red imported fire ants in citrus orchard floors.