

Supplemental Labeling

ACCEPTED

SEP 28 1992

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 62719-32



DowElanco

DowElanco

Quad IV, 9002 Purdue Road

P.O. Box 681428

Indianapolis, Indiana 46268-1189 USA

RESTRICTED USE PESTICIDE

Sale and Use Restricted to Certified Applicators or
Persons Under their Direct Supervision

Telone* II Soil Fumigant

(EPA Reg. No. 62719-32)

(For Distribution and Use Only in Idaho, Nevada, Oregon and Washington,
and in Modoc and Siskiyou Counties of California)

For Nematode and Wireworm Control In Potatoes

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Telone II before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Telone II according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Telone II.

Directions for Use

BEST AVAILABLE COPY

For control of Columbia root knot nematode, apply Telone II as described below at the rate of 20 gallons/acre.

For control of Northern root knot nematode, apply Telone II as described below at the rate of 15 to 18 gallons/acre. For high populations, use the higher recommended application rate.

For control of wireworms, apply Telone II as described below at the rate of 20 gallons/acre.

Application Methods and Equipment: Apply Telone II using either chisel (shank), Nobel (sweep) or modified ParaTill application equipment. Under most conditions, use either chisel equipment with ripper-type shanks or ParaTill equipment modified so that outlet spacing is evenly distributed under the tool bar. With chisel or ParaTill equipment, a shank spacing of 12 to 24 inches is recommended. Do not exceed a shank spacing of 24 inches. Outlet depth should be at least 18 inches below the final soil surface.

Nobel plow equipment should be used when shallow soils containing excessive live root material such as alfalfa, prevents the use of shank application. Nobel plow application should be made to a depth of at least 15 inches. Fumigant penetration may be limited if a plow pan exists below the depth of the Nobel blade. Plow-sole application is not recommended.

Note: Use a disc and cultipacker to seal the soil surface immediately after fumigant application.

Precautions:

- Do not plow the ground in the spring in such a way that inverts the soil prior to a spring fumigation. Such tillage operations should be conducted in the fall to allow winter kill of residual nematode populations in the top 1 to 2 inches of the soil profile
- Refer to the product label for recommended soil conditions and additional use precautions for optimum fumigant performance.

*Trademark of DowElanco

(Label code) Approved ___/___/___
Initial printing.

Amendments

- 1) Restricts use rates for control of wireworm and northern root knot nematode in potatoes.
- 2) Eliminates plow-sole applications in potatoes.