

PM 21

62719-32

12/7/98

page 18 of 24



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

DEC 7 1998

Bruce A. Houtman  
Regulatory Manager  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, IN 46268

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

SUBJECT: Review of Amended Labeling  
Telone™ C-17 EPA Reg. No. 62719- 12  
Telone™ II EPA Reg. No. 62719- 32  
Telone™ C-35 EPA Reg. No. 62719-302  
Your Submissions Dated October 29, 1998

Dear Mr. Houtman:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, are acceptable subject to the following conditions:

1. Make the following correction for all of the subject product labels: under "Ground water advisory" section, change the word "When" to "Where" in the last sentence (...Montana: **Where** groundwater...).
2. For Telone™ II (EPA Reg. No. 62719- 32): on page 5, modify the sentence under the "Agricultural Use Requirements" box to read "Refer to label booklet for **additional precautionary information and Directions for Use.**" because there are additional precautionary language contained in the booklet.
3. For Telone™ C-35 (EPA Reg. No. 62719-302): on page 6, modify the sentence under the "Agricultural Use Requirements" box to read "Refer to label booklet for **additional precautionary information and Directions for Use including Storage and Disposal.**" because there are additional precautionary language contained in the booklet.

Stamped copies of the labeling are enclosed for your records. Please submit one (1) copy each of your final printed labeling before you release the products for shipment. If you have any questions, then please contact Terri Stowe of my staff at (703) 305-6117.

Sincerely,

*Mary L. Waller*

Mary L. Waller  
Product Manager (21)  
Fungicide Branch  
Registration Division (7505C)

Enclosures

2724

(Base label):

**RESTRICTED USE PESTICIDE**

Due to high acute inhalation toxicity and carcinogenicity.  
For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

(logo) Dow AgroSciences LLC

**Telone\* II**

**A liquid fumigant for preplant treatment of soil to control plant parasitic nematodes and certain other soil pests in cropland**

**Not for use in greenhouses or other enclosed areas.**

Active Ingredient:	(by weight)
1,3-dichloropropene.....	94%
Inert Ingredients.....	6%
Total .....	100%

1 gallon of Telone weighs 10.1 lb at 70°F.

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated:  
DEC 7 1998**

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.**

62719-32

**Keep Out of Reach of Children**

**WARNING AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**Precautionary Statements**

**Hazards to Humans and Domestic Animals**

**Keep Out of Reach of Children**

**WARNING AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**Hazardous Liquid And Vapor**

- Do not swallow any of this product. May be fatal if swallowed.
- Do not get in eyes. Causes substantial, but temporary eye injury.
- Do not get on skin. May be fatal if absorbed through the skin. Causes skin irritation and, if confined, skin burns. May cause allergic skin reaction.
- Do not breathe vapor. May be fatal if inhaled. May cause lung, liver, and kidney damage and respiratory system irritation upon prolonged contact.
- The use of this product may be hazardous to your health. This product contains 1,3-dichloropropene, which has been determined to cause tumors in laboratory animals. Risks can be reduced by exactly following directions for use, precautionary statements, by wearing the personal protective equipment specified in this labeling.

3 8 24

## Personal Protective Equipment (PPE)

**Chemical-Resistant Materials:** Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category H on an EPA chemical resistance category selection chart. PPE constructed of Saranex, neoprene, and chlorinated polyethylene provide short-term contact or splash protection against liquid in this product. Longer-term protection is provided by PPE constructed of Viton, Teflon, and EVAL barrier laminates (for example Responder suits manufactured by Life-guard or Silvershield gloves manufactured by North). Where chemical-resistant materials are required, leather, canvas, or cotton materials offer no protection from this product and must not be worn when contact with this product is possible. Coveralls must be loose-fitting and constructed of woven fabrics (e.g. tight knot cotton or cotton/polyester), non-woven fabrics (e.g. Tyvek or Sontara), or fabrics containing microporous Teflon.

### 1. Handlers Performing Direct Contact Tasks

Direct contact tasks are tasks performed outdoors or in a well-ventilated area. They include:

- equipment calibration or adjustment
- equipment clean-up and repair
- product sampling
- any activity less than 6 feet from an unshielded pressurized hose containing this product
- removal of tarp or plastic film
- rinsate disposal
- fumigant transfer
- clean-up of small spills
- preparing containers for aeration
- any other task not otherwise listed in (2), (3), (4) or (5) below

Handlers performing direct-contact tasks must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves, such as barrier laminate (EVAL) or viton.
- Chemical-resistant footwear plus socks
- Face-sealing goggles, unless full-face respirator is worn
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron
- Respirator with either an organic-vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G). See further respirator requirements in the "User Safety Requirements" section on this label.

### 2. Handlers in Enclosed Cabs

Applicators and other handlers in enclosed cabs must wear:

- Coveralls
- Shoes and socks
- A half-face respirator with either an organic-vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G). A respirator is not required if the occupants are within an enclosed cab that is in conformance with one of the following: 1) ASAE Standard S525 sections 7.1.5, 7.1.7, 7.2.3, and 9, or 2) the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides -- 40 CFR 170.240(d)(5). The cab must be equipped with a vapor-adsorptive filter containing a minimum of 1000 grams activated charcoal. The filter must be changed after no more than 50 hours of application time. See further respirator requirements in the "User Safety Requirements" section on this label.
- In addition, the PPE specified in (1) for direct-contact activities must be immediately available in the enclosed cab and must be worn if the handler leaves the enclosed cab to perform any direct contact activity

4224

### 3. Applicators Outside an Enclosed Cab

Applicators applying Telone\* II soil fumigant (or sealing the soil following application of Telone II) who are not inside an enclosed cab that meets requirements specified above must wear:

- Coveralls over short-sleeved shirt and short pants,
- Chemical-resistant gloves, such as barrier laminate (EVAL) or viton,
- Chemical-resistant footwear plus socks,
- Face-sealing goggles, unless full-face respirator is worn
- Chemical-resistant headgear for overhead exposure,
- A respirator with either an organic-vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G). Face-sealing goggles must be worn unless full-face respirator is worn. See further respirator requirements in the "User Safety Requirements" section on this label.

### 4. Handlers in Treated Area within 5 days After Application

Only the following handler tasks may be performed in the treated area within 5 days after the application is complete:

- Assessing/adjusting the soil seal
- Assessing pest control, application technique, or application efficacy
- Sampling air or soil for this product

**All other tasks are prohibited until the 5 day period has expired.**

Unless in an enclosed cab as described in (2) above, handlers performing the above tasks in the treated area within 5 days after application must wear:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate (EVAL) or viton
- Chemical-resistant footwear and socks
- A respirator with either an organic-vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G). See further respirator requirements in the "User Safety Requirements" section on this label.

### 5. Handlers Exposed to High Concentrations

Handlers exposed to high airborne concentrations of this product, such as cleanup following large spills and exposure to this product in poorly ventilated areas must wear:

- Chemical-resistant suit
- Chemical-resistant gloves, such as barrier laminate (EVAL) or viton
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear
- Supplied-air respirator with MSHA/NIOSH approval number prefix TC-19C or self-contained breathing apparatus (SCBA) with MSHA/NIOSH approval number prefix TC-13F. See further respirator requirements in the "User Safety Requirements" section on this label.

**Note:** In-tank cleaning of bulk tanks must be performed only by persons who have been specifically trained for this activity. Refer to OSHA 29 CFR Part 1910.146 and Telone User's Guide section on Storage Tanks.

### Engineering Controls Requirements

**Mechanical Transfer System:** Personal protective equipment specified for "Direct Contact Activities" must be worn by the operator of the mechanical transfer system. The operator of the mechanical transfer system must follow instructions on proper operation of the system found in the "Telone Soil Fumigants - A Guide to Application" manual. Contact your distributor for Telone II for more information or these materials.

**End-Row Spillage Control:** The dispensing system must shut off the feed stream when chisels are raised out of the ground. Do not stop or park near any area where dribble from chisel tips has fallen. The applicator must follow instructions on proper operation and maintenance of the system found in the "Telone Soil Fumigants - A Guide to Application" manual. Contact your distributor for Telone II for more information or these materials.

- A flow shutoff device must be placed as close as is technically feasible to the fluid discharge point. This can be a ball, poppet, or diaphragm check valve, or full flow shutoff device such as an electric or pneumatically actuated valve.
- Check valves must be replaced immediately if continuous drip occurs.
- Place check valves above the orifice.
- Isolate the check valve from upstream pressure by installing a main line shut off or bypass valve prior to the manifold.
- Do not exceed 1/4 inch diameter tubing.
- Do not use any method of end-row spillage control other than that stated on this label.
- An alternative to shutoff devices is a purge system which clears the line of all liquid. Consult your Dow AgroSciences representative for purge system description.

**With all bulk and mini-bulk containers, Telone II must be transferred through connecting hoses, pipes, and/or couplings sufficiently tight to prevent workers or other persons from coming in contact with liquid Telone II.**

1. All hoses, piping, and tanks used in connection with Telone II shall be of type appropriate for use under the pressure and vacuum conditions to be encountered.
2. External sight gauges shall be equipped with valves so that pipes to sight gauge can be shut off in case of breakage or leakage.
3. The mechanical transfer system must be adequate to make necessary measurements of the pesticide being used.
4. Shut-off devices must be installed on the exit end of all hoses and at all disconnect points to prevent leakage of Telone product when the transfer is stopped and hose is removed or disconnected. A dry coupler that will minimize pesticide leakage must be installed at the disconnect point.
5. The pressure in hoses used to move Telone II beyond a pump must not exceed the manufacturer's maximum pressure specification.

### User Safety Requirements

1. **Respirator Requirements:** When a respirator is required for use with this product, the following criteria must be met:
  - a. Cartridges or canisters must be replaced daily or when odor or irritation from this product becomes apparent, whichever is sooner.
  - b. Respirators must be fit-tested and fit-checked using a program that conforms to OSHA's requirements (described in 29 CFR Part 1910.134).
  - c. Respirator users must be trained using a program that conforms to OSHA's requirements (described in 29 CFR Part 1910.134).
  - d. Respirator users must be examined by a qualified medical practitioner to ensure physical ability to safely wear the style of respirator to be worn.
2. **Dispose of Contaminated Clothing:** Discard clothing and other absorbent materials that have been drenched or heavily contaminated with liquid from this product. Do not reuse them.
3. **Clean and Maintain PPE:** Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Wash PPE after each day's use.
4. **Contact With Mouth:** Never siphon this product by mouth or use mouth to blow out clogged lines, nozzles, etc.

6724

**5. Heat Illness Avoidance:** Use measures to avoid or minimize heat illness while using this product. These measures include gradual adjustment to heat and respirator stress, fans for cooling, cooling vests, frequent breaks to cool down, frequent intake of drinking water, and maintaining weight from day to day.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

**If inhaled:** Remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. If breathing is difficult, give oxygen. Call a physician.

**If on skin:** Immediately flush skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Call a physician. If water is not immediately available, remove excess chemical from skin with adsorbent material such as towel or dry soil, then proceed at once to a location where water is available and thoroughly wash contaminated skin with plenty of water. Call a physician.

**If in eyes:** Immediately flush eyes with plenty of water for at least 15 minutes. Call a physician.

**If swallowed:** Do not induce vomiting. Call a physician or Poison Control Center immediately. Never give anything by mouth to an unconscious person.

**Note to physician:** Because rapid absorption may occur through lungs if product is aspirated and cause systemic effects, the decision to induce vomiting or not should be made by a physician. If lavage is performed, endotracheal and/or esophageal control is suggested. Danger from lung aspiration must be weighed against toxicity when considering emptying the stomach.

### Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposal of equipment washwaters. See "Storage, Shipment and Disposal" section. In case of spills properly dispose of contaminated materials.

**Ground water advisory:** 1,3-dichloropropene is known to move through soil and under certain conditions has the potential to reach ground water as a result of agricultural use. Application in areas where soils are permeable and ground water is near the surface, or in karst geology, could result in ground water contamination. Do not apply within 100 feet of any well used for potable water. Do not apply in areas overlying karst geology. In North Dakota, South Dakota, Wisconsin, Minnesota, New York, Maine, New Hampshire, Vermont, Massachusetts, Utah, and Montana: When groundwater aquifers exist at a depth of 50 feet or less from the surface, do not apply this product where soils are Hydrologic Group A.

### Physical or Chemical Hazards

**Flammable - Do not use, pour, spill, or store near heat or open flame. Do not cut or weld container.**

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to label booklet for additional precautionary information including Personal Protective Equipment (PPE), User Safety Recommendations and Directions for Use, including Storage and Disposal.



5724

(Label Booklet cover):

**RESTRICTED USE PESTICIDE**  
 Due to high acute inhalation toxicity and carcinogenicity.  
 For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

(logo) Dow AgroSciences LLC

# Telone\* II

**A liquid fumigant for preplant treatment of soil to control plant parasitic nematodes and certain other soil pests in cropland**

**Not for use in greenhouses or other enclosed areas.**

Active Ingredient:	(by weight)
1,3-dichloropropene.....	94%
Inert Ingredients.....	6%
Total .....	100%

1 gallon of Telone weighs 10.1 lb at 70°F.

**Keep Out of Reach of Children**

## WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**Agricultural Use Requirements**  
 Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

**Refer to label booklet for additional precautionary information including Personal Protective Equipment (PPE), User Safety Recommendations and Directions for Use including Storage and Disposal.**

**Notice:** Read the entire label. Use only according to label directions. **Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.**

*In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at [www.dowagro.com](http://www.dowagro.com).*

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-32

28      17      45      44      69  
 EPA Est. 464-TX-1; 464-MI-1; 464-CA-1; 464-CA-2; 54686-WA-1  
 Superscripts correspond to places 7 & 8 of lot number.

\*Trademark of Dow AgroSciences LLC



9 7 24

Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

# Soil Fumigant

Net Contents \_\_\_

10 7 94

(Page 1 through end):

<b>Table of Contents</b>	<b>Page</b>
Precautionary Statements	
Hazards to Humans and Domestic Animals	
Personal Protective Equipment	
Engineering Controls Requirements	
User Safety Recommendations	
First Aid	
Environmental Hazards	
Physical or Chemical Hazards	
Directions for Use	
Agricultural Use Requirements	
Storage, Shipment and Disposal	
General Information	
General Use Precautions	
Application Directions	
Application Timing	
Soil Conditions	
Soil Moisture	
Soil Preparation	
Placement of Fumigant	
Application Methods and Equipment	
Sealing the Soil After Application	
Soil Fumigation Interval	
Approved Uses	
Control of Nematodes: Vegetable Crops, Field Crops, Fruit and Nut Crops, Nursery Crops	
Broadcast Treatment Rates for Nematode Control (Table 1)	
Rate Conversion Chart for Various Row Spacings and Fumigant Flow Rates (Table 2)	
Plant Diseases	
Bacterial Canker of Peaches	
<i>Fusarium</i> Wilt of Cotton	
Sugar Beet <i>Rhizomania</i> Disease	
<i>Verticillium</i> Wilt of Mint	
Soil Insects	
Symphylans (Garden Centipedes)	
Wireworms	
Warranty Disclaimer	
Inherent Risks of Use	
Limitation of Remedies	

---

## Precautionary Statements

---

### Hazards to Humans and Domestic Animals

## WARNING — AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

#### Hazardous Liquid And Vapor

- Do not swallow any of this product. May be fatal if swallowed.
- Do not get in eyes. Causes substantial, but temporary eye injury.
- Do not get on skin. May be fatal if absorbed through the skin. Causes skin irritation and, if confined, skin burns. May cause allergic skin reaction.
- Do not breathe vapor. May be fatal if inhaled. May cause lung, liver, and kidney damage and respiratory system irritation upon prolonged contact.
- The use of this product may be hazardous to your health. This product contains 1,3-dichloropropene, which has been determined to cause tumors in laboratory animals. Risks can be reduced by exactly following directions for use, precautionary statements, by wearing the personal protective equipment specified in this labeling.

#### Personal Protective Equipment (PPE)

**Chemical-Resistant Materials:** Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category H on an EPA chemical resistance category selection chart. PPE constructed of Saranex, neoprene, and chlorinated polyethylene provide short-term contact or splash protection against liquid in this product. Longer-term protection is provided by PPE constructed of Viton, Teflon, and EVAL barrier laminates (for example Responder suits manufactured by Life-guard or Silvershield gloves manufactured by North). Where chemical-resistant materials are required, leather, canvas, or cotton materials offer no protection from this product and must not be worn when contact with this product is possible. Coveralls must be loose-fitting and constructed of woven fabrics (e.g. tight knot cotton or cotton/polyester), non-woven fabrics (e.g. Tyvek or Sontara), or fabrics containing microporous Teflon.

#### 1. Handlers Performing Direct Contact Tasks

Direct contact tasks are tasks performed outdoors or in a well-ventilated area. They include:

- |  |   |
|--|---|
| <ul style="list-style-type: none"> <li>• equipment calibration or adjustment</li> <li>• equipment clean-up and repair</li> <li>• product sampling</li> <li>• any activity less than 6 feet from an unshielded pressurized hose containing this product</li> <li>• removal of tarp or plastic film</li> </ul> | <ul style="list-style-type: none"> <li>• rinsate disposal</li> <li>• fumigant transfer</li> <li>• clean-up of small spills</li> <li>• preparing containers for aeration</li> <li>• any other task not otherwise listed in (2), (3), (4) or (5) below</li> </ul> |
|--|---|

Handlers performing direct-contact tasks must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves, such as barrier laminate (EVAL) or viton
- Chemical-resistant footwear plus socks
- Face-sealing goggles, unless full-face respirator is worn
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron

12 7 24

- Respirator with either an organic-vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G). See further respirator requirements in the "User Safety Requirements" section on this label.

## 2. Handlers in Enclosed Cabs

Applicators and other handlers in enclosed cabs must wear:

- Coveralls
- Shoes and socks
- A half-face respirator with either an organic-vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G). A respirator is not required if the occupants are within an enclosed cab that is in conformance with one of the following: 1) ASAE Standard S525 sections 7.1.5, 7.1.7, 7.2.3, and 9, or 2) the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides -- 40 CFR 170.240(d)(5). The cab must be equipped with a vapor-adsorptive filter containing a minimum of 1000 grams activated charcoal. The filter must be changed after no more than 50 hours of application time. See further respirator requirements in the "User Safety Requirements" section on this label.
- In addition, the PPE specified in (1) for direct-contact activities must be immediately available in the enclosed cab and must be worn if the handler leaves the enclosed cab to perform any direct-contact activity

## 3. Applicators Outside an Enclosed Cab

Applicators applying Telone\* II soil fumigant (or sealing the soil following application of Telone II) who are not inside an enclosed cab that meets requirements specified above must wear:

- Coveralls over short-sleeved shirt and short pants,
- Chemical-resistant gloves, such as barrier laminate (EVAL) or viton,
- Chemical-resistant footwear plus socks,
- Face-sealing goggles, unless full-face respirator is worn
- Chemical-resistant headgear for overhead exposure,
- A respirator with either an organic-vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G). Face-sealing goggles must be worn unless full-face respirator is worn. See further respirator requirements in the "User Safety Requirements" section on this label.

## 4. Handlers in Treated Area within 5 days After Application

Only the following handler tasks may be performed in the treated area within 5 days after the application is complete:

- Assessing/adjusting the soil seal
- Assessing pest control, application technique, or application efficacy
- Sampling air or soil for this product

**All other tasks are prohibited until the 5 day period has expired.**

Unless in an enclosed cab as described in (2) above, handlers performing the above tasks in the treated area within 5 days after application must wear:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate (EVAL) or viton
- Chemical-resistant footwear and socks
- A respirator with either an organic-vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G). See further respirator requirements in the "User Safety Requirements" section on this label.

### 5. Handlers Exposed to High Concentrations

Handlers exposed to high airborne concentrations of this product, such as cleanup following large spills and exposure to this product in poorly ventilated areas must wear:

- Chemical-resistant suit
- Chemical-resistant gloves, such as barrier laminate (EVAL) or viton
- Chemical -resistant footwear plus socks
- Chemical-resistant headgear
- Supplied-air respirator with MSHA/NIOSH approval number prefix TC-19C or self-contained breathing apparatus (SCBA) with MSHA/NIOSH approval number prefix TC-13F. See further respirator requirements in the "User Safety Requirements" section on this label.

**Note:** In-tank cleaning of bulk tanks must be performed only by persons who have been specifically trained for this activity. Refer to OSHA 29 CFR Part 1910.146 and Telone User's Guide section on Storage Tanks.

### Engineering Controls Requirements

**Mechanical Transfer System:** Personal protective equipment specified for "Direct Contact Activities" must be worn by the operator of the mechanical transfer system. The operator of the mechanical transfer system must follow instructions on proper operation of the system found in the "Telone Soil Fumigants - A Guide to Application" manual. Contact your distributor for Telone II for more information or these materials.

**End-Row Spillage Control:** The dispensing system must shut off the feed stream when chisels are raised out of the ground. Do not stop or park near any area where dribble from chisel tips has fallen. The applicator must follow instructions on proper operation and maintenance of the system found in the "Telone Soil Fumigants - A Guide to Application" manual. Contact your distributor for Telone II for more information or these materials.

- A flow shutoff device must be placed as close as is technically feasible to the fluid discharge point. This can be a ball, poppet, or diaphragm check valve, or full flow shutoff device such as an electric or pneumatically actuated valve.
- Check valves must be replaced immediately if continuous drip occurs.
- Place check valves above the orifice.
- Isolate the check valve from upstream pressure by installing a main line shut off or bypass valve prior to the manifold.
- Do not exceed 1/4 inch diameter tubing.
- Do not use any method of end-row spillage control other than that stated on this label.
- An alternative to shutoff devices is a purge system which clears the line of all liquid. Consult your Dow AgroSciences representative for purge system description.

**With all bulk and mini-bulk containers, Telone II must be transferred through connecting hoses, pipes, and/or couplings sufficiently tight to prevent workers or other persons from coming in contact with liquid Telone II.**

1. All hoses, piping, and tanks used in connection with Telone II shall be of type appropriate for use under the pressure and vacuum conditions to be encountered.
2. External sight gauges shall be equipped with valves so that pipes to sight gauge can be shut off in case of breakage or leakage.
3. The mechanical transfer system must be adequate to make necessary measurements of the pesticide being used.
4. Shut-off devices must be installed on the exit end of all hoses and at all disconnect points to prevent leakage of Telone product when the transfer is stopped and hose is removed or disconnected. A dry coupler that will **minimize pesticide leakage** must be installed at the disconnect point.

5. The pressure in hoses used to move Telone II beyond a pump **must not exceed** the manufacturer's maximum pressure specification.

### User Safety Requirements

1. **Respirator Requirements:** When a respirator is required for use with this product, the following criteria must be met:
  - a. Cartridges or canisters must be replaced daily or when odor or irritation from this product becomes apparent, whichever is sooner.
  - b. Respirators must be fit-tested and fit-checked using a program that conforms to OSHA's requirements (described in 29 CFR Part 1910.134).
  - c. Respirator users must be trained using a program that conforms to OSHA's requirements (described in 29 CFR Part 1910.134).
  - d. Respirator users must be examined by a qualified medical practitioner to ensure physical ability to safely wear the style of respirator to be worn.
2. **Dispose of Contaminated Clothing:** Discard clothing and other absorbent materials that have been drenched or heavily contaminated with liquid from this product. Do not reuse them.
3. **Clean and Maintain PPE:** Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Wash PPE after each day's use.
4. **Contact With Mouth:** Never siphon this product by mouth or use mouth to blow out clogged lines, nozzles, etc.
5. **Heat Illness Avoidance:** Use measures to avoid or minimize heat illness while using this product. These measures include gradual adjustment to heat and respirator stress, fans for cooling, cooling vests, frequent breaks to cool down, frequent intake of drinking water, and maintaining weight from day to day.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

**If inhaled:** Remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. If breathing is difficult, give oxygen. Call a physician.

**If on skin:** Immediately flush skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Call a physician. If water is not immediately available, remove excess chemical from skin with adsorbent material such as towel or dry soil, then proceed at once to a location where water is available and thoroughly wash contaminated skin with plenty of water. Call a physician.

**If in eyes:** Immediately flush eyes with plenty of water for at least 15 minutes. Call a physician.

**If swallowed:** Do not induce vomiting. Call a physician or Poison Control Center immediately. Never give anything by mouth to an unconscious person.

**Note to physician:** Because rapid absorption may occur through lungs if product is aspirated and cause systemic effects, the decision to induce vomiting or not should be made by a physician. If lavage is performed, endotracheal and/or esophageal control is suggested. Danger from lung aspiration must be weighed against toxicity when considering emptying the stomach.

15 7 24

### Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposal of equipment washwaters. See "Storage, Shipment and Disposal" section. In case of spills properly dispose of contaminated materials.

**Ground water advisory:** 1,3-dichloropropene is known to move through soil and under certain conditions has the potential to reach ground water as a result of agricultural use. Application in areas where soils are permeable and ground water is near the surface, or in karst geology, could result in ground water contamination. Do not apply within 100 feet of any well used for potable water. Do not apply in areas overlying karst geology. In North Dakota, South Dakota, Wisconsin, Minnesota, New York, Maine, New Hampshire, Vermont, Massachusetts, Utah, and Montana: When groundwater aquifers exist at a depth of 50 feet or less from the surface, do not apply this product where soils are Hydrologic Group A.

### Physical or Chemical Hazards

**Flammable - Do not use, pour, spill, or store near heat or open flame. Do not cut or weld container.**

---

### Directions for Use

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals, and notification to workers. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

#### Entry Restriction:

Entry (including early entry that would otherwise be permitted under the WPS) by any person -- other than a correctly trained and equipped handler who is performing a handling task permitted on this labeling -- is **prohibited** from the start of application until 5 days after application. **In addition**, if tarps are used for the application, non-handler entry is prohibited while tarps are being removed.

#### Notification:

Notify workers of the application by warning them orally and by posting fumigant warning signs at entrances to treated areas. The sign must bear the skull and crossbones symbol and state: (1) "DANGER/PELIGRO," (2) Areas under fumigation, **DO NOT ENTER/NO ENTRE**," (3) the date and time of fumigation, (4) "Telone II Fumigant in use," and (5) name, address, and telephone number of the applicator." Post the fumigant warning sign instead of the WPS sign for this application, but follow all WPS requirements pertaining to location, legibility, size and timing of posting and removal.

#### PPE For Reentry During The Entry-Restricted Period:

PPE for entry that is permitted by this labeling is listed in the "Hazards to Humans and Domestic Animals" section of this labeling.

16724

### Storage, Shipment and Disposal

**Shipment:** Do not ship or store with food, feeds, drugs, or clothing.

**Storage:** Store in tightly-closed original container in a cool place away from dwellings. Do not allow contamination of seeds, plants, fertilizers, or other pesticide chemicals. Do not contaminate food, feedstuffs, drugs, or domestic water supplies.

**Disposal:** Pesticide wastes are toxic. Improper disposal of excess pesticide and rinsates is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA regional office for guidance.

Because Telone II is corrosive under certain conditions, flush all application equipment with fuel oil, kerosene or a similar type of petroleum solvent immediately after use. Fill pumps and meters with new motor oil or a 50% motor oil/fuel oil mixture before storing. **Do not use water.** Dispose of rinsate by applicable Federal, state and local regulations. Never introduce rinsate or unused Telone II into surface or underground water supplies.

**Metal Container Disposal:** To dispose of container emptied during application operation, remove bungs, invert container in the field just treated and ensure that the container is free of liquid. Orient container such that ventilation of bung holes is not restricted. Allow containers to aerate for at least 14 days.

Replace bungs prior to transport. After aeration, offer container to qualified reconditioner or dispose of as directed by State or local regulations.

**Refillable Containers:** Follow cleaning and handling directions in the Telone User's Guide.

### General Information

Telone II is a liquid fumigant for preplant treatment of soil to control plant parasitic nematodes and certain other soil pests and plant diseases in cropland.

Telone II may be applied as a preplant soil treatment to control the following types of plant parasitic nematodes: burrowing, citrus, cyst (golden, sugar beet, soybean, carrot and wheat), dagger, lance, pin, needle, reniform, ring, root knot, root lesion, spiral, sting and stubby root.

Telone II can also be used to control garden centipedes (symphylans) and wireworms, suppress sugar beet Rhizomania disease, *Fusarium* wilt of cotton and *Verticillium* wilt of mint and aid in the control of bacterial canker of peaches.

Before fumigation, soil sampling for the type and number of pests present is recommended. In fields where pre-treatment soil samples indicate the presence of high population levels of nematodes, a successful fumigation cannot be expected to eradicate entire populations. Therefore, post-treatment sampling is recommended to determine the need for additional pest management practices.

Consult State Agricultural Experiment Station or Extension Service specialists for information on other practices such as post-harvest destruction of crop residues, weed control or other cultural practices, and use of nematode resistant crop varieties that may aid in reducing crop losses from soil borne pests.

### General Use Precautions

Soil fumigation using Telone II should be conducted only according to directions and conditions of use described in this labeling.

**Formulator use of Telone II:** Labeling for end use products containing Telone II that are prepared and sold by formulators must comply with all precautionary statements, use precautions, environmental hazards, handling and protective equipment requirements, maximum application rates and other exposure mitigation measures specified in this product labeling. Any product formulated from this product and/or any product which is formulated from the repackaging of this product must be labeled only as a pre-plant



17724

soil injected and/or a soil fumigant product. Each formulator is responsible for obtaining EPA registration for each end use product. Note: A revised groundwater advisory in accordance with EPA agreement dated Sept. 30, 1998, must be included on the label of any product reformulated from Telone II after August 1, 1999.

**Recontamination prevention:** Telone II will control pests that are present in the soil treatment zone at time of fumigation. It will not control pests that are introduced into soil after fumigation. To avoid reinfestation of treated soil do not use irrigation water, transplants, seed pieces, or equipment that could carry soil borne pests from infested land. Avoid contamination from moving infested soil onto treated beds through cultivation, movement of soil from below the treated zone, dumping contaminated tare soil in treated fields and soil contamination from equipment or crop remains. Clean equipment carefully before entering treated fields.

**Do not use containers, pumps or other transfer equipment made of aluminum, magnesium or their alloys, as under certain conditions Telone II may be severely corrosive to such metals.**

**Equipment Clean-up:** Because Telone II is corrosive under certain conditions, flush all application equipment with fuel oil, kerosene or a similar type of petroleum solvent immediately after use. Fill pumps and meters with new motor oil or a 50% motor oil/fuel oil mixture before storing. **Do not use water.** Dispose of rinsate by incorporation into field just treated or by other approved means. Never introduce rinsate or unused Telone II into surface or underground water supplies.

**Chemigation:** Do not apply Telone II through any type of irrigation system.

**Fertility Interactions:** Fumigation may temporarily raise the level of ammonia nitrogen and soluble salts in the soil. This is most likely to occur when heavy rates of fertilizer and fumigant are applied to soils that are either cold, wet, acid, or high in organic matter. To avoid injury to certain crops including red beets, carrots, corn, radishes, cole crops, legumes (beans), lettuce, onions, and sugar beets, fertilize as indicated by soil tests made after fumigation. To avoid ammonia injury or nitrate starvation (or both) to crops grown on high organic soils, do not use fertilizers containing ammonium salts. Use only fertilizers containing nitrates until after the crop is well established and the soil temperature is above 65°F.

When using high rates of Telone II as required by certain state nursery regulations, liming of highly acid soils before fumigation may stimulate nitrification and reduce the possibility of ammonia toxicity. Certain nursery crops such as citrus seedlings, *Comus* sp., *Crataegus* sp., spruce, and vegetable crops such as cauliflower have shown evidence of phosphorus deficiency following fumigation. To avoid this possible effect, additional phosphate fertilizer (foliar applied) is recommended where experience indicates a deficiency may occur.

## Application Directions

### Application Timing

Telone II can be applied at any time of the year when soil conditions permit. Conditions that allow rapid diffusion of the fumigant as a gas through the soil normally give best results. Because Telone II does not provide residual control of soil pests, it should be used as a preplant application before planting each crop. The following soil temperature and moisture conditions should exist at time of treatment. Failure to meet these conditions may result in unsatisfactory product performance:

### Soil Conditions

**Soil temperature** at the depth of application must be between 40°F and 80°F. In areas where the soil temperature in the spring may not reach 40°F in time to allow application of Telone II prior to planting, late summer or early fall treatment is recommended.

18724

### Soil moisture

It is critical to manage soil moisture properly before fumigation. Plan fumigation for seasons, crop rotations, or irrigation schedules which leave moisture in the soil. For application depths greater than 18 inches, the soil should be moist within a 16 inch radius upwards from the point of injection as determined by the feel method (see below). ~~For all other applications, the soil must be moist from two inches below the soil surface to at least 12 inches deep as determined by the feel method (see below).~~ The amount of moisture needed in this zone will vary according to soil type. The surface soil generally dries very rapidly and should not be considered in this determination. If there is insufficient moisture at the two to six inch depth, the soil moisture must be adjusted. If irrigation is not available and there is adequate soil moisture below six inches, it may be brought to the surface by disking or plowing before or during the injection. To conserve existing soil moisture, pretreatment or treatment tillage practices should be done as close to the time of application as possible. For fields with more than one soil texture, soil moisture content in the lightest textured (most sandy) areas must comply with this soil moisture requirement. Whenever possible, the field should be divided into areas of similar soil texture and the soil moisture of each area should be adjusted as needed. Coarser textured soils can be fumigated under conditions of higher soil moisture than finer textured soils; however, if the soil moisture is too high, fumigant movement will be retarded and effectiveness of the treatment will be reduced. Previous and/or local experience with the soil to be treated or the crop to be planted can often serve as a guide to conditions that will be acceptable. If you do not know how to determine the soil moisture content of the area to be treated, consult your local extension service or soil conservation service specialist or pest control advisor (ag consultant) for assistance.

In general, no irrigation should immediately precede subsoiling or fumigation; however, when irrigation is available and surface soil moisture conditions are not likely to provide an adequate seal against fumigant loss, a very light sprinkler irrigation to wet the top 1 to 2 inches of soil may be used to bring soil moisture content to the desired level.

**The following descriptions will aid in determining acceptable soil moisture conditions by the "feel method".** For coarse soils (sand and loamy sand), there must be enough moisture to allow formation of a weak ball when compressed in the hand. Due to soil texture, this ball is easily broken with little disturbance. In loamy, moderately coarse, or medium textured soils (coarse sandy loam, sandy loam, and fine sandy loam), a soil sample with the proper moisture content can be formed into a ball which holds together with moderate disturbance, but does not stick between the thumb and forefinger. Fine textured soils (clay loam, silty clay loam, sandy clay, silty clay, sandy clay loam and clay), should be pliable and not crumbly, but should not form a ribbon when compressed between the thumb and forefinger.

### Soil Preparation

The soil should be free of clods. Large clods can prevent effective soil sealing and reduce effectiveness of Telone II. Plant residues should be thoroughly incorporated into the soil prior to treatment to avoid interfering with application. Undecomposed plant material may harbor pests that will not be controlled by fumigation. Little or no crop residue should be present on the soil surface. Crop residue that is present should lie flat to permit the soil to be sealed effectively. Compacted soil layers within the desired treatment zone should be fractured before or during application of the fumigant. Deviation from the above conditions may result in unsatisfactory results.

### Placement of Fumigant

Telone II may be applied as either a broadcast (overall) or row treatment. It must be placed at least 12 inches below the final soil surface. When soil conditions allow, placement a minimum of 14 inches below the final soil surface is recommended. Deeper placement is recommended when fumigating soil to be planted to deep-rooted plants, such as perennial fruit and nut crops, or to control deeply distributed pests. For row application, the fumigant must be placed at least 12 inches from the nearest soil/air interface (e.g. furrow).

### Application Methods and Equipment

**Broadcast Application:** Use chisel (shank), offset wing shank, Nobel (sweep) plow or plow-sole application equipment. For best results when using chisel equipment, use ripper-type, forward-swept

19 8 24

shanks. Nobel plow equipment is particularly useful for fall fumigation when the soil still contains some undecomposed standing plant material. Subsoiling may be necessary before application as described under "Soil Preparation". Choose application equipment which allows the deepest application and best soil seal under existing conditions.

The fumigant outlet spacing varies with the type of application equipment used.

**With chisel equipment** a fumigant shank spacing of 12 to 24 inches is recommended. The outlet spacing for this equipment may be up to 1 1/2 times the application depth but generally should be equal to the application depth and should not exceed the soil-shattering capability of the chisels. The maximum outlet spacing should not exceed 24 inches.

**With plow-sole equipment** a 12-inch outlet spacing is recommended. Do not exceed an outlet spacing of 18 inches.

**With Nobel (sweep) plow equipment** use an outlet spacing of 9-12 inches along the sweeps.

Broadcast application can be made in the same direction or at an angle to the direction of row planting. Refer to Table 1 for broadcast treatment rates for various crops.

**Row Application (for row spacing greater than 24 inches):** Use chisel equipment to treat a band of soil where the crop is to be planted, i.e. the plant row. In general, when one chisel is used, apply Telone II at twice the flow rates given in Table 1. When multiple chisels per plant row are used, space the chisels (fumigant outlets) 8 to 12 inches apart and use the flow rates given in Table 1 per outlet (see footnote 1, Table 2). Regardless of the number or spacing of chisels used, the fumigant must be placed at least 12 inches from the nearest soil/air interface (e.g. furrow). With certain deeper rooted crops such as potatoes and sugar beets, higher flow rates may be necessary to ensure adequate treatment of the zone of soil where primary root growth occurs; however, **in no case should the amount of fumigant applied per acre exceed the maximum gallons per acre rates given in Table 1.** To determine the amount (gallons) of Telone II required per acre for various plant row spacings and flow rates, refer to Table 2. Note that as the distance between the plant rows increases the amount of fumigant required decreases and vice versa.

To prevent seed germination problems caused by improper seed-to-soil contact or improper seeding depth, do not place the seed directly over the furrow left by the applicator chisel(s). When one chisel is used per plant row, place the seed about 4 inches to one side of the chisel furrow. When two chisels are used per plant row, plant the seed offset from the chisel trace.

#### **Sealing the Soil After Application**

**Immediately** after chisel application of Telone II the soil must be "sealed" to prevent fumigant loss and ensure that an effective concentration of fumigant is maintained within the soil for a period of several days. **For broadcast treatment (flat fumigation),** sealing can be accomplished with equipment that will uniformly mix the soil to a depth of 3 to 4 inches to effectively eliminate chisel or plow traces which can allow direct escape of the fumigant. A tandem disc or similar equipment may be used for this purpose. To maximize sealing, steps should also be taken to compact the soil surface to further retard the rate of fumigant loss by following with a ring roller, cultipacker or roller in combination with tillage equipment. Compaction of the soil surface alone does not effectively disrupt chisel or plow traces.

**For row treatment,** forming the beds at the time of application should be accomplished in a manner that places the fumigant at least 12 inches from the nearest soil/air interface (e.g. furrow). The closest soil/air interface could be the furrow for multiple knife applications or the top of the bed for single knife applications. Row treatments into preformed beds must be sealed by disrupting the chisel trace using press sealers, ring rollers or by reforming the beds and following with such equipment.

20724

Sealing can also be improved by applying non-perforated plastic film, such as polyethylene, over the entire area or in strips. Use of a film to seal the soil surface does not eliminate the need to eliminate chisel traces prior to application of the plastic film.

Proper soil conditions at the time of application (see Soil Preparation section) are important to ensure proper placement of fumigant (see Placement of Fumigant section) and obtaining adequate sealing. Prior tillage should be adequate to eliminate clods and thoroughly mix crop residues into the soil.

### Soil Fumigation Interval

Leave the soil undisturbed and unplanted for at least 7 days after application of the fumigant. A longer undisturbed interval is required if the soil becomes cold or wet, and for deep-rooted tree, shrub and vine planting sites.

After the fumigation interval, to prevent phytotoxicity, allow the fumigant to dissipate completely before planting the crop. Under optimum soil conditions for dissipation, 1 week for each 10 gallons/acre is recommended. To hasten dissipation, especially if heavy rains or low temperatures occur during the treatment period, till the soil to the depth of fumigant application. Use a knife-like chisel without turning the soil to reduce the possibility of recontaminating the treated soil. Dissipation is usually complete when the odor of Telone II is no longer evident at the application depth. Seed may be used as a bioassay to determine if Telone II is present in the soil at concentrations sufficient to cause plant injury. Do not plant if the odor of Telone II is present within the zone of fumigation.

**Buffer Zone:** An application of Telone II shall not be made within 300 feet of an occupied structure, such as a school, hospital, business or residence. No person shall be present at this structure at any time during the seven consecutive day period following application. **This buffer zone does not apply to use on soils to be planted with perennial crops that will not experience an additional 1,3-D treatment for at least three years. For example, on soils to be planted with fruit trees, nut and nursery crops, perennial vines, hops, mint or pineapple, perennial vines, hops, mint, fruit and nut trees.**

## Approved Uses

### Control of Nematodes

Telone II is recommended for control of nematodes in soils to be planted to vegetable crops, field crops, fruit and nut crops, and nursery crops. Refer to Table 1 for broadcast application rates and to Table 2 for row treatment flow rates for specific row spacings.

**Maximum application rates for nematode control (except cyst nematode) in mineral soils for the following crop groups:**

	(Gallons/acre)
Vegetable Crops	12
Field Crops	12
Fruit and Nut Crops	35
Nursery Crops	55

**Table 1. Broadcast Treatment Rates for Nematode Control**

Crop	Soil Type or Texture	Application Rates†	
		Broadcast Gal/Acre	Fl Oz/ 1000 Ft of Row/Outlet <sup>1</sup>
Vegetable crops <sup>2</sup>	Mineral <sup>3</sup>	9 - 12 <sup>4</sup>	26 - 35
	Muck or Peat <sup>5</sup>	25	74
Field crops	Mineral	9 - 12 <sup>4</sup>	26 - 35
	Muck or Peat	18	53

21 3 24

Fruit and Nut Crops <sup>6,7,8,9,10</sup>	Mineral	27 - 35	79 - 103
Nursery Crops <sup>10-11</sup>	Mineral	42 - 55	124 - 162

†Do not exceed specified maximum application rates.

<sup>1</sup>Flow rates are based on 12-inch outlet spacing. Flow rates for alternate spacings can be calculated using the following formula: fl oz/1000 ft of row/outlet = 0.245 x rate in gallons/acre x outlet spacing in inches. For row treatment refer to Table 2.

<sup>2</sup>Potatoes:

Before fumigation, soil sampling for the type and number of pests present is recommended and can help to determine the need for additional treatment with a contact nematicide. Preharvest tuber sampling for nematodes also is recommended. If the nematode population is high enough to damage the crop, potatoes can be harvested early. Do not store potatoes with a detectable nematode infestation.

Row treatment is not recommended for potatoes in irrigated areas of western and northwestern states.

**In Colorado, Idaho, Nevada, Oregon, Utah and Washington, and in Modoc and Siskiyou counties of California** refer to supplemental labeling for Telone II entitled: "For the Control of Nematode Nematodes and the Suppression of Wireworm Wireworms Control in Soils to be Planted to Potatoes or Onions" for directions for use.

<sup>3</sup>Mineral soil includes sand, sandy loam, silt, and clay loam. Use the higher rates for finer textured (heavier) soils.

<sup>4</sup>For cyst-forming nematodes use 18 gallons per acre (53 fl oz/1000 ft/outlet).

<sup>5</sup>Greater than 20% organic matter content.

<sup>6</sup>Pineapple: Application may be made at the time of planting. For best results, seal the soil with polyethylene film, which acts as a gas permeability barrier.

<sup>7</sup>Tree Planting Sites: Use 24 fl oz (1.5 pints) of Telone II by application of the fumigant at a single point in the center of each planting site at a depth of 5 feet below the final soil surface. Sites prepared by backhoeing to break up restrictive soil layers that may retard fumigant movement should be dug in the approximate dimensions of 10 x 10 x 10 feet. The hole should then be backfilled to a depth of 5 feet, the fumigant applied using a closed-system application tube and the remainder of the soil previously removed immediately added to the hole. For sites where no restrictive soil layers are present, the fumigant can be applied to a depth of 5 feet using an injection auger. For best results, prepare and treat planting sites in the fall and plant in the spring.

<sup>8</sup>For shallow-rooted plants grown only one year, use 15 to 27 gallons per acre (44 to 79 fl oz/1000 ft of row/outlet).

<sup>9</sup>Citrus Fruits: For burrowing nematode control, inject Telone II on 18-inch centers at least 12 inches deep. For buffers within existing groves or for tree planting sites within existing groves, do not apply within 5 feet of living trees. Keep the field free of plants susceptible to this nematode at least two years before planting to citrus.

<sup>10</sup>Stone Fruits: Within existing groves or for tree planting sites within existing groves, do not apply within 5 feet of living trees.

22 9 24

~~40-11~~ When used according to state nursery regulations, Telone II may be used in the production of certified nursery stock.

**Table 2. Rate Conversion Chart for Various Row Spacings and Fumigant Flow Rates<sup>1</sup>**

**Note:** In no case should the amount of fumigant applied per acre exceed the gallons per acre rates for broadcast treatment given in Table 1.

Fl Oz/ 1000 Ft of Row	Plant Row Spacing (Inches)								
	28	32	36	40	44	48	52	56	60
	(Gallons per Acre)								
52	7.6	6.6	5.9	5.3	4.8	4.4	4.1	3.8	3.5
60	8.8	7.7	6.8	6.1	5.6	5.1	4.7	4.4	4.1
68	9.9	8.7	7.7	6.9	6.3	5.8	5.3	4.9	4.6
76	11.1	9.7	8.6	7.8	7.0	6.5	6.0	5.5	5.2
84	12.3	10.7	9.5	8.6	7.8	7.1	6.6	6.1	5.7
92	13.4	11.7	10.4	9.4	8.5	7.8	7.2	6.7	6.3
100	14.6	12.8	11.3	10.2	9.3	8.5	7.8	7.3	6.8
108	15.8	13.8	12.2	11.0	10.2	9.2	8.5	7.9	7.3
116	16.9	14.8	13.2	11.8	10.8	9.9	9.1	8.5	7.9
124	18.1	15.8	14.1	12.7	11.5	10.5	9.7	9.0	8.4
132	19.3	16.8	15.0	13.5	12.2	11.2	10.4	9.6	9.0
140	20.4	17.9	15.9	14.3	13.0	11.9	11.0	10.2	9.5
148	21.6	18.9	16.8	15.1	13.7	12.6	11.6	10.8	10.1
156	22.8	19.9	17.7	15.9	14.5	13.3	12.2	11.4	10.6
164	23.9	20.9	18.6	16.7	15.2	13.9	12.9	11.9	11.2
172	25.1	21.9	19.5	17.6	16.0	14.6	13.5	12.5	11.7
180	26.3	23.0	20.4	18.4	16.7	15.3	14.1	13.1	12.2
188	27.4	24.0	21.3	19.2	17.4	16.0	14.8	13.7	12.8
196	28.6	25.0	22.2	20.0	18.2	16.7	15.4	14.3	13.3
204	29.8	26.0	23.1	20.8	18.9	17.4	16.0	14.9	13.9
212	30.9	27.0	24.0	21.6	19.7	18.0	16.6	15.4	14.4

<sup>1</sup> For row spacing of 24 inches or less, apply as a broadcast treatment. For treatments with row spacing greater than 24 inches, refer to Table 1 for the rate needed for a specific crop and/or soil texture. To determine gallons per acre for row treatments, double the flow rate in Table 1 and look up the corresponding gallons per acre in Table 2.

For single chisel applications: the flow rates are double those listed in Table 1. For example, for vegetable crops in mineral soil, the flow rate for a single chisel row treatment is 52 to 70 fl oz per 1000 ft of row (note the broadcast rate is 26 to 35 fl oz per 1000 ft of row).

For multiple chisel applications: use the flow rate given in Table 1 per outlet. For example, for vegetable crops in mineral soil using 2 chisels per row, the flow rate per outlet is 52 to 70/2, or 26 to 35 fl oz per 1000 ft of row per outlet.

To obtain the gallons per acre used for a row spacing not shown in this table, use the following equation:

$$\frac{\text{fl oz/1000 ft of row}}{\text{row spacing (inches)}} \times 4.08^a = \text{gallons per acre}$$

$$4.08 = \frac{12 \text{ inches} \times 43.56 \text{ (no. 1000 ft/acre)}}{128 \text{ (fl oz per gallon)}}$$

## Plant Diseases

**Bacterial Canker of Peaches:** To aid in the control of this disease apply Telone II as a preplant broadcast treatment to light (sandy) soils at the rate of 35 gallons per acre (103 fl oz/1000 ft row/outlet) preferably in the fall when the soil is warm (55 to 80°F at injection depth) and moist. Inject the fumigant at a depth of 12 to 14 inches with chisels mounted on 12-inch centers.

**Fusarium Wilt of Cotton:** The effects of this disease can be suppressed by controlling the root knot nematodes associated with this disease/nematode complex. Use Telone II as a row treatment at the rate of 12 gallons per acre (35 fl oz/1000 ft row/outlet).

**Sugar Beet *Rhizomania* Disease:** Use Telone II to suppress the effects of this disease by preplant broadcast application at the rate of 10 to 18 gallons per acre broadcast equivalent (29 to 53 fl oz/1000 ft row/outlet). Use the higher rates for heavier (finer textured) soils and/or for higher levels of disease infestation. Telone II is believed to reduce the activity of *Polymyxa beta*, which has been identified as the vector of the *Rhizomania* disease virus.

**Verticillium Wilt of Mint:** To aid in the control of this disease, apply Telone II as a broadcast treatment at 25 to 30 gallons per acre (73 to 88 fl oz/1000 ft/outlet) in the spring, or preferably in the fall.

## Soil Insects

**Symphylans (Garden Centipedes):** Use Telone II for treatment of soil to be planted to crops where these pests have been shown to be a problem. Apply the fumigant only as a broadcast treatment at the rate of 18 to 35 gallons per acre (53 to 103 fl oz/1000 ft row/outlet) when soil temperature is warm (55 to 80°F) at the application depth.

**Wireworms:** Use Telone II for treatment of soil to be planted to crops where these pests have been shown to be a problem. Apply the fumigant as a broadcast treatment at 20 gallons per acre by injection at least 14 inches below the final soil surface.

For wireworm suppression control in Colorado, Idaho, Nevada, Oregon, Utah and Washington, and in Modoc and Siskiyou counties of California refer to supplemental labeling for Telone II entitled: "For the Control of Nematode Nematodes and the Suppression of Wireworm Wireworms Control in Soils to be Planted to Potatoes or Onions" for directions for use.

---

## Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

## Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

24 7 24

---

### **Limitation of Remedies**

---

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

\*Trademark of Dow AgroSciences LLC  
EPA-Accepted:   /  /