

62719-34

6-3-2003

1/2



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

JUN 3 2003

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

A. Sterett Robertson  
Regulatory Leader  
Regulatory Success-Americas  
Dow AgroSciences LLC  
9330 Zionsville, IN 46268-1054

Subject: Supplemental Labeling  
Lorsban 4E  
EPA Reg. No. 62719-34  
Submissions dated March 4, 2003.

Dear Mr. Robertson:

The labeling referred to above submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable with the following comment:

At your next label printing, or within eighteen months of the date of this letter, whichever comes first, you must incorporate this supplemental labeling into the main product labeling.

A stamped copy "accepted with comments" of the label is enclosed for your records.

Please submit two (2) copies of the final printed label for our file. If you have any questions please call me at (703) 308-8328.

Sincerely yours,

A handwritten signature in black ink, appearing to read "Akiva Abramovitch".

Akiva Abramovitch, Ph.D.

Chemist

Insecticide/Rodenticide Branch  
Registration Division (7505C)

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Lorsban\* 15G

EPA Reg. No. 62719-34

### Use of Lorsban 15G Prior to Post-Application of Lumax or Camix to Field Corn

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Lorsban\* 15G granular insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Lorsban 15G according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Lorsban 15G.

#### Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

Soil applied Lorsban 15G, as a T-band or in-furrow application, may be followed by post-applied Lumax or Camix herbicide, applied in accordance with their label recommendations. Apply Lorsban 15G at-plant at a rate of 8 to 12 oz per 1000 feet of row to control soil insects such as corn rootworm, cutworm, wireworm, white grub, seed corn maggot, and others listed on the product label.

**Do not** use Lorsban 15G prior to the use of Lumax or Camix for corn grown for seed production due to the unknown nature of the response of inbred lines to Lumax or Camix.

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated:**

JUN 3 2003

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.

62719-34

\*Trademark of Dow AgroSciences LLC

A1A / Lorsban 15G / Sec 3 Supp Lumax, Camix on Corn / 03-04-03

D06-061-XXX

EPA accepted:   /  /  

Initial printing.