

BEST AVAILABLE COPY

Specimen Label

ACCEPTED

APR 23 1991

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 62719-39

Lorsban* 50W

Wettable Powder Insecticide

For Control of Various Insects on Certain Fruit and Nut Crops

Active Ingredient(s):

Chlorpyrifos (O-diethyl O-(3,5,6-trichloro-2-pyridinyl) phosphorothioate) 50%

Inert Ingredient(s): 50%

E.P.A. Registration No. 464-552

E.P.A. Est. 477-TX-1

KEEP OUT OF REACH OF CHILDREN

WARNING

AVISO:

PRECAUCION AL USUARIO:

Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada en español.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

MAY BE FATAL IF SWALLOWED

• **HARMFUL IF ABSORBED**

THROUGH SKIN OR INHALED

• **CAUSES EYE IRRITATION**

Avoid Contact With Eyes, Skin, Or Clothing

• **Avoid Breathing Dust Or Spray Mist**

• **Wash Thoroughly With Soap And Water**

After Handling And Before Eating Or

Smoking • When Loading The Spray Tank

Or Handling The Powder, Wear A Pesticide

Respirator Jointly Approved By The Mining

Enforcement And Safety Administration

(Formerly The U.S. Bureau Of Mines) And

By The National Institute For Occupational

Safety And Health Under The Provisions of

30 CFR Part II • Wear Protective Clothing

Such As Long Pants, Long Sleeved Shirt,

Gloves, Hat, And Boots When Loading The

Spray Tank Or Handling The Powder And

During Application Of The Spray Mixture

• **Work To Windward To Stay Out Of Spray**

Drift Or Mist • Remove Contaminated

Clothing And Wash Before Reuse • Do Not

Apply This Product In Such a Manner as to

Directly or Through Drift Expose Workers or

Other Persons • The Area Being Treated

Must Be Vacated by Unprotected Persons

• **Keep Away From Food, Feedstuffs And**

Water Supplies

STATEMENT OF PRACTICAL TREATMENT:

If Swallowed: Call a physician or Poison Control Center.

Drink 1 or 2 glasses of water and induce vomiting by

touching back of throat with finger. Do not induce vomiting

or give anything by mouth to an unconscious person.

If On Skin: Wash with plenty of soap and water. Get

medical attention if irritation persists. **If In Eyes:** Flush with

plenty of water for 5 minutes. Get medical attention if

irritation persists. **If Inhaled:** Remove to fresh air if effects

occur. Get medical attention.

NOTE TO PHYSICIANS: Chlorpyrifos is a cholinesterase

inhibitor. Treat symptomatically. If exposed, plasma and

red blood cell cholinesterase tests may indicate

significance of exposure (baseline data are useful).

Atropine, only by injection, is the preferable antidote.

Oximes, such as 2-PAM/protopam, may be therapeutic

if used early; however, use only in conjunction with

atropine. In case of severe acute poisoning, use antidote

immediately after establishing an open airway and

respiration.

Environmental Hazards

This pesticide is toxic to birds and wildlife, and extremely

toxic to fish and aquatic organisms. Do not apply directly

to water. Drift and runoff from treated areas may be

hazardous to aquatic organisms in adjacent aquatic areas.

Cover or incorporate spills. Do not contaminate water

when disposing of equipment washwaters. This product is

highly toxic to bees exposed to direct treatment or

residues on blooming crops or weeds. Avoid use when

bees are actively foraging or if bees are likely to come in

contact with wet or re-wetted treated surfaces. Protective

information may be obtained from your Cooperative

Extension Service. This product may pose a hazard to

certain Federally endangered fish species. Refer to the

Endangered Species section of this label for any

geographic restrictions.

NOTICE

Read and understand the entire label before using. Use

only according to label directions.

Before buying or using this product, read **WARRANTY**

LIMITATIONS AND DISCLAIMER elsewhere on this label.

If terms are not acceptable, return unopened package or

order to seller for full refund of purchase price paid.

Otherwise, use by the buyer or any other user constitutes

acceptance of the terms under **WARRANTY / LIMITATIONS**

AND DISCLAIMER.

IN CASE OF AN EMERGENCY

endangering life or property involving this

product, call collect 817-638-4400

AGRICULTURAL CHEMICAL

Do Not Ship or Store with Food, Feeds,

Drugs, or Clothing

Lorsban* 50W

ENDANGERED SPECIES RESTRICTIONS

The use of LORSBAN 50W may pose a hazard to certain Federally endangered fish species. This product cannot be used on sour cherry trees in the following counties:

COLORADO: Delta, Larmer and Mesa

OHIO: Washington

UTAH: Emery, Grand, Utah and Washington

WISCONSIN: Iowa, Pierce, Richland and Vernon

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

General Information

LORSBAN 50W insecticide is a wettable powder formulation designed for use as a cover spray to control various pests. The pests controlled are listed in the accompanying table. LORSBAN 50W is compatible with insecticides, miticides, and fungicides commonly recommended in orchard spray schedules except for alkaline materials such as Bordeaux mixture and lime.

Test compatibility of the intended tank mixture before adding LORSBAN 50W to the spray or mix tank. Add proportionate amounts of each ingredient to a pint or quart jar, cap, shake, and let set 15 minutes. Formation of precipitates that do not readily redisperse indicates an incompatible mixture that should not be used.

Mixing Instructions

LORSBAN 50W mixes readily with water to form a suspension. Mix the required amount of LORSBAN 50W with a small quantity of water and add this premix while filling the spray tank. Alternatively, fill the tank with the required amount of water and then add the premix. Maintain sufficient agitation during both mixing and application to ensure uniformity of the spray mixture. LORSBAN 50W contains a wetting agent. Do not use additional wetting agents, spreaders or stickers.

Use and Dosage Recommendations

Use LORSBAN 50W at the dosages indicated by application as a foliar spray to control pests listed in the following table. Where a rate range is listed in the use directions, use the higher rate for more severe infestations. Mix the required dosage in sufficient water to ensure adequate coverage and apply as a concentrate or dilute spray using conventional, power-operated ground spray equipment or by aircraft (fixed-wing or helicopter). For dilute sprays applied to tree fruits and nuts, mix the required dosage in sufficient water to allow for spray to run off. For concentrate sprays, apply an equivalent amount of LORSBAN 50W per acre. Treat when pests appear or in accordance with local conditions. Mow or beat down orchard blooming weed cover crops before applying spray to tree fruit crops to reduce hazard to bees. This step is especially important in relation to early Spring cover sprays when morning conditions may be wet from dew. LORSBAN 50W will perform best in an almond or walnut pest control program that includes winter sanitation (removal of mummy nuts) and regular monitoring of target pests. To control apple achid infesting apples, spray every 10 to 14 days on a preventative schedule. To suppress pecan leaf scorch mite on pecans, use in a preventative program. For best pecan *Phylloxera* control, use two applications 7 to 10 days apart starting at bud swell and ending by early leaf expansion. For ant control on pecans, do not apply where weed growth or other obstructions would impede uniform coverage of the orchard floor. Consult your State Agricultural Experiment Station, Certified Pest Control Advisor, or Extension Service specialist for specific use information in your area.

LORSBAN 50W may also be used on almonds and walnuts as a dormant/delayed dormant treatment applied at the dosages indicated in a dilute or concentrate spray. Proper equipment must be used to obtain complete coverage of the entire bark area. If scale infestations are severe, dilute sprays should be used to ensure complete coverage. Use oil as recommended by your State Agricultural Experiment Station, State Extension Service Specialist, or Certified Pest Control Advisor. Do not use oil with LORSBAN 50W on walnuts. Because dry or cold conditions may cause LORSBAN 50W plus oil sprays to injure trees resulting in bud damage or drop, do not apply until winter rains or irrigation has replenished soil moisture such that bark and twigs are not desiccated.

BEST AVAILABLE COPY

APPLES: Use LORSBAN 50W to control the pests listed in the following table at the rates indicated.

Geographic Location	Pests Controlled	Dosage Recommendations	Remarks
<p>Restrictions (All use Areas) Make no more than 8 applications per season. Do not apply last treatment within 28 days before harvest or apply last two treatments closer than 21 days apart. Do not allow livestock to graze in treated orchards.</p> <p>Additional Restrictions (by Geographic Location) In California: Do not do work involving contact with trees within 2 days after treatment.</p>			
All Areas	Apple aphid ¹ , Apple maggot, Climbing cutworm, Codling moth, Dogwood borer (trunk spray), European apple sawfly, Eyespotted bud moth, European corn borer, Fruittree leafroller, Green fruitworm, Lesser appleworm, Obliquebanded leafroller, Orange tortrix, Pandemis leafroller, Oriental fruit moth, Periodical cicada, Plum curculio, Redbanded leafroller, Rosy apple aphid, San Jose scale, Tarnished plant bug, Tufted apple bud moth, Variegated leafroller, Woolly apple aphid.	3 lb/acre	*For apple aphid control LORSBAN 50W should not be used alone as a rescue treatment.
Connecticut, Delaware, Maine, Maryland, Massachusetts, Michigan, New Hampshire, New Jersey, New York, N. Carolina, Ohio, Pennsylvania, Rhode Island, South Carolina, Vermont, Virginia, West Virginia.	Apple aphid ¹ , Apple maggot ¹ , Codling moth, Dogwood borer (trunk spray) ¹ , European apple sawfly ¹ , European corn borer ¹ , European red mite ¹ , Eyespotted bud moth, Fruittree leafroller, Green fruitworm, Lesser appleworm, Obliquebanded leafroller ¹ , Oriental fruit moth, Periodical cicada ¹ , Plum curculio ¹ , Potato leafhopper, Redbanded leafroller, Rosy apple aphid, San Jose scale, Spotted tentiform leafminer ¹ , Tarnished plant bug, Tufted apple bud moth ¹ , Variegated leafroller, White apple leafhopper ¹ , Woolly apple aphid	2-3 lb/acre (8-12 oz/100 gal dilute) ¹ OR A tank mix of 1 1/2 lb/acre (6 oz/100 gal dilute) with one of the following: 1 1/2 pints/acre LANNATE 1.8L (6 fl oz/100 gal dilute) ¹ 6 oz/acre LANNATE 90SP (1 1/2 oz/ 100 gal dilute) ¹ 2 lb/acre IMIDAN 50W (8 oz/100 gal dilute) ¹ 1 1/2 lb/acre GUTHON 35W (6 oz/ 100 gal dilute) ¹ 2 pints/acre PENNCAP M (8 fl oz/100 gal dilute) ¹	*For apple aphid control, use 2-3 lbs/acre LORSBAN 50W or a tank mix with LANNATE 1.8L or LANNATE 90SP as recommended. LORSBAN 50W should not be used alone or in a tank mix as a rescue treatment. Treatments should be applied at regular intervals (1-2 weeks apart). *For apple maggot control in Michigan or in other areas after July, use 3 lbs/acre LORSBAN 50W or a tank mix with IMIDAN 50W, GUTHON 35W or PENNCAP M as recommended. *No tank mix recommended. *For European apple sawfly control, use 2-3 lbs/acre LORSBAN 50W or a tank mix with IMIDAN 50W, GUTHON or PENNCAP M as recommended. *For control of European corn borer or obliquebanded leafroller, use 2-3 lbs/acre LORSBAN 50W or a tank mix with LANNATE 1.8L or LANNATE 90SP as recommended. *European red mite suppression with LORSBAN 50W is achieved when a regularly scheduled preventative spray program is followed or when used with other mite management practices such as biological control. *For plum curculio control in Michigan, use 3 lbs/acre LORSBAN 50W or a tank mix with IMIDAN 50W, GUTHON 35W or PENNCAP M as recommended. For other areas, use 2-3 lbs/acre LORSBAN 50W or one of the above-mentioned tank mixes. *For control of spotted tentiform leafminer or white apple leafhopper, use a tank mix with LANNATE 1.8L or LANNATE 90SP as recommended. LORSBAN 50W alone is not recommended for control of these pests. *For tufted apple bud moth control, use 3 lbs/acre LORSBAN 50W or a tank mix with LANNATE 1.8L or LANNATE 90SP as recommended. *Based on 400 gal dilute/acre.

BEST AVAILABLE COPY

DOSAGE AND USE RECOMMENDATIONS
For LORSBAN 50W Wettable Powder Insecticide

Crop	Insects Controlled	Dosage LORSBAN 50W	Restrictions
Almonds	Nevel orangeworm, Peach twig borer, San Jose scale	4 lb per acre	Make only one dormant/delayed dormant spray application and no more than three foliar spray applications on almonds per season, one dormant/delayed dormant spray application and no more than two foliar spray applications on walnuts per season, and no more than three foliar spray applications on filberts per season. Do not apply within 14 days of harvest. Do not allow livestock to graze in treated orchards. Do not apply as a dormant spray application on almonds in the following counties in California: Butte, Colusa, Glenn, Solano, Sutter, Tehama, Yolo and Yuba.
Walnuts	Codling moth, Walnut scale, Walnut husk fly	4 lb per acre	
Filberts	Eyespotted bud moth, Filbert aphid, Filbert leafroller, Filbertworm Obliquibanded leafroller, Omnivorous leafminer, Winter moth	3 to 4 lb per acre	
Pecans	Black pecan aphid, Hickory shuckworm, Pecan leaf scorch mite (suppression)	2 lb per 100 gallons	Make no more than 5 applications per season. Do not apply within 28 days of harvest. Do not allow livestock to graze in treated orchards.
	Pecan nut casebearer, Phylloxera spp., Fall webworm, Spittlebug	1 lb per 100 gallons	
	Fire ants and other ant species (orchard floor)	2 lb per acre	
Braessca (cole) Broccoli Brussels sprouts Cabbage Cauliflower Collards Kale Kohlrabi	Cabbage aphid Cutworms Imported cabbageworm Striped flea beetle (adult)	2 lb per acre	Make no more than 6 applications per season. Do not apply within 21 days before harvest. To avoid phytotoxicity in vegetable crops, do not use with other pesticide products or treat plants that are under extreme heat and drought stress.
Sour Cherries	Borers American plum Lesser peachtree Pacific fletheaded Peach twig Peachtree Shot-hole Climbing cutworm Green fruitworm Leafrollers Fruitree Obliquibanded Pandemis Redbanded Lesser appleworm Tarnished plant bug Moth Eyespotted bud Miner Scale European fruit lecanium San Jose Stink bug	2 to 3 lb per acre	Make no more than 8 applications per season. Do not apply within 14 days of harvest. Do not allow livestock to graze in treated orchards. In California do not do work involving contact with trees within 2 days after treatment. Do not use on sweet cherries. Refer to the Endangered Species Section of this label for geographic restrictions on the use of LORSBAN 50W on sour cherries.
	Black cherry aphid Cherry fruit fly Plum curculio Rose chafar	3 lb per acre	

BEST AVAILABLE COPY

REENTRY STATEMENT

Do not allow reentry into treated areas within 24 hours unless protective clothing is worn. Protective clothing consists of at least a hat or other suitable head covering, a long-sleeved shirt and long-legged trousers or a coverall type garment (all of closely woven fabric covering the body, including the arms and legs), shoes and socks. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. These oral warnings shall inform workers of areas to be entered without protective clothing and actions to take in case of accidental exposure as shown by the Statement of Practical Treatment section on this label. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written warnings must include the following information: **WARNING.** Area treated with LORSBAN 50W wettable powder insecticide on (insert date of application). Do not enter without protective clothing for 24 hours. In case of accidental exposure (insert actions to take as shown by the Statement of Practical Treatment section on this label).

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY LIMITATIONS AND DISCLAIMER

The Dow Chemical Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions therein under normal conditions of use. **THIS IS THE ONLY WARRANTY MADE ON THIS PRODUCT. NO OTHER EXPRESS AND NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE IS MADE OUTSIDE OF THIS LABEL.** Therefore, neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), under abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes, etc.) or under conditions not reasonably foreseeable to or beyond the control of seller.

When buyer or user suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), buyer or user must promptly notify in writing The Dow Chemical Company of any claims to be eligible to receive either remedy given below. The **EXCLUSIVE REMEDY OF THE BUYER OR USER** and the **LIMIT OF LIABILITY** of The Dow Chemical Company or any other seller will be one of the following, at the election of The Dow Chemical Company:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

The seller will not be liable for consequential or incidental damages or losses.

The terms of this Warranty Limitations And Disclaimer cannot be varied by any written or verbal statements or agreements. Any employee or sales agent of the seller is not authorized to vary or exceed the terms of this Warranty Limitations And Disclaimer in any manner.

45386-L1

J989

THE DOW CHEMICAL COMPANY
Midland, Michigan 48674 U.S.A.
*Trademark of THE DOW CHEMICAL COMPANY

BEST AVAILABLE COPY