

(Product Label)

(Logo) DowElanco

Turflon* Ester

For the Control of Annual and Perennial Broadleaf Weeds and Kikuyugrass in Ornamental Turf

Active Ingredients:

Triclopyr: 3,5,6-trichloro-2-pyridinyloxyacetic acid, triethylamine salt.....	61.6%
Inert Ingredients	38.4%
Total.....	100.0%

Acid equivalent: Triclopyr - 44.3% - 4 lb/gal

Contains petroleum distillates

Keep Out of Reach of Children

CAUTION

AVISO:

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

See back panel for additional precautionary statements.

Notice: Read the entire label. Use only according to label directions. Before buying or using this product, read Warranty Disclaimer and Limitation of Remedies sections elsewhere on this label.

In case of emergency endangering life or property involving this product, call collect 517-636-4400

Agricultural Chemical: Do Not Ship or Store with Food, Feeds, Drugs, or Clothing.

EPA Reg. No. 62719-40


EPA Est. 464-MI-1

*Trademark of DowElanco

DowElanco • Indianapolis, IN 46268, U.S.A.

Herbicide

Net Contents - XX Gallons



Precautionary Statements

Hazards to Humans and Domestic Animals**CAUTION**

Harmful If Swallowed, Inhaled Or Absorbed Through The Skin

Avoid contact with eyes, skin or clothing. Avoid breathing mists or vapors. Avoid contamination of food. Wash thoroughly after handling. Remove and wash contaminated clothing before reuse.

First Aid

if case of skin contact, flush skin with plenty of water. Get medical attention if irritation persists. If Swallowed, do not induce vomiting. Call a physician.

Environmental Hazards

This pesticide is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water when disposing of equipment washwaters.

Physical or Chemical Hazards

Combustible - Do not use or store near heat or open flame. Do not cut or weld container.

Directions for Use

It is a violation of Federal Law to use this product in manner inconsistent with its labeling. Read All Directions Carefully Before Applying.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store above 28°F or agitate before use.

Pesticide Disposal: Pesticide spray mixture or rinsate water that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures.

Plastic containers may also be disposed of by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General: Consult federal, state, or local disposal authorities for approved alternative procedures.

General Information

Turfion Ester herbicide is recommended for the control of actively growing annual and perennial broadleaf weeds and kikuyugrass in perennial bluegrass, perennial ryegrass or tall fescue turf.

General Use Precautions (Read Carefully Before Using)

Local conditions may affect the use of herbicides. Consult your local specialist for advice in selecting treatments from this label to best fit local conditions.

Turfion Ester may injure certain turfgrass species. Do not apply to bahiagrass, bentgrass, bermudagrass, centipedegrass, St. Augustinegrass or zoysiagrass, unless turf injury can be tolerated.

Do not apply Turfion Ester herbicide directly to or otherwise permit it to come in contact with or permit spray mists containing Turfion Ester to drift onto cotton, grapes, tobacco, vegetable crops, flowers, fruit or orchard trees, shrubs or other desirable broadleaf plants.

Do not apply Turfion Ester to exposed roots of shallow rooted trees and shrubs.

BEST AVAILABLE COPY

For spot treatments, do not apply more than 2 quarts of Turflon Ester (400 gallons of spray) per acre in a single application.

Avoid Injurious Spray Drift: Applications should be made only when there is little or no hazard from spray drift. Very small quantities of spray, which may not be visible may seriously injure susceptible plants. Do not spray when wind is blowing toward susceptible crops or ornamental plants near enough to be injured. Spray drift can be reduced by spraying when wind velocity is low, keeping the spray boom as low as possible, using nozzles that deliver larger spray droplets, and by using higher volume sprays and lower pressures.

Do not apply to ditches used to transport irrigation water. Do not apply where runoff or irrigation water may flow onto susceptible crops.

Chemigation: Do not apply this product through any type of irrigation system.

Do not use this product for manufacturing or formulating.

Be sure that use of this product conforms to all applicable regulations.

Broadleaf weeds controlled by Turflon Ester include:

Black medic	Lespedeza
Bull thistle	Matchweed
Burdock	Mustard
Canada thistle	Oxalis
Chicory	Plantain
Clover	Ragweed
Creeping beggarweed	Smartweed
Curly dock	Sweet clover
Dandelion	Vetch
Field bindweed	Wild carrot
Goldenrod	(Queen Annes lace)
Ground ivy	Wild lettuce
Kikuyugrass†	Wild violet
Lambsquarters	Yarrow

†See control of Kikuyugrass under "Application Directions".

Mixing Instructions

When Turflon Ester is mixed with water it forms an emulsion (not a solution) and separation may occur unless the spray mixture is agitated continuously.

Add about one-half the required amount of clean water to the spray tank. Start agitation and add the recommended amount of Turflon Ester. Provide moderate agitation while completing the addition of water and during application.

Application Directions

Foliar sprays should be applied during warm weather, from early spring through fall, when weeds are actively growing. Application under drought conditions may provide less than desirable results. Use low pressure sprays to minimize spray drift. When making applications to mature plants, hard to control species, or during drought conditions, use higher rates.

Note: Broadleaf weeds germinate at different times. Only emerged weeds present at the time of application will be controlled. Newly seeded turf should be mowed two or three times before being treated. Do not water for 24 hours after application. Do not rese. 1 for 3 weeks after application.

BEST AVAILABLE COPY

Broadcast Treatment of Ornamental Turf

Apply 1/2 to 1 quart of Turflon Ester in 20 to 200 gallons of water per acre to control actively growing broadleaf weeds growing in perennial bluegrass, perennial ryegrass, or tall fescue. Do not use on other turfgrass species (see General Use Precautions) unless injury can be tolerated. To minimize turf injury, do not treat if turf is under heat or drought stress and make repeat applications at least four weeks apart.

Tank Mixing: To improve the spectrum of activity, Turflon Ester may be tank mixed at a rate of 1/2 to 1 pint per acre with recommended rates of low volatile amine or ester formulations of 2,4-D or other labeled postemergence broadleaf herbicides. Refer to tank mix product labels for specific use directions, precautions and limitations before use.

Spot Treatment of Ornamental Turf

Mix 1/4 to 1/2 fluid ounce of Turflon Ester in enough water to make three gallons of spray (1/2 to 1 pint per 100 gallons of spray) and apply at any time broadleaf weeds are susceptible by wetting foliage of susceptible plants to point of runoff. **Note:** Do not apply more than 2 quarts of Turflon Ester (400 gallons of spray) per acre in a single application.

Control of Kikuyugrass

Apply Turflon Ester at a rate of 1/2 to 1 quart per acre. To improve activity, MSMA herbicide may be tank mixed with the 1/2 quart per acre rate of Turflon Ester. Three to four additional applications at 4 to 6 week intervals may be required to achieve control of kikuyugrass.

WARRANTY DISCLAIMER

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. DOWELANCO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of DowElanco or the seller. All such risks shall be assumed by Buyer.

LIMITATION OF REMEDIES

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at DowElanco's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

DowElanco shall not be liable for losses or damages resulting from handling or use of this product unless DowElanco is promptly notified of such loss or damage in writing. In no case shall DowElanco be liable for consequential or incidental damages or losses. The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of DowElanco or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

rsb 1/30/92

BEST AVAILABLE COPY

Amendments


- 1) Label edited to reflect DowElanco transfer
- 2) Revised instructions to prevent spray drift.
- 3) Added tank mixing instructions for broadcast foliar treatments.
- 4) Added use instructions for control of kikuyugrass.

(Product Label)

(Logo) DowElanco

Turflon* Ester**For the Control of Annual and Perennial Broadleaf Weeds and Kikuyugrass in Ornamental Turf****Active Ingredients:**

Triclopyr: 3,5,6-trichloro-2-pyridinyloxyacetic acid, triethylamine salt.....	61.6%
Inert ingredients	38.4%
Total.....	100.0%

Acid equivalent: Triclopyr - 44.3% - 4 lb/gal**Contains petroleum distillates****Keep Out of Reach of Children****CAUTION****AVISO:****PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.****See back panel for additional precautionary statements.****Notice: Read the entire label. Use only according to label directions. Before buying or using this product, read Warranty Disclaimer and Limitation of Remedies sections elsewhere on this label.****In case of emergency endangering life or property involving this product, call collect 517-636-4400****Agricultural Chemical: Do Not Ship or Store with Food, Feeds, Drugs, or Clothing.****EPA Reg. No. 62719-40****EPA Est. 464-MI-1*****Trademark of DowElanco****DowElanco • Indianapolis, IN 46268, U.S.A.****Herbicide****Net Contents - XX Gallons**

2 of 8

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful If Swallowed, Inhaled Or Absorbed Through The Skin

Avoid contact with eyes, skin or clothing. Avoid breathing mists or vapors. Avoid contamination of food. Wash thoroughly after handling. Remove and wash contaminated clothing before reuse.

First Aid

If case of skin contact, flush skin with plenty of water. Get medical attention if irritation persists. If Swallowed, do not induce vomiting. Call a physician.

Environmental Hazards

This pesticide is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water when disposing of equipment washwaters.

Physical or Chemical Hazards

Combustible - Do not use or store near heat or open flame. Do not cut or weld container.

Directions for Use

It is a violation of Federal Law to use this product in manner inconsistent with its labeling. Read All Directions Carefully Before Applying.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store above 28°F or agitate before use.

Pesticide Disposal: Pesticide spray mixture or rinsate water that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures.

Plastic containers may also be disposed of by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General: Consult federal, state, or local disposal authorities for approved alternative procedures.

General Information

Turfion Ester herbicide is recommended for the control of actively growing annual and perennial broadleaf weeds and kikuyugrass in perennial bluegrass, perennial ryegrass or tall fescue turf.

General Use Precautions (Read Carefully Before Using)

Local conditions may affect the use of herbicides. Consult your local specialist for advice in selecting treatments from this label to best fit local conditions.

Turfion Ester may injure certain turfgrass species. Do not apply to bahiagrass, bentgrass, bermudagrass, centipedegrass, St. Augustinegrass or zoysiagrass, unless turf injury can be tolerated.

Do not apply Turfion Ester herbicide directly to or otherwise permit it to come in contact with or permit spray mists containing Turfion Ester to drift onto cotton, grapes, tobacco, vegetable crops, flowers, fruit or orchard trees, shrubs or other desirable broadleaf plants.

Do not apply Turfion Ester to exposed roots of shallow rooted trees and shrubs.

BEST AVAILABLE COPY

For spot treatments, do not apply more than 2 quarts of Turflon Ester (400 gallons of spray) per acre in a single application.

Avoid Injurious Spray Drift: Applications should be made only when there is little or no hazard from spray drift. Very small quantities of spray, which may not be visible may seriously injure susceptible plants. Do not spray when wind is blowing toward susceptible crops or ornamental plants near enough to be injured. Spray drift can be reduced by spraying when wind velocity is low, keeping the spray boom as low as possible, using nozzles that deliver larger spray droplets, and by using higher volume sprays and lower pressures.

Do not apply to ditches used to transport irrigation water. Do not apply where runoff or irrigation water may flow onto susceptible crops.

Chemigation: Do not apply this product through any type of irrigation system.

Do not use this product for manufacturing or formulating.

Be sure that use of this product conforms to all applicable regulations.

Broadleaf weeds controlled by Turflon Ester include:

Black medic	Lespedeza
Bull thistle	Matchweed
Burdock	Mustard
Canada thistle	Oxalis
Chicory	Plantain
Clover	Ragweed
Creeping beggarweed	Smartweed
Curly dock	Sweet clover
Dandelion	Vetch
Field bindweed	Wild carrot
Goldenrod	(Queen Annes lace)
Ground ivy	Wild lettuce
Kikuyugrass†	Wild violet
Lambsquarters	Yarrow

†See control of Kikuyugrass under "Application Directions".

Mixing Instructions

When Turflon Ester is mixed with water it forms an emulsion (not a solution) and separation may occur unless the spray mixture is agitated continuously.

Add about one-half the required amount of clean water to the spray tank. Start agitation and add the recommended amount of Turflon Ester. Provide moderate agitation while completing the addition of water and during application.

Application Directions

Foliar sprays should be applied during warm weather, from early spring through fall, when weeds are actively growing. Application under drought conditions may provide less than desirable results. Use low pressure sprays to minimize spray drift. When making applications to mature plants, hard to control species, or during drought conditions, use higher rates.

Note: Broadleaf weeds germinate at different times. Only emerged weeds present at the time of application will be controlled. Newly seeded turf should be mowed two or three times before being treated. Do not water for 24 hours after application. Do not reseed for 3 weeks after application.

BEST AVAILABLE COPY

Broadcast Treatment of Ornamental Turf

Apply 1/2 to 1 quart of Turflon Ester in 20 to 200 gallons of water per acre to control actively growing broadleaf weeds growing in perennial bluegrass, perennial ryegrass, or tall fescue. Do not use on other turfgrass species (see General Use Precautions) unless injury can be tolerated. To minimize turf injury, do not treat if turf is under heat or drought stress and make repeat applications at least four weeks apart.

Tank Mixing: To improve the spectrum of activity, Turflon Ester may be tank mixed at a rate of 1/2 to 1 pint per acre with recommended rates of low volatile amine or ester formulations of 2,4-D or other labeled postemergence broadleaf herbicides. Refer to tank mix product labels for specific use directions, precautions and limitations before use.

Spot Treatment of Ornamental Turf

Mix 1/4 to 1/2 fluid ounce of Turflon Ester in enough water to make three gallons of spray (1/2 to 1 pint per 100 gallons of spray) and apply at any time broadleaf weeds are susceptible by wetting foliage of susceptible plants to point of runoff. Note: Do not apply more than 2 quarts of Turflon Ester (400 gallons of spray) per acre in a single application.

Control of Kikuyugrass

Apply Turflon Ester at a rate of 1/2 to 1 quart per acre. To improve activity, MSMA herbicide may be tank mixed with the 1/2 quart per acre rate of Turflon Ester. Three to four additional applications at 4 to 6 week intervals may be required to achieve control of kikuyugrass.

WARRANTY DISCLAIMER

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. DOWELANCO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of DowElanco or the seller. All such risks shall be assumed by Buyer.

LIMITATION OF REMEDIES

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at DowElanco's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

DowElanco shall not be liable for losses or damages resulting from handling or use of this product unless DowElanco is promptly notified of such loss or damage in writing. In no case shall DowElanco be liable for consequential or incidental damages or losses. The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of DowElanco or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

rsb 1/30/92

BEST AVAILABLE COPY

Amendments

- 1) Label edited to reflect DowElanco transfer
- 2) Revised instructions to prevent spray drift.
- 3) Added tank mixing instructions for broadcast foliar treatments.
- 4) Added use instructions for control of kikuyugrass.



Turflon* Ester

Specialty Herbicide

For the Control of Annual and Perennial Broadleaf Weeds and Kikuyugrass in Ornamental Turf

Active Ingredients:

Triclopyr: 3,5,6-trichloro-2-pyridinyloxyacetic acid, triethylamine salt61.6%
Inert Ingredients38.4%
Total100.0%

Acid equivalent: Triclopyr - 44.3% - 4 lb/gal
Contains petroleum distillates

EPA Reg. No. 62719-40
EPA Est. 464-MI-1
Net Contents 2.5 gal

Precautionary Statements

Hazards to Humans and Domestic Animals
Keep Out Of Reach Of Children

CAUTION PRECAUCION:

Precaucion al usuario: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada apropiadamente.

Harmful If Swallowed, Inhaled Or Absorbed Through The Skin

Avoid contact with eyes, skin or clothing. Avoid breathing mists or vapors. Avoid contamination of food. Wash thoroughly after handling. Remove and wash contaminated clothing before reuse.

First Aid

If case of skin contact: Flush skin with plenty of water. Get medical attention if irritation persists.
If Swallowed: DO NOT induce vomiting. Call a physician.

Environmental Hazards

This pesticide is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water when disposing of equipment washwaters.

Physical or Chemical Hazards

Combustible - Do not use or store near heat or open flame. Do not cut or weld container.

Notice: Read the entire label. Use only according to label directions. Before buying or using this product, read Warranty Disclaimer and Limitation of Remedies sections elsewhere on this label.

In case of emergency endangering life or property involving this product, call collect 517-636-4400

Agricultural Chemical: Do Not Ship or Store with Food, Feeds, Drugs, or Clothing.

In Accordance with FR Notice 12-
Based on Draft Labeling Dated 1/29/92

Turflon* Ester

Directions for Use

It is a violation of Federal Law to use this product in manner inconsistent with its labeling.
Read All Directions Carefully Before Applying.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store above 20°F or agitate before use.

Pesticide Disposal: Pesticide spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures. Plastic containers may also be disposed of by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General: Consult federal, state, or local disposal authorities for approved alternative procedures.

General Information

Turfion Ester herbicide is recommended for the control of actively growing annual and perennial broadleaf weeds and kikuyugrass in perennial bluegrass, perennial ryegrass or tall fescue turf.

General Use Precautions

(Read Carefully Before Using)

Local conditions may affect the use of herbicides. Consult your local specialist for advice in selecting treatments from this label to best fit local conditions.

Turfion Ester may injure certain turfgrass species. Do not apply to bahiagrass, bentgrass, bermudagrass, centipedegrass, St. Augustinegrass or zoysiagrass, unless turf injury can be tolerated.

Do not apply Turfion Ester herbicide directly to or otherwise permit it to come in contact with or permit spray mists containing Turfion Ester to drift onto cotton, grapes, tobacco, vegetable crops, flowers, fruit or orchard trees, shrubs or other desirable broadleaf plants.

Do not apply Turfion Ester to exposed roots of shallow rooted trees and shrubs.

For spot treatments, do not apply more than 2 quarts of Turfion Ester (400 gallons of spray) per acre in a single application.

Avoid Injurious Spray Drift: Applications should be made only when there is little or no hazard from spray drift. Very small quantities of spray, which may not be visible may seriously injure susceptible plants. Do not spray when wind is blowing toward susceptible crops or ornamental plants near enough to be injured. Spray drift can be reduced by spraying when wind velocity is low, keeping the spray boom as low as possible, using nozzles that deliver larger spray droplets, and by using higher volume sprays and lower pressures.

Do not apply to ditches used to transport irrigation water. Do not apply where runoff or irrigation water may flow onto susceptible crops.

Chemigation: Do not apply this product through any type of irrigation system.

Do not use this product for manufacturing or formulating.

Be sure that use of this product conforms to all applicable regulations.

Broadleaf weeds controlled by Turfion Ester include:

Black medic	Lespedeza
Bull thistle	Matchweed
Burdock	Mustard
Canada thistle	Oxalis
Chicory	Plantain
Clover	Ragweed
Creeping beggarweed	Smartweed
Curly dock	Sweet clover
Dandelion	Vetch
Field bindweed	Wild carrot
Goldenrod	(Queen Anne's lace)
Ground ivy	Wild lettuce
Kikuyugrass†	Wild violet
Lambsquarters	Yarrow

† See control of Kikuyugrass under "Application Directions".

Mixing Instructions

When Turfion Ester is mixed with water it forms an emulsion (not a solution) and separation may occur unless the spray mixture is agitated continuously.

Add about one-half the required amount of clean water to the spray tank. Start agitation and add the recommended amount of Turfion Ester. Provide moderate agitation while completing the addition of water and during application.

Application Directions

Foliar sprays should be applied during warm weather, from early spring through fall, when weeds are actively growing. Application under drought conditions may provide less than desirable results. Use low pressure sprays to minimize spray drift. When making applications to mature plants, hard to control species, or during drought conditions, use higher rates.

Note: Broadleaf weeds germinate at different times. Only emerged weeds present at the time of application will be controlled. Newly seeded turf should be mowed two or three times before being treated. Do not water for 24 hours after application. Do not reseed for 3 weeks after application.

Broadcast Treatment of Ornamental Turf

Apply 1/2 to 1 quart of Turfion Ester in 20 to 200 gallons of water per acre to control actively growing broadleaf weeds growing in perennial bluegrass, perennial ryegrass, or tall fescue. Do not use on other turfgrass species (see General Use Precautions) unless injury can be tolerated. To minimize turf injury, do not treat if turf is under heat or drought stress and make repeat applications at least four weeks apart.

Tank Mixing: To improve the spectrum of activity, Turfion Ester may be tank mixed at a rate of 1/2 to 1 pint per acre with recommended rates of low volatile amine or ester formulations of 2,4-D or other labeled post-emergence broadleaf herbicides. Refer to tank mix product labels for specific use directions, precautions and limitations before use.

Spot Treatment of Ornamental Turf

Mix 1/4 to 1/2 fluid ounce of Turfion Ester in enough water to make three gallons of spray (1/2 to 1 pint per 100 gallons of spray) and apply at any time broadleaf weeds are susceptible by wetting foliage of susceptible plants to point of runoff. **Note:** Do not apply more than 2 quarts of Turfion Ester (400 gallons of spray) per acre in a single application.

Control of Kikuyugrass

Apply Turfion Ester at a rate of 1/2 to 1 quart per acre. To improve activity, MSMA herbicide may be tank mixed with the 1/2 quart per acre rate of Turfion Ester. Three to four additional applications at 4 to 6 week intervals may be required to achieve control of kikuyugrass.

BEST AVAILABLE COPY

WARRANTY DISCLAIMER

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. DOWELANCO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of DowElanco or the seller. All such risks shall be assumed by Buyer.

LIMITATION OF REMEDIES

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at DowElanco's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought,
or
- (2) Replacement of amount of product used.

DowElanco shall not be liable for losses or damages resulting from handling or use of this product unless DowElanco is promptly notified of such loss or damage in writing. In no case shall DowElanco be liable for consequential or incidental damages or losses. The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of DowElanco or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

*Trademark of DowElanco
DowElanco • Indianapolis, IN 46268, U.S.A.

LABEL CODE 113-64-008 DATE CODE 992
EPA APPROVAL 06/29/92 REPLACES 113-64-007

Revisions Include:

- 1) Revised instructions to prevent spray drift.
- 2) Added tank mixing instructions for broadcast foliar treatments.
- 3) Added use instructions for control of kikuyugrass.