

PM 19

62719-72

Yellow Copy
10/12

NOV 13 1992

Mr. Michael W. Melichar
DowElanco
Quad IV--9002 Purdue Road
Indianapolis, Indiana 46268-1189

Subject: Revised Labeling--User Limitation
Dursban 50W in Water Soluble Packets
EPA Registration Number 62719-72
Your Application (Fax) Dated November 3, 1992

Dear Mr. Melichar:

Your proposed change of the user limitation phrase immediately following the product name on the first page of the label is acceptable. You may delete the phrase as it now appears and substitute the following (your "Insert 1."):

To be applied only by or under the supervision of pest control operators or other trained personnel responsible for insect control programs.

The proposed changes in the directions for "Pest Control on Outside Surfaces, Around and Under Structures or in Crawl Spaces" (your "Inserts 2. and 3.") are not acceptable. We feel that the proposed directions could be interpreted as allowing excessive application of the product. You have indicated that you are willing to postpone resolution of this issue.

Accordingly, we are conditionally accepting the labeling. That is, it is acceptable provided you make no changes in the "Pest Control on Outside Surfaces, Around and Under Structures or in Crawl Spaces" section as it now appears on the label.

Sincerely yours,

DHE

Dennis H. Edwards, Jr.
Product Manager (19)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosure

DowElanco

Dursban* 50W

In Water Soluble Packets

Specialty Insecticide

Insert 1.

To be applied only by or under the direct supervision of commercial applicators responsible for insect control programs

Active Ingredient:

chlorpyrifos: O,O-diethyl O-(3,5,6-trichloro-2-pyridinyl) phosphorothioate.....	50%
Inert Ingredient.....	50%

EPA Registration No. 62719-72

EPA Est. 33967-NJ-1

Net Contents 4 oz *Trademark of DowElanco

Precautionary Statements

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

WARNING AVISO:

Precaucion al usuario: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

May Be Fatal If Swallowed • Harmful If Inhaled Or Absorbed Through Skin • Causes Eye Irritation

Do not get in eyes, on skin or clothing. Avoid breathing dust and spray mist. Wash thoroughly with soap and water after handling and before eating or smoking. Remove contaminated clothing and wash before reuse. Keep away from food, feedstuffs, and water supplies.

First Aid

If swallowed: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not give anything by mouth to an unconscious person.

If on skin: Wash with plenty of soap and water. Get medical attention.

If in eyes: Flush with plenty of water for 5 minutes. Get medical attention if irritation persists.

If inhaled: Remove to fresh air if symptoms of cholinesterase inhibition appear and get medical attention immediately.

Note to physician: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate significance of exposure (baseline data are useful). Atropine, only by injection, is the preferable antidote. Oximes, such as 2-PAM/protopam, may be therapeutic if used early; however, use only in conjunction with atropine. In case of severe acute poisoning, use antidote immediately after establishing an open airway and respiration.

Environmental Hazards

This pesticide is toxic to birds and wildlife, and extremely toxic to fish and aquatic organisms. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Cover or incorporate spills. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Notice: Read the entire label. Use only according to label directions. **Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.**

In case of emergency endangering health or the environment involving this product, call collect 517-636-4400.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

REGISTERED
with COMMENTS
in EPA Letter Dated

NOV 13 1992

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

Dursban* 50W In Water Soluble Packets

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system. Do not formulate this product into other end-use products.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers. Do not store above 122°F for extended periods of time. If container is damaged or spill occurs, use product immediately or dispose of product and damaged container as indicated below. Packets may become brittle when stored below 32°F. Handle carefully to avoid breakage.

Pesticide Disposal: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: When all packets are used, dispose of empty package in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General Information

This outer protective package contains four 4 oz water-soluble packets. Do not allow packets to become wet prior to adding to the spray tank. Wear gloves when handling water-soluble packets. Do not handle with wet gloves. Do not excessively handle water-soluble packets since this may cause breakage. Reseal outer package to protect remaining packets. Packets may become brittle when stored below 32°F. Handle carefully to avoid breakage.

Dursban 50W insecticide is a formulation designed for use as a spray to control various pests injurious to trees, turf and ornamental plants. The pests controlled are listed in the accompanying tables. Dursban 50W is compatible with insecticides, miticides and fungicides commonly recommended except for alkaline materials such as Bordeaux mixture and lime. A small amount of spray mixture should be prepared to check for compatibility before a large volume of spray is mixed. Dursban 50W mixes readily with water to form a suspension. Dursban 50W contains a wetting agent. Do not use additional wetting agents, spreaders or stickers.

Applicator Safe Handling Procedures

When exposure to spray mist is likely to occur, wear protective equipment and clothing (eye protections, long-sleeved coveralls and water-proof gloves) to prevent exposure of eyes and skin. To prevent breathing of spray mist during application in confined areas, wear a respirator and cartridge(s) approved by NIOSH/MSHA for pesticides.

Attention: Do not allow adults, children or pets on treated areas until spray has dried. Keep out of fishpools and other bodies of water. Do not treat vegetable gardens. Do not allow livestock to graze in treated areas. Do not feed treated grass cuttings (hay) or seed screenings to livestock nor use hay for livestock bedding. Do not use in poultry houses.

Mixing Directions

For paddle, sparger and bypass agitation systems.

1. Fill the spray tank 1/3 full or water.
Note: For paddle agitation systems, paddles should be covered with water.
2. Activate agitation system.
3. Add required number of water-soluble packets to obtain desired dilution.
4. Complete tank filling while the packet dissolves and Dursban 50W disperses.
5. Before beginning to spray, make sure Dursban 50W has dispersed and water-soluble packet has dissolved.

Note: Depending on the water temperature and the degree of agitation, the packets and Dursban 50W should be completely dispersed within approximately 5 minutes from the time they were added to the water.

Approved Uses

Turf and Other Outdoor Uses

Use Dursban 50W to control pests infesting lawns, golf courses, parks, perennial turf grasses grown for seed or sod, etc., listed in the following table by application at the recommended dosages. Dilute Dursban 50W in enough water to obtain complete and uniform coverage of pest-infested areas and apply 40 to 200 gallons of water per acre in a manner to obtain complete and uniform coverage of pest-infested areas. Use a coarse, low-pressure spray using suitable application equipment. For best results, the lawn should be moist at time of treatment. Spray when pests first appear; re-treat when needed, but not more often than once every 7 days.

Pest†	Amount of Dursban 50W per Acre	Specific Directions
ants (including foraging fire ants)	2 lb (eight 4 oz packets)	1. Use Dursban 50W for area control of ticks and chiggers infesting non-cropland areas such as roadsides, footpaths, and trails, picnic and camping sites, parks and other recreational areas where these pests are present and create a nuisance or a possible public health problem. Do not allow public use of treated areas during application or until spray has dried. Apply Dursban 50W in water at the rate of 1/2 lb/acre (equivalent to 1/2 oz per 2700 sq ft) using a hydraulic sprayer, mist applicator, knapsack sprayer or other suitable hand- or power-operated spray equipment. Treat low underbrush, grassy areas, weeds, ground surfaces and debris using enough spray volume to obtain thorough coverage, usually 40 to 100 gallons per acre.
armyworms (beet) (fall) (yellowstriped)		
centipedes	leafhoppers	2. Mosquitoes coming to rest on areas treated for control of turf pests will be killed for varying periods of time after treatment, depending on exposure of treated areas to weathering conditions.
chiggers	Lucerne moths	
chinch bugs	millipedes	
clover mites	mites	
cutworms	(clover)	
crickets	(Bermuda grass stunt)	
earwigs	(winter grain)	
fiery skipper	mosquitoes	
fleas	pillbugs	
gnats	sod webworm	
grasshoppers	(lawn moths)	
greenbugs	sowbugs	
June beetles	springtails	
	ticks	

† Superscripts refer to "Specific Directions"

Pest†	Amount of Dursban 50W per Acre	Specific Directions
European crane fly	3 lb (twelve 4 oz packets)	
Turfgrass weevil (<i>Hyperodes</i>)	4 lb (sixteen 4 oz packets)	1. Spray suspected problem areas in mid-April and again in mid-May or as recommended by your Local Agricultural Extension Service Specialist.
White grubs (black turfgrass <i>ateneus</i>) (European chafer) (southern masked chafer) (Japanese beetle larvae)	4 to 8 lb (sixteen to thirty-two 4 oz packets)	1. For white grubs, spray when grubs are young and actively feeding near the soil surface, usually during late July and August or as recommended by your Local Agricultural Extension Service Specialist. Immediately after spraying, irrigate the treated area with 1/2 to 1 inch of water to wash the insecticide into the underlying soil.
Bluegrass billbugs	2 to 8 lb (eight to thirty-two 4 oz packets)	1. For bluegrass billbugs, spray early in the season when adults first appear. Re-treat as needed.
Mole crickets	4 to 6 lb (sixteen to twenty-four 4 oz packets)	1. Mole crickets: Apply Dursban 50W in a minimum spray volume of 50 gallons of water per acre using broadcast or suitable hand held application equipment. Apply when early stage nymphs are active. Efficacy may be enhanced by spraying late in the afternoon or early evening and irrigating the turf prior to treatment to move mole crickets close to the soil surface.
Imported fire ants in commercial sod	16 lb (sixty-four 4 oz packets)	1. Imported fire ants: Apply Dursban 50W as a spray to the area of sod to be cut. Immediately after applying the spray soak in with at least 1/2 acre-inch of water or a sufficient volume to thoroughly soak below the cut line. Do not apply a volume or rate of water to cause runoff. Sod can be cut when vegetation has dried and when soil is moist but not wet. Precaution: Personnel handling treated sod should wear elbow-length chemically resistant gloves and apron. Protective clothing and boots should be worn to prevent repeated skin exposure to treated sod. Certification Period: 42 days from day of application.
Chiggers (turkey pens)	8 lb (thirty-two 4 oz packets)	1. For control of chiggers infesting outdoor turkey pens, apply Dursban 50W as a spray to the soil using 100 to 150 gallons of water per acre. To prevent turkeys from becoming infested with chiggers, spray the soil in pens just before turkeys are transferred to them. For turkeys already infested with chiggers, spray the soil in pens thoroughly. Allow at least three weeks for chigger lesions on the turkeys to heal before turkeys are slaughtered. For optimum results, spray a 10 to 15 ft protective area around the periphery of the treated pens. Treatment is limited to 2 applications to the soil in turkey pens at intervals of not less than 4 weeks. Do not treat soil in turkey pens later than 7 days before turkeys are to be slaughtered. Avoid getting the spray on feed or in water troughs. Open feed should be covered during spraying operations, and water troughs should be flushed out immediately after the spraying operations. Do not spray Dursban 50W directly on the turkeys.

† Superscripts refer to "Specific Directions"

Insert 2.

Pest Control on Outside Surfaces of and Around Buildings
 Apply Dursban 50W by application as a residual spray to outside surfaces of buildings including porches, window frames, eaves, patios, garages, refuse dumps and other areas where pests congregate or have been seen. Repeat treatment as needed to maintain effectiveness. Avoid application to surface where visible residues are objectionable.

Pest†	Amount of Dursban 50W in Water to Make 50 Gallons of Diluted Spray	Specific Directions
ants bees boxelder bugs carpenter ants clover mites cockroaches crickets earwigs elm leaf beetles (adults) firebrats fleas flies hornets millipedes mosquitoes pillbugs scorpions silverfish sowbugs spiders springtails symphylans ticks wasps yellowjackets	Perimeter Treatment	1. To help prevent infestation of buildings, treat a band of soil 6 to 10 feet wide around and adjacent to buildings, also the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use 4 to 16 ounces (one to four 4 oz packets) of Dursban 50W per 50 gallons of water and apply as a coarse spray at the rate of about 10 gallons spray mixture per 1 000 sq ft to thoroughly and uniformly wet the band area. 2. For scorpions, remove accumulations of lumber, firewood and other materials which serve as harborage sites. Dursban 50W may be applied as a residual spray to surfaces immediately below such materials. Perimeter treatments may reduce pest immigration from surrounding areas.
	1/4 to 1 lb (one to four 4 oz packets)	
	On Outside Surfaces	
	4 lb (sixteen 4 oz packets)	

Insert 3. →

† Superscripts refer to "Specific Directions".

Ornamentals (Outdoor, Greenhouse, Shadehouse, Nurseries, etc.)


Use Dursban 50W to treat flowers, shrubs, evergreens, vines and shade and flowering trees found to be infested with pests listed in the following table. Dilute Dursban 50W with water according to directions given in the table and apply using suitable hand- or power-operated spray equipment in a manner to provide complete and uniform coverage. For best results apply a coarse spray to thoroughly wet both upper and lower leaf surfaces and infested limb and trunk areas. Attempt to penetrate dense foliage but avoid overspraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals, if needed.

Consult your State Agricultural Experiment Station or Extension Service Specialist for application timing and other specific use information.

Note for Greenhouse Grown Ornamentals: Environmental factors significantly affect phytotoxicity. While Dursban 50W has been tested on numerous ornamental plants without causing serious phytotoxicity, because of the numerous varieties grown, it is recommended that a small group of plants be treated at the recommended rate under the expected growing conditions and observed for phytotoxic symptoms. Two test sprays should be made 7 days apart and the plants observed for another 7 days after the second treatment. Significant phytotoxicity has been observed in the following species following treatment with Dursban 50W. Apply to these species at your own risk.

Plant Species

Common Name	Scientific Name
copperleaf	<i>Acalypha wilkesiana</i>
zebra plant	<i>Aphelandra squarosa</i>
schefflera	<i>Brassia actinophylla</i>
black olive	<i>Bucca buceras</i>
papayas	<i>Carica papaya</i>
ossus	<i>Cissus antarctica</i>
croton	<i>Codiaeum variegatum</i>
figus	<i>Ficus nitida</i>
weeping fig	<i>Ficus pumila</i>
Cuban laurel	<i>Ficus religiosa</i>
yellow hibiscus	<i>Hibiscus cadybeus</i>
red/Chinese hibiscus	<i>Hibiscus rosa-sinensis</i>
impatiens	<i>Impatiens</i> spp.
Boston fern, fluffly ruffe fern	<i>Nephrolepis exaltata</i>
petunia	<i>Petunia</i> spp.
Belinda, caramia, lavande, Jack Frost	<i>Rosa</i> spp.



Pest†	Amount of Dursban 50W in Water to Make:		Specific Directions	
	50 gal	100 gal		
adelgids (cooley spruce gall) (eastern spruce gall) (pine bark) ants (including foraging fire ants) aphids (apple) (chrysanthemum) (cottonwood) (elm leaf) (peach) (rose) (spirea) armyworms (fall) (yellowstriped) bagworms boxelder bugs cankerworms catalpa sphinx chiggers citrus mealybugs clover mite cockroaches (American) (Asian wood) (smokey brown) elm spanworms fall webworms ¹ grasshoppers green fruitworm hornworms jackpine budworm juniper webworms katydids	lace bugs leafhoppers leafrollers maple leafcutter mites (red spider) (southern red) oleander caterpillars orange tortrix periodical cicada plant bugs poplar tentmaker phyllids puss caterpillars rose chafer sawflies, exposed (pine) (pin oak) (redheaded) sowbugs spittlebugs spring elm caterpillars spruce budworms tent caterpillars (eastern) (forest) (western) thornbugs walnut caterpillars western spruce budworms whiteflies yellownecked caterpillars	1/4 lb (one 4 oz packet)	1/2 lb (two 4 oz packets)	<ol style="list-style-type: none"> 1. For bagworms, treat when insects are in the crawler stage. 2. For effective control of leafrollers, spray should be applied before leaves are tightly rolled. 3. For effective control of maple leafcutter on maple trees, apply spray to larvae as cases are being formed. Do not treat sugar maple trees intended for maple syrup production. 4. For effective control of spider mites when large numbers of eggs are present, apply a second spray 3 to 5 days in the south or 7 to 10 days in the north after initial treatment to control newly-hatched nymphs. 5. For effective control of Fall webworm, direct spray into web and immediately surrounding foliage.
armyworms (beet) beetles (native elm bark) cutworms leafhoppers mahogany webworms mealybugs mimosa webworms moths (browntail) (Cypress tip) (Douglas fir tussock) (European pine shoot) (gypsy) ² (holly bug) (Nantucket pine tip) (Pandora) (pitch pine tip) (subtropical pine tip) (tussock) oakworms (California orangestriped) (redhumped) redhumped caterpillars thrips (exposed) weevils (blackvine) ¹ (yellow poplar) (pine reproduction)	1/2 lb (two 4 oz packets)	1 lb (four 4 oz packets)	<ol style="list-style-type: none"> 1. To reduce twig and branch feeding by bark beetles, applications should be made in the spring or early summer. 2. To control migrating and invading gypsy moth larvae, treat trunks and foliage. 3. Blackvine weevils are night feeders. Late afternoon spraying will give control in some areas. 	

† Superscripts refer to "Specific Directions"

Pest†	Amount of Dursban 50W in Water to Make:		Specific Directions
	50 gal	100 gal	
foliar feeding beetles (blister) (cottonwood leaf) ¹ (elm leaf) (flea) (fuller rose) (Japanese) (June) (willow leaf)	1/2 to 1 lb (two to four 4 oz packets)	1 to 2 lb (four to eight 4 oz packets)	1. For cottonwood leaf beetles, use Dursban 50W in water to control larvae and adults infesting cottonwoods. Make the treatment when field counts indicate damaging beetle populations are developing or present.
borers: clearwing moths (ash) (dogwood) (lesser peachtree) (Hlac) (oak) (rhododendron) metallic wood (bronze birch) (flatheaded apple tree) (twolined chestnut) longhorned beetles (locust) (red oak) leafminers needleminers (Jeffrey pine) (lodgepole pine) (spruce) scale crawlers (cottonycushion) (cottony maple) (euonymus) (Fletcher) (Florida wax) (golden oak) (hemispherical) (lecanium) (magnolia) (oak kermes) (oystershell) (pine needle) (San Jose) (tea) (white birch)	1 lb (four 4 oz packets)	2 lb (eight 4 oz packets)	1. For borers, apply Dursban 50W to the trunks and lower limbs of trees and shrubs when the adults begin to emerge. Consult your State Agricultural Experiment Station or Extension Service Specialist for proper time to treat. Apply uniformly as a coarse low-pressure spray.
borers (cottonwood) (peachtree)	3 lb (twelve 4 oz packets)	6 lb (twenty-four 4 oz packets)	1. For peachtree borers, apply Dursban 50W in water to flowering trees and shrubs of the genus <i>Prunus</i> as a trunk spray before newly-hatched larvae enter the trees. Apply as a coarse, low-pressure spray. Thoroughly wet all bark areas from ground level to scaffold limbs.
wood infesting beetles ¹ (ambrosia) (Anobiidae) (black turpentine) (cottonwood leaf) (European elm bark) (native elm bark) (willow)	8 lb (thirty-two 4 oz packets)	16 1/2 lb (sixty-six 4 oz packets)	1. For preventive treatment, apply the spray to the main trunk of trees in the early spring or when threat of attack exists from nearby infested trees. For remedial treatment, apply the spray to the main trunk of infested trees when damage occurs but before adult beetles begin to emerge.

† Superscripts refer to "Specific Directions".

Tree Pests (Nurseries, Plantations and Felled Trees)

Use Dursban 50W to treat shade and flowering trees, plantation trees, transplant trees and evergreens found to be infested with pests listed in the following table. Felled trees should be treated as necessary. Dilute Dursban 50W with water according to directions given in the table and apply using suitable hand- or power-operated spray equipment in a manner to provide complete and uniform coverage. For best results, apply a coarse spray to thoroughly wet both upper and lower leaf surfaces and to infested limb and trunk areas. Attempt to penetrate dense foliage, but avoid overspraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals, if needed.

Attention: For felled trees, treat after cutting. Do not handle until spray has dried, or wear protective clothing. Treat only fruit and nut trees identified in the "Tree Fruits and Nuts" section of this label.

Consult your State Agricultural Experimental Station or Extension Service Specialist for application timing and other specific use information.

Pest†	Amount of Dursban 50W in Water to Make:		Specific Directions
	50 gal	100 gal	
adelgids (cooley spruce gall) (eastern spruce gall) (pine bark) aphids (apple) (chrysanthemum) (cottonwood) (elm leaf) (peach) (rose) (spirea) bagworms boxelder bugs cankerworms carpenter ants [†] catalpa sphinx citrus mealybugs clover mite elm sawworms fall webworms [†] fire ants greenstriped mapleworms jackpine budworm juniper webworms katydids lace bugs	1/4 lb (one 4 oz packet)	1/2 lb (two 4 oz packets)	1. For bagworms, treat when insects are in the crawler stage. 2. For effective control of leafrollers, spray should be applied before leaves are tightly rolled. 3. For maple leafcutter on maple trees, apply spray to larvae as cases are being formed. Do not treat sugar maple trees intended for maple syrup production. 4. For effective control of spider mites when large numbers of eggs are present, apply a second spray 3 to 5 days in the south or 7 to 10 days in the north after initial treatment to control newly-hatched nymphs. 5. For effective control of Fall webworm, direct spray into web and immediately surrounding foliage. 6. If possible, locate carpenter ant nests and drench thoroughly.
beetles (native elm bark) leafhoppers mahogany webworms mealybugs mimosa webworms moths (browntail) (Cypress tip) (Douglas fir tussock) (European pine shoot) (gypsy) [†] (holly bud) (Nantucket pine tip) (Pandora) (pitch pine tip) (subtropical pine tip) (tussock)	1/2 lb (two 4 oz packets)	1 lb (four 4 oz packets)	1. To reduce twig and branch feeding on uninfested trees deemed to be of high value, apply a water spray to the tree crown. Application should be made in the spring or early summer using a sprayer that will give thorough coverage to the tree crown. 2. To kill migrating and invading gypsy moth larvae, treat trunks and foliage. 3. Blackvine weevils are night feeders. Late afternoon spraying will give control in some areas.
foliar feeding beetles (blister) (cottonwood leaf) (elm leaf) (flea) (fuller rose) (Japanese) (June) (willow)	1/2 to 1 lb (two to four 4 oz packets)	1 to 2 lb (four to eight 4 oz packets)	1. For cottonwood leaf beetles, use Dursban 50WSP in water to control larvae and adults infesting cottonwoods. Make the treatment when field counts indicate damaging beetle populations are developing or present. For seedlings, use 8 to 20 gallons spray volume per acre.

† Superscripts refer to "Specific Directions"

Pest†	Amount of Dursban 50W in Water to Make:		Specific Directions
	50 gal	100 gal	
borers: clearwing moths (ash) (dogwood) (lesser peachtree) (lilac) (oak) (rhododendron) metallic wood (bronze birch) (flatheaded apple tree) (two-lined chestnut) longhorned beetles (locust) (red oak) Cranberry girdler larvae ² leafminers needleminers (Jeffrey pine) (lodgepole pine) (spruce)	1 lb (four 4 oz packets)	2 lb (eight 4 oz packets)	1 For borers, apply Dursban 50WSP to the trunks and lower limbs of trees when the adults begin to emerge. Consult your State Agricultural Experiment Station or Extension Service Specialist for proper time to treat. Apply uniformly as a coarse low-pressure spray. 2 For cranberry girdler larvae infesting Douglas fir seedlings, apply 2 lb of Dursban 50WSP per acre. Direct spray at the lower crown and stems using 50 gallons of water per acre. Irrigate immediately after application for soil penetration of 1 to 2 inches. Treat after egg laying during the summer.
borers (cottonwood) (peachtree)	3 lb (twelve 4 oz packets)	6 lb (twenty-four 4 oz packets)	1 For peachtree borers, apply Dursban 50WSP in water to flowering trees of the genus <i>Prunus</i> as a trunk spray before newly-hatched larvae enter the trees. Apply as a coarse, low-pressure spray. Thoroughly wet all bark areas from ground level to scaffold limbs.
beetles ¹ (elm leaf) (flea) (fuller rose) (native elm bark) ² (willow leaf)	4 lb (sixteen 4 oz packets)	8 1/3 lb (thirty-three 4 oz packets)	1 For preventive treatment, apply the spray to the main trunk of trees in the early spring or when threat of attack exists from nearby infested trees. For remedial treatment, apply the spray to the main trunk of infested trees or logs when damage occurs but before adult beetles begin to emerge. For plantation trees, apply to individual trees using suitable hand- or power-operated spray equipment. 2 To prevent native elm bark beetles from overwintering in uninfested trees, apply Dursban 50WSP in water to the bottom 9 feet of the trunk. Wet the trunk thoroughly, but do not spray to runoff. Care should be taken to apply the spray right to the base of the root flare. Application can be made with either a backpack mistblower or a hydraulic pressure sprayer from spring through to early fall. To reduce the twig and branch feeding on uninfested trees deemed to be of high value, apply a water spray to the tree crown. Application should be made in the spring or early summer using a sprayer that will give thorough coverage to the tree crown.
beetles ¹ (ambrosia) (Anobiidae) (black turpentine) (European elm bark) (Japanese) (Junc) ¹ (southern pine) (spruce)	8 1/3 lb (thirty-three 4 oz packets)	16 2/3 lb (sixty-seven 4 oz packets)	1 For preventive treatment, apply the spray to the main trunk of trees in the early spring when threat of attack exists from nearby infested trees. For remedial treatment, apply the spray to the main trunk of infested trees or logs when damage occurs or before adult beetles begin to emerge. For plantation trees, apply to individual trees using suitable hand- or power-operated spray equipment.
weevils ¹ (northern pine) (pales) (prich-eating) (twig)	16 1/2 lb (sixty-six 4 oz packets)	33 1/3 lb (133 4 oz packets)	1 For southern pine seedlings, treat immediately after transplanting. Treat each seedling with enough spray to thoroughly wet the foliage and stems to the point of runoff but do not use more than 6 gallons of spray per acre.


† Superscripts refer to "Specific Directions"

Tree Fruits and Nuts

Use Dursban 50W at the dosages indicated by application as a foliar spray to control pests listed in the following table. Mix the required dosage of Dursban 50W in water. Spray for thorough and complete coverage of the foliage, using suitable hand- or conventional power-operated spray equipment. Treat when pests first appear or in accordance with local conditions. Consult your State Agricultural Experiment Station or Extension Service Specialist for specific use information. Do not allow livestock to graze in treated areas.

Attention: In California, do not work with or contact trees within 2 days after treatment.

Crop	Pest	Amount of Dursban 50W in Water to Make		Specific Directions
		50 Gallons	100 Gallons	
Apples	apple aphid apple maggot codling moth European apple sawfly eyespotting bud moth fruittree leafroller green fruitworm lesser appleworm obliquebanded leafroller orange tortrix oriental fruit moth pandemis leafroller plum curculio redbanded leafroller rosy apple aphid San Jose scale tarnished plant bug tufted apple budmoth variegated leafroller woolly apple aphid	1 1/2 lb (six 4 oz packets)	3 lb (twelve 4 oz packets)	Make no more than 8 applications per season. Do not apply last treatment within 28 days before harvest or apply last two treatments closer than 21 days apart.
Almonds	navel orangeworm peach twig borer San Jose scale	2 lb (eight 4 oz packets)	4 lb (sixteen 4 oz packets)	Make no more than 3 applications per season to almonds and filberts and no more than 2 applications per season to walnuts. Do not apply within 14 days before harvest.
Walnuts	codling moth walnut scale	2 lb (eight 4 oz packets)	4 lb (sixteen 4 oz packets)	
Pecans	black pecan aphid hickory shuckworms pecan leaf scorch mite pecan nut casebearer pecan weevil <i>Phylloxera</i> spp.	1 lb (four 4 oz packets)	2 lb (eight 4 oz packets)	Make no more than 5 applications per season. Do not apply within 28 days of harvest. Do not allow livestock to graze in treated orchards.



Warranty Disclaimer

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. DOWELANCO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of DowElanco or the seller. All such risks shall be assumed by Buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at DowElanco's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

DowElanco shall not be liable for losses or damages resulting from handling or use of the product unless DowElanco is promptly notified of such loss or damage in writing. In no case shall DowElanco be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of DowElanco or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Copyright © DowElanco 1992
~~Trademark of DowElanco~~
DowElanco • Indianapolis, IN 46268, U.S.A.

LABEL CODE ~~WG-48-846~~ DATE CODE ~~888~~
EPA APPROVAL ~~684888-448484~~
REPLACES ~~WG-48-888-438-48-846~~

-Revisions:
-4) Added supplemental label for control of mole crickets in subgrasses.

Amendments:

- 1) Front panel statement requiring "... application only by or under the direct supervision of commercial applicators ..." was modified to "... application only by or under the supervision of pest control operators or other trained personnel ..."
- 2) Added use directions for "Pest Control on Outside Surfaces, Around and Under Structures or Crawl Spaces"

Dursban 50WSP Inserts

Insert 1.

To be applied only by or under the supervision of pest control operators or other trained personnel responsible for insect control programs

Insert 2.

Pest Control on Outside Surfaces, Around and Under Structures or in Crawl Spaces

Apply Dursban 50WSP by application as a residual spray to outside surfaces and under structures of buildings including porches, window frames, eaves, crawl space areas, patios, garages, refuse dumps and other areas where pests congregate or have been seen.

Repeat treatment as needed to maintain effectiveness. Avoid application to surfaces where visible residues are objectionable.

Insert 3.

Pest	Amount of Dursban 50WSP in Water to Make 50 Gallons of Diluted Spray:	Specific Directions
	Under Structures or in Crawl Spaces ³ 1/4 to 1 lb (One to four 4 oz packets)	3. Under Structures or in Crawl Spaces: In treating under structures such as crawl spaces, apply spray to foundation areas, cracks and crevices, along sill line, foundation vents, door frames and other areas where pests may find entry or harborage