


(Base label):

**RESTRICTED USE PESTICIDE**  
 For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

(Logo) Dow AgroSciences LLC

# Dursban\* 50W

in Water Soluble Packets



Active Ingredient:

chlorpyrifos: O,O-diethyl O-(3,5,6-trichloro-2-pyridinyl) phosphorothioate .....	50%
Inert Ingredients .....	50%
Total .....	100%

Keep Out of Reach of Children

## DANGER PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

### Precautionary Statements

#### Hazards to Humans and Domestic Animals

Corrosive • Causes Irreversible Eye Damage • May Be Fatal If Swallowed • Harmful If Inhaled Or Absorbed Through Skin

Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing dust and spray mist. Keep away from food, feedstuffs, and water supplies.

#### Personal Protective Equipment (PPE)

**WPS Uses:** Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) -- in general, agricultural-plant uses are covered -- must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear (goggles, face shield, or safety glasses with front, brow, and temple protection)
- Chemical-resistant headgear for overhead exposure

**Non-WPS Uses:** Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) -- in general, only agricultural-plant uses are covered by the WPS -- must wear:

When exposure to spray mist is likely to occur, protective equipment and clothing:

- Long-sleeved shirt and long pants
- Protective eyewear (goggles, face shield, or safety glasses with front, brow, and temple protection)
- Waterproof gloves
- Shoes plus socks
- To prevent breathing of spray mist during application in confined areas, wear a respirator and cartridge(s) approved by MSHA/NIOSH for pesticides (approval number prefix TC-21C).

2/15

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

#### Organophosphate

**If in eyes:** Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

**If swallowed:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

**If on skin:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**If inhaled:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

**Note to physician:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate significance of exposure (baseline data are useful). Atropine, only by injection, is the preferable antidote. Oximes, such as 2-PAM/protopam, may be therapeutic if used early; however, use only in conjunction with atropine. In case of severe acute poisoning, use antidote immediately after establishing an open airway and respiration.

### Environmental Hazards

This pesticide is toxic to birds and wildlife, and extremely toxic to fish and aquatic organisms. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Cover or incorporate spills. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

**Refer to inside of label booklet for additional precautionary information including Personal Protective Equipment (PPE), User Safety Recommendations and Directions for Use including Storage and Disposal.**

**Notice:** Read the entire label. Use only according to label directions. **Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at [www.dowagro.com](http://www.dowagro.com).

**Agricultural Chemical:** Do not ship or store with food, feeds, drugs or clothing.

3/15

EPA Reg. No. 62719-72  
\*Trademark of Dow AgroSciences LLC  
Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

EPA Est. 464-MI-1

## Specialty Insecticide

Net Weight \_\_\_ lb

(Datapack cover):

**RESTRICTED USE PESTICIDE**  
 For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

(Logo) Dow AgroSciences LLC

# Dursban\* 50W in Water Soluble Packets

Active Ingredient:

chlorpyrifos: O,O-diethyl O-	
(3,5,6-trichloro-2-pyridinyl)	
phosphorothioate .....	50%
Inert Ingredients .....	50%
Total .....	100%

**Keep Out of Reach of Children**

## DANGER PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**Agricultural Use Requirements**  
 Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

**Refer to inside of label booklet for additional precautionary information including Personal Protective Equipment (PPE), User Safety Recommendations and Directions for Use including Storage and Disposal.**

**Notice:** Read the entire label. Use only according to label directions. **Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at [www.dowagro.com](http://www.dowagro.com).

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-72  
\*Trademark of Dow AgroSciences LLC  
Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

EPA Est. 464-MI-1

## Specialty Insecticide

Net Weight \_\_ lb

(Page 1 through end):

---

## Precautionary Statements

---

### Hazards to Humans and Domestic Animals

# DANGER

**Corrosive • Causes Irreversible Eye Damage • May Be Fatal If Swallowed • Harmful If Inhaled Or Absorbed Through Skin**

**Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing dust and spray mist. Keep away from food, feedstuffs, and water supplies.**

### Personal Protective Equipment (PPE)

**WPS Uses: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) -- in general, agricultural-plant uses are covered -- must wear:**

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear (goggles, face shield, or safety glasses with front, brow, and temple protection)
- Chemical-resistant headgear for overhead exposure

**Non-WPS Uses: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) -- in general, only agricultural-plant uses are covered by the WPS -- must wear:**

**When exposure to spray mist is likely to occur, protective equipment and clothing:**

- Long-sleeved shirt and long pants
- Protective eyewear (goggles, face shield, or safety glasses with front, brow, and temple protection)
- Waterproof gloves
- Shoes plus socks
- To prevent breathing of spray mist during application in confined areas, wear a respirator and cartridge(s) approved by MSHA/NIOSH for pesticides (approval number prefix TC-21C).

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

#### Organophosphate

**If in eyes:** Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

**If swallowed:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

**If on skin:** Wash with plenty of soap and water. Get medical attention if irritation persists.

6/15

**If inhaled:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

**Note to physician:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate significance of exposure (baseline data are useful). Atropine, only by injection, is the preferable antidote. Oximes, such as 2-PAM/protopam, may be therapeutic if used early; however, use only in conjunction with atropine. In case of severe acute poisoning, use antidote immediately after establishing an open airway and respiration.

### Environmental Hazards

This pesticide is toxic to birds and wildlife, and extremely toxic to fish and aquatic organisms. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Cover or incorporate spills. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

---

### Directions for Use

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

**Do not apply this product through any type of irrigation system.**  
**Do not formulate this product into other end-use products.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

### Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

- Adults, children, and pets should not contact treated surfaces until the spray has dried.

### Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

**Storage:** Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers. Do not store above 122°F for extended periods of time. If container is damaged or spill occurs, use product immediately or dispose of product and damaged container as indicated below. Packets may become brittle when stored below 32°F. Handle carefully to avoid breakage.

**Pesticide Disposal:** Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** When all packets are used, dispose of empty package in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### General Information

Dursban\* 50W In Water Soluble Packets insecticide is a wettable powder formulation designed for use as a spray to control various pests injurious to trees, turf, and ornamental plants. The pests controlled are listed in the accompanying tables. Dursban 50W is compatible with insecticides, miticides, and fungicides commonly recommended except for alkaline materials such as Bordeaux mixture and lime. A small amount of spray mixture should be prepared to check for compatibility before a large volume of spray is mixed.

**Attention:** Keep out of fishpools and other bodies of water. Do not treat vegetable gardens. Do not allow livestock to graze in treated areas. Do not feed treated grass cuttings (hay) or seed screenings to livestock or use hay for livestock bedding. Do not use in poultry houses or greenhouses.

### Handling Precautions for Water Soluble Packets

Do not allow packets to become wet prior to adding to the spray tank. Wear gloves when handling water-soluble packets. Do not handle with wet gloves. Do not handle water-soluble packets excessively since this may cause breakage. Reseal outer package to protect remaining packets. Packets may become brittle when stored below 32°F. Handle carefully to avoid breakage.

### Mixing Directions

For paddle, sparger, and bypass agitation systems.

1. Fill the spray tank 1/3 full of water.  
**Note:** For paddle agitation systems, paddles should be covered with water.
2. Activate agitation system.
3. Add required number of water-soluble packets to obtain desired dilution.
4. Complete tank filling while the packet(s) dissolve and Dursban 50W disperses.
5. Before beginning to spray, make sure water-soluble packet(s) have dissolved and Dursban 50W has completely dispersed.

**Note:** Depending on the water temperature and amount of agitation, the packets and Dursban 50W should be completely dispersed within approximately 5 minutes after adding to water.

## Approved Uses

### **Pest Control on Outside Surfaces and Around Buildings (Industrial Plant Sites Only)**

Apply Dursban 50W by application as a residual spray to outside surfaces, around buildings including window frames, eaves, patios, garages, refuse dumps, and other areas where pests congregate or have been seen. Repeat treatment as needed to maintain effectiveness. Avoid application to surfaces where visible residues are objectionable.

**Note:** Numbers in parentheses following a listed pest or use site refer to Pest-Specific Directions following table.

<b>Pest</b>	<b>Amount of Dursban 50W in Water to Make 50 Gallons of Diluted Spray</b>
Ants	<b>Perimeter Treatment (1)</b> 1/4 to 1 lb (one to four 4 oz packets) <i>Note: One 4 oz packet of            Dursban 50W in 50 gallons            of water equals a            chlorpyrifos concentration of            0.03%</i>
Bees	
Boxelder bugs	
Carpenter ants	
Clover mites	
Cockroaches	
Crickets	
Earwigs	
Elm leaf beetles (adults)	
Firebrats	
Fleas	
Flies	
Hornets	
Millipedes	
Mosquitoes	
Pillbugs	
Scorpions (2)	
Silverfish	
Sowbugs	
Spiders	
Springtails	
Symphylans	
Ticks	
Wasps	
Yellowjackets	
	<b>On Outside Surfaces</b> 4 lb (sixteen 4 oz packets)

#### **Pest-Specific Directions**

- To help prevent infestation of buildings, treat a band of soil 6 to 10 feet wide around and adjacent to buildings, also the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use 4 to 16 ounces (one to four 4 oz packets) of Dursban 50W per 50 gallons of water and apply as a coarse spray at the rate of about 10 gallons spray mixture per 1,000 sq ft to thoroughly and uniformly wet the band area.
- For **scorpions**, remove accumulations of lumber, firewood, and other materials which serve as harborage sites. Dursban 50W may be applied as a residual spray to surfaces immediately below such materials. Perimeter treatments may reduce pest immigration from surrounding areas.
- Apply spray to foundation areas, cracks and crevices, along sill line, foundation vents, door frames and other areas where pests may find entry or harborage. Apply up to 10 gallons per 1,000 sq ft depending on surface texture and type of surface. Use low pressure to minimize splashback. Avoid puddling.



9/15

## Ornamentals Grown in Nurseries

**Agricultural Use Requirements:** When used in nurseries, following reentry interval and PPE requirements for early entry in the "Agricultural Use Requirements" box under "Directions for Use".

Use Dursban 50W insecticide to treat flowers, shrubs, evergreens, vines, shade and flowering trees, and non-bearing fruit trees found to be infested with the types of pests listed in the following tables including: (1) Sucking Insects and Mites; (2) Defoliators and Leafminers; (3) Borers, Bark Beetles and Weevils; and (4) Ants, Termites, and Miscellaneous Pests.

Dilute Dursban 50W insecticide with water according to directions given in the table and apply using suitable hand or power-operated spray equipment in a manner to provide complete and uniform coverage. Attempt to penetrate dense foliage, but avoid over spraying to the point of excessive runoff. For nurseries: when using spray equipment delivering less than 200 gallons of finished spray per acre, use the rate recommended in the per acre column. Uniform coverage is critical for effective insect control.

**Note:** Numbers in parentheses following a listed pest refers to Pest-Specific Directions following each table.

Consult your State Agricultural Experiment Station or Extension Service Specialist for application timing and other specific use information

**Note: Phytotoxicity:** Environmental factors and varietal differences have significant effects on phytotoxic expression. Dursban 50W insecticide has been tested on numerous ornamental plants without causing serious phytotoxicity at recommended use rates. Before treating large numbers of plants, it is recommended that a small block of plants be sprayed and observed for 7 to 10 days to determine phytotoxic potential.

### (1) Sucking Insects and Mites (Ornamentals Grown in Nurseries)

Pest	Amount of Dursban 50W per			
	100 gallons	Acre		
Adelgids (such as: Cooley and Eastern spruce galls, Pine bark)	0.5 to 1 lb (two to four 4 oz packets)	1 to 2 lb (four to eight 4 oz packets)		
Aphids (Such as: Apple, Balsam twig, Black pecan, Chrysanthemum, Cottonwood, Crape myrtle, Elm leaf, Melon, Peach, Rose, Spirea, White pine, Woolly apple, Yellow pecan)				
Boxelder bugs				
Lace bugs (such as: Hawthorn)				
Leafhoppers				
Periodical cicada				
Plant bugs				
Psyllids				
Spittlebugs				
Thornbug				
Whiteflies				
Leafhoppers			1 lb (four 4 oz packets)	2 lb (eight 4 oz packets)
Mealybugs (Such as: Citrus, Taxus)				
Mites (Such as: Clover, Red spider, Southern red) (1) Thrips (exposed)				
Scale insects (Such as: Cottonycushion, Cottony maple, Dearness, Euonymus, Fletcher, Florida wax, Golden oak, Hemispherical, Lecanium, Magnolia, Oak kermes, Oystershell, Pine needle, San Jose, Tea, White birch) (2)	2 lb (eight 4 oz packets)	2 to 4 lb (eight to sixteen 4 oz packets)		

### Pest-Specific Directions

1. For effective control of **spider mites** when large numbers of eggs are present, apply a second spray 3 to 5 days in the South or 7 to 10 days in the North after initial treatment to control newly hatched nymphs.
2. Time applications for control of **scale insects** when crawlers or first two stages of settled scales are present.

**(2) Defoliators and Leafminers (Ornamentals Grown in Nurseries)**

Pest	Amount of Dursban 50W per		
	100 gallons	Acre	
Armyworms (Such as: Fall, Yellowstriped) Bagworms (1) Cankerworms Catalpa sphinx Elm spanworms Fall webworms Grasshoppers Greenstriped mapleworms Green fruitworms Hornworms Jackpine budworms Juniper webworms Katydid Leafrollers (2) Maple leafcutters (3) Oak skeletonizers Oleander caterpillars	Orange tortrix Poplar tentmaker Puss caterpillars Rose chafers Sawflies, exposed (Such as: European pine, Pine, Pin oak, Redheaded) Spring elm caterpillars Springtails Spruce budworms Tent caterpillars (Such as: Eastern, Forest, Western) Walnut caterpillars Western spruce budworms Yellownecked caterpillars	0.5 to 1 lb (two to four 4 oz packets)	1 to 2 lb (four to eight 4 oz packets)
Balsam gall midge Beet armyworms Beetles (Such as: Fuller rose, Native elm bark) (5) Browntail moth Cutworms Cypress tip moth Douglas fir tussock moth European pine shoot moth Gypsy moth (4) Holly bud moth Mahogany webworms	Mimosa webworms Nantucket pine tip moth Oakworms (such as: California, Orangestriped, Redhumped) Pandora moth Pitch pine moth Redhumped caterpillars Subtropical pine tip moth Tussock moth	1 lb (four 4 oz packets)	2 lb (eight 4 oz packets)
Beetles (Such as: Cottonwood leaf, Elm leaf, Flea, Willow leaf) (6)		1 to 2 lb (four to eight 4 oz packets)	2 lb (eight 4 oz packets)
Leafminers Needleminers (Such as: Jeffrey pine, Lodgepole pine, Spruce) Pine needle midge		2 lb (eight 4 oz packets)	2 to 4 lb (eight to sixteen 4 oz packets)

11/15

**Pest-Specific Directions**

1. For **bagworms**, treat when larvae are small and actively feeding.
2. For effective control of **leafrollers**, spray should be applied before leaves are tightly rolled.
3. For **maple leafcutter** on maple trees, apply spray to larvae as cases are being formed. Do not treat sugar maple trees intended for maple syrup production.
4. To control migrating and invading **gypsy moth larvae**, treat trunks and foliage.
5. To reduce twig and branch feeding by bark beetles, applications should be made in the spring or early summer.
6. For cottonwood leaf beetles, use Dursban 50W in water to control larvae and adults infesting cottonwoods. Make the treatment when field counts indicate damaging beetle populations are developing or present.

**(3) Borers, Bark Beetles and Weevils (Ornamentals Grown in Nurseries)**

Pest	Amount of Dursban 50W per	
	100 gallons	Acre
Weevils (such as: Blackvine (1), Cranberry girdler (2), Yellow poplar, Pine reproduction)	1 lb (four 4 oz packets)	2 to 4 lb (eight to sixteen 4 oz packets)
Borers: Clearwing moths (Such as: Ash, Dogwood, Lesser peachtree, Lilac, Oak, Peachtree, Rhododendron), Metallic wood (Such as: Bronze birch, Flatheaded appletree, Twolined chestnut), Longhorned beetles (Such as: Cottonwood, Locust, Red oak) (3) Pales weevil adults Zimmerman pine moth	2 lb (eight 4 oz packets)	-
Pales weevil (4) Northern pine weevil (4)	6 lb (twenty-four 4 oz packets)	-
Beetles (Such as: Ambrosia, Anobiidae, Black turpentine, Blister, Cottonwood leaf, Elm leaf, European elm bark, Fuller rose, Japanese, June, Mountain pine, Native elm bark, Southern pine, Spruce, Western pine, Willow leaf) (5)	16 lb (sixty-four 4 oz packets)	-
Weevils (Such as: Northern pine, Pitch eating, Twig) (6)	32 lb (one hundred twenty-eight 4 oz packets)	-

**Pest-Specific Directions**

1. **Blackvine weevils** are night feeders. Late afternoon spraying will give control in some areas.
2. For **cranberry girdler larvae** infesting Douglas fir seedlings, direct spray at lower crown and stems following egg laying during summer. For nurseries: apply a minimum of 50 gallons of dilute spray per acre and irrigate immediately after application to move the insecticide into the top 1 to 2 inches of soil.
3. For **borers**, spray trunks and lower limbs of trees and shrubs when adults begin to emerge. For **peachtree borers**, spray flowering trees and shrubs of the genus *Prunus* as a trunk spray before newly-hatched larvae enter the trees and thoroughly wet all bark areas from ground level to scaffold limbs. Pheromone traps may aid in detection of adult clearwing moths. Consult your State Agricultural Experiment Station or Extension Service specialist for proper time to treat.
4. For control of **pales and northern pine weevil larvae**, apply as a cut stump spray or drench.
5. For **preventive treatment**, spray the main trunk of trees in the early spring or when threat of attack exists from nearby infested trees. For **remedial treatment**, spray the main trunk of infested trees or logs when damage occurs but before adult beetles begin to emerge.

6. For pine seedlings, treat immediately after transplanting. Treat each seedling with enough spray to thoroughly wet the foliage and stems to the point of runoff. For nurseries: do not use more than 6 gallons of spray dilution per acre.

**(4) Ants, Termites, and Miscellaneous Pests (Ornamentals Grown in Nurseries)**

Pest	Amount of Dursban 50W per	
	100 gallons	Acre
Ants Cockroaches (Such as: American, Asian wood, Brownbanded, German, Oriental, Smokey brown) Fire ants (foraging workers) Fire ants (mounds) (1) Sowbugs Springtails	1 lb (four 4 oz packets)	2 lb (eight 4 oz packets)
Carpenter ants (2) Termites	16 lb (sixty-four 4 oz packets)	-

**Pest-Specific Directions**

1. For individual **fire ant mounds** apply Dursban 50W insecticide as a drench. Dilute 0.75 lb of Dursban 50W (three 4 oz packets) per 50 gallons of water. Gently sprinkle 1 to 2 gallons of the diluted insecticide over the surface of each mound and surrounding areas to a 2 foot diameter. For best results, apply in cool weather, 65-80° F, or in early morning or late evening hours. Treat new mounds as they appear. Pressurized sprays may disturb the ants and cause migration, reducing product effectiveness.
2. If possible, locate carpenter ant nests and drench thoroughly.

**Ornamentals (Pre-Plant Incorporation Treatment of Field Grown Nursery Stock)**

Apply Dursban 50W insecticide to soil and incorporate before transplanting or planting to control white grubs during transplant or seedling establishment. Apply to the soil surface as a broadcast spray at a rate of 8 pounds per acre (thirty-two 4 oz packets) using sufficient water to obtain adequate coverage. Do not aerially apply. On the same day of the treatment, incorporate the insecticide into the top 2 to 4 inches of the soil using a tandem disc, field cultivator, or equivalent incorporation equipment capable of thorough soil mixing.

**Turf in Golf Courses,  
Road Medians and Industrial Plant Sites  
(Maximum Use Rate of 2 lb (1 lb a.i./Acre)**

Labeled turf uses within this section of the label are not within the scope of the WPS. Follow reentry requirements in the "Non-agricultural Use Requirements" box under "Directions for Use".

Use Dursban 50W to control the pests listed in the following table by application at the recommended dosages.

Pest	Amount of Dursban 50W per Acre
Ants Armyworm (such as: Beet, Fall, Yellowstriped) Chiggers (1) Chinch bugs Crickets Cutworms Deer ticks (2) Earwigs European crane fly larvae Fiery skipper Fire ants (foraging workers) Fire ants (mounds) (3) Fleas	Grasshoppers Greenbugs Green June beetle grubs Leafhoppers Lucerne moth Millipedes Mites (Such as: Clover, Bermudagrass stunt, Winter grain) Mosquitoes (4) Pillbugs Sod webworms (lawn moths) (5) Sowbugs Springtails Ticks (1)
	2 lb (eight 4 oz packets)

### General Use Directions

Dilute Dursban 50W insecticide in water and apply using suitable application equipment. For best results, turf should be moist at time of treatment. **Note:** Numbers in parentheses following a listed pest refers to Pest-Specific Directions.

### Pest-Specific Directions

1. Use Dursban 50W insecticide for area control of **ticks** and **chiggers** infesting golf courses, road medians and industrial plant sites where these pests are present and create a nuisance or a possible public health problem. Do not allow public use of treated areas during application or until spray has dried. Apply Dursban 50W insecticide in water at the rate of 0.25 lb per acre (one 4 oz packet) using a hydraulic sprayer, mist applicator, backpack sprayer, or other suitable hand or power operated spray equipment.
2. For control of **deer ticks** apply Dursban 50W insecticide in water at the rate of 2 lb per acre (eight 4 oz packets). Treat low underbrush, turf, grassy areas, weeds, ground surface and debris, using enough spray volume to obtain thorough coverage.
3. For individual **fire ant mounds** apply Dursban 50W insecticide as a drench. Dilute 0.75 lb Dursban 50W (three 4 oz packets) per 50 gallons of water. Gently sprinkle 1 to 2 gallons of the diluted insecticide over the surface of each mound and surrounding areas to a 2 foot diameter. For best results, apply in cool weather, 65-80° F, or in early morning or late evening hours. Treat new mounds as they appear. Pressurized sprays may disturb the ants and cause migration, reducing product effectiveness.
4. **Mosquitoes** coming to rest on areas treated for control of turf pests will be controlled for varying periods of time after treatment depending on exposure of treated areas to weathering conditions.
5. For **sod webworms**, delay watering or mowing of the treated area for 12 to 24 hours after treatment.

### Turfgrass Grown for Sod or Seed (Agricultural Use)

**Agricultural Use Requirements:** When used on turfgrasses grown for seed or sod farms, follow reentry interval and PPE requirements for early entry in the "Agricultural Use Requirements" box under "Directions for Use".

Use Dursban 50W to control the pests listed in the following table by application at the recommended dosages.

Pest	Amount of Dursban 50W per Acre
Ants Armyworm (such as: Beet, Fall, Yellowstriped) Chinch bugs Crickets Cutworms Earwigs European crane fly larvae Fiery skipper Fire ants (foraging workers) Fire ants (mounds) (1) Fleas Grasshoppers Greenbugs Green June beetle grubs Leafhoppers Lucerne moth Millipedes Mites (Such as: Clover, Bermudagrass stunt, Winter grain) Pillbugs Sod webworms (lawn moths) (2) Sowbugs Springtails	2 lb (eight 4 oz packets)
Billbugs (Such as: Bluegrass, Denver, Hunting) (3)	2 to 4 lb (eight to sixteen 4 oz packets)
Black turfgrass ataeenius (4) Annual bluegrass weevil ( <i>Hyperodes</i> ) (5)	4 lb (sixteen 4 oz packets)
Mole Crickets (6)	4 to 6 lb (sixteen to twenty four 4 oz packets)
White grubs (Such as: Black turfgrass ataeenius, European chafer, Southern masked chafer and Japanese beetle larvae) (7)	4 to 8 lb (sixteen to thirty-two 4 oz packets)
Imported fire ants in commercial sod (8)	16 lb (sixty-four 4 oz packets)

### General Use Directions

Dilute Dursban 50W insecticide in water and apply using suitable application equipment. For best results, turf should be moist at time of treatment. **Note:** Numbers in parentheses following a listed pest refers to Pest-Specific Directions.

### Pest-Specific Directions

- For individual **fire ant mounds** apply Dursban 50W insecticide as a drench. Dilute 0.75 lb Dursban 50W (three 4 oz packets) per 50 gallons of water. Gently sprinkle 1 to 2 gallons of the diluted insecticide over the surface of each mound and surrounding areas to a 2 foot diameter. For best results, apply in cool weather, 65-80° F, or in early morning or late evening hours. Treat new mounds as they appear. Pressurized sprays may disturb the ants and cause migration, reducing product effectiveness.
- For **sod webworms**, delay watering or mowing of the treated area for 12 to 24 hours after treatment.
- For **billbugs**, spray early in the season just prior to, or coinciding with first appearance of adults as recommended by your local Agricultural Extension Service Specialist.
- For **black turfgrass ataeenius** adults, spray early in the season as recommended by your local Agricultural Extension Service Specialist. A repeat application may be needed 1 to 2 weeks later.
- To control **annual bluegrass weevil**, spray suspected problem areas in mid-April and again in Mid-May, or as recommended by your local Agricultural Extension Service Specialist.
- To control **mole crickets** in turfgrass, apply Dursban 50W insecticide in a minimum spray volume of 50 gallons of water per acre using broadcast or suitable hand held application equipment. Apply when early stage nymphs are active. Efficacy may be enhanced by spraying late in the afternoon or early evening and irrigating the turf prior to treatment to move mole crickets to the soil surface.

15/15

7. For **white grubs**, spray when grubs are young and actively feeding near the soil surface, usually during late July and August or as recommended by your local Agricultural Extension Service Specialist. For best results, soil should be moist prior to treatment. **For best results, immediately after spraying, irrigate the treated area with 1/2 to 1 inch of water to wash the insecticide into the underlying soil.**
8. Imported **fire ants on sod farms**: Apply Dursban 50W as a spray to the area of sod to be cut. Immediately after application, irrigate treated area with at least 1/2 acre-inch of water or a sufficient volume to thoroughly soak below the cut line. Do not apply a volume or rate of water to cause runoff. Sod can be cut when vegetation has dried and when soil is moist but not wet.  
**Precaution (Additional PPE Requirements):** In addition to PPE listed in this label for WPS uses, elbow-length waterproof gloves, chemical-resistant apron and chemical-resistant footwear plus socks must be worn to prevent repeated skin exposure to treated sod.

**Certification Period:** 42 days from day of application.

---

### Warranty Disclaimer

---

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences **MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

---

### Inherent Risks of Use

---

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

---

### Limitation of Remedies

---

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the "Warranty Disclaimer" above and this "Limitation of Remedies" cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the "Warranty Disclaimer" or this "Limitation of Remedies" in any manner.

\*Trademark of Dow AgroSciences LLC  
EPA Accepted 10/23/00