

(Product Label)

(Logo) DowElanco

Turflon* II Amine

For the Control of Annual and Perennial Broadleaf Weeds in Ornamental Turf

Active Ingredients:

2,4-D: 2,4-Dichlorophenoxyacetic acid, dimethylamine salt	34.2%
Triclopyr: 3,5,6-trichloro-2-pyridinyloxyacetic acid, triethylamine salt	15.2%
Inert Ingredients	50.6%
Total	100.0%

2,4-D acid equivalent 28.4% - 2.78 lb/gal
Triclopyr acid equivalent 10.9% - 1.07 lb/gal

Keep Out of Reach of Children

DANGER

PELIGRO:

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

See back panel for additional precautionary statements.

Notice: Read the entire label. Use only according to label directions. Before buying or using this product, read Warranty Disclaimer and Limitation of Remedies sections elsewhere on this label.

In case of emergency endangering life or property involving this product, call collect 517-636-4400

Agricultural Chemical: Do Not Ship or Store with Food, Feeds, Drugs, or Clothing.


EPA Reg. No. 62719-75

EPA Est. 464-MI-1

*Trademark of DowElanco
DowElanco • Indianapolis, IN 46268, U.S.A.

Herbicide

Net Contents - XX Gallons



Precautionary Statements

Hazards to Humans and Domestic Animals

DANGER

Causes Irreversible Eye Damage • May Be Fatal If Absorbed Through Skin • Harmful If Swallowed or Inhaled

Avoid breathing vapors. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash thoroughly with soap and water after handling and before eating or smoking. Remove contaminated clothing and wash before reuse.

First Aid

If case of skin contact, immediately flush eyes with plenty of water for at least 15 minutes and get medical attention promptly. Flush skin with water and get medical attention if signs of irritation develop. If Swallowed, promptly drink a large quantity of milk, egg whites, gelatin or , if these are not available, drink large quantities of water. Avoid alcohol. Call a physician.

Environmental Hazards

This pesticide is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Directions for Use

It is a violation of Federal Law to use this product in manner inconsistent with its labeling. Read All Directions Carefully Before Applying.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store above 28°F or agitate before use.

Pesticide Disposal: Pesticide spray mixture or rinsate water that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

Refillable Container Disposal: Seal all openings which have been opened during use.

Return the empty container to a collection site designated by DowElanco. If the container has been damaged and cannot be returned according to the recommended procedures, contact DowElanco Customer Service Center at 1-800258-1470 to obtain proper handling instructions.

Non-Refillable Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures. Plastic containers may also be disposed of by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General: Consult federal, state, or local disposal authorities for approved alternative procedures.

General Information

Turfion II Amine is a broad spectrum herbicide recommended for the control of actively growing annual and perennial broadleaf weeds in perennial bluegrass, perennial ryegrass or tall fescue turf.

General Use Precautions (Read Carefully Before Using)

Local conditions may affect the use of herbicides. Consult your local specialist for advice in selecting treatments from this label to best fit local conditions.

Turfion II Amine may injure certain turfgrass species. Do not apply to bahiagrass, bentgrass, bermudagrass, centpedegrass, St. Augustinegrass or zoysiagrass, unless turf injury can be tolerated.

Do not apply Turflon II Amine herbicide directly to or otherwise permit it to come in contact with or permit spray mists containing Turflon II Amine to drift onto cotton, grapes, tobacco, vegetable crops, flowers, fruit or orchard trees, shrubs or other desirable broadleaf plants.

Do not apply Turflon II Amine to exposed roots of shallow rooted trees and shrubs.

For spot treatments, do not apply more than 8 quarts of Turflon II Amine (400 gallons of spray) per acre in a single application.

Avoid Injurious Spray Drift: Applications should be made only when there is little or no hazard from spray drift. Very small quantities of spray, which may not be visible may seriously injure susceptible plants. Do not spray when wind is blowing toward susceptible crops or ornamental plants near enough to be injured. Spray drift can be reduced by spraying when wind velocity is low, keeping the spray boom as low as possible, using nozzles that deliver larger spray droplets, and by using higher volume sprays and lower pressures.

Do not apply to ditches used to transport irrigation water. Do not apply where runoff or irrigation water may flow onto susceptible crops.

Chemigation: Do not apply this product through any type of irrigation system.

Do not use this product for manufacturing or formulating.

Be sure that use of this product conforms to all applicable regulations.

Application Rates and Broadleaf weeds controller

Weeds Controlled		Rate/Acre	Rate/1000 sq ft
Black Medic Carpet weed Catnip Chamise Chickweeds Chicory Cinquefoil Cocklebur Coffee weed Cornflower Creeping beggarweed Dandelion Dayflower Dock Dwarf beggarweed Field bindweed Goldenrod	Henbit Hop clover Knawel Lambsquarter Lespedeza Little starwort Mallow Matchweed Plantain Purslane speedwell Smartweed Sowthistle Spiderwort Spotted catsear Spurweed Vetch Yarrow	2 1/2 pints	1 fl oz or 2 tbsp
Burdock Buttercup Clovers Mustard	Poison ivy Poison oak Wild carrot	3 to 3 1/2 pints	1.3 fl oz or 2 1/2 tbsp
Corn speedwell Ground ivy Oxalis (<i>stricta</i> and <i>corniculata</i>)	Parsley-piert Prostrate spurge Wild violet	3 pints (2 applications 4 to 6 weeks apart)	1.3 fl oz or 2 1/2 tbsp (2 applications 4 to 6 weeks apart)

Mixing Instructions

Add about one-half the required amount of clean water to the spray tank. Start agitation and add the recommended amount of Turflon II Amine. Provide moderate agitation while completing the addition of water and during application.

Application Directions

Foliar sprays should be applied during warm weather, from early spring through fall, when weeds are actively growing. Application under drought conditions may provide less than desirable results. Use low pressure sprays to minimize spray drift. When making applications to mature plants, hard to control species, or during drought conditions, use higher rates.

Note: Broadleaf weeds germinate at different times. Only emerged weeds present at the time of application, will be controlled. Newly seeded turf should be mowed two or three times before being treated. Do not water for 24 hours after application. Do not reseed for 3 weeks after application.

Broadcast Treatment of Ornamental Turf

Apply 2 1/2 to 3 1/2 pints of Turflon II Amine in 20 to 200 gallons of water per acre to control actively growing broadleaf weeds growing in perennial bluegrass, perennial ryegrass, or tall fescue. Do not use on other turfgrass species (see General Use Precautions) unless injury can be tolerated. To minimize turf injury, do not treat if turf is under heat or drought stress and make repeat applications at least four weeks apart.

Spot Treatment of Ornamental Turf

Mix 1 to 2 fluid ounces of Turflon II Amine in enough water to make three gallons of spray (1 to 2 quarts per 100 gallons of spray) and apply at any time broadleaf weeds are susceptible by wetting foliage of susceptible plants to point of runoff. **Note:** Do not apply more than 8 quarts of Turflon II Amine (400 gallons of spray) per acre in a single application.

WARRANTY DISCLAIMER

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. DOWELANCO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of DowElanco or the seller. All such risks shall be assumed by Buyer.

LIMITATION OF REMEDIES

The exclusive remedy, for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at DowElanco's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

DowElanco shall not be liable for losses or damages resulting from handling or use of this product unless DowElanco is promptly notified of such loss or damage in writing. In no case shall DowElanco be liable for consequential or incidental damages or losses. The terms of the Warranty

Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of DowElanco of the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

rsb 1/30/92

Amendments

- 1) Label edited to reflect DowElanco transfer
- 2) Revised instructions to prevent spray drift.