

Supplemental Labeling



DowElanco

Quad IV, 9002 Purdue Road

P O Box 681428

Indianapolis, Indiana 46268-1189 USA

Treflan® E.C. (EPA Reg. No. 2719-93)
~~Treflan M.T.F. (EPA Reg. No. 62719-116)~~
~~Treflan S (EPA Reg. No. 62719-118)~~
~~Treflan 800G (EPA Reg. No. 62719-216)~~

Treflan for Weed Control in Soybeans Grown Under Reduced or Conservation Tillage Conditions

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow use precautions and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all precautions and limitations imposed by the label affixed to the container for Treflan.

Directions For Use

Treflan can be applied either in the fall or in the spring as a preplant incorporated treatment for weed control in soybeans grown under reduced or conservation tillage conditions. Make only one application per crop cycle.


Make applications to tilled land or standing or chopped stubble from the previous season's crop. Use incorporation equipment that thoroughly and uniformly mixes Treflan into the top 2-3 inches of the final seedbed and leaves a maximum amount of crop residue on the soil surface. Incorporation must occur within 24 hours after application. For fall applications, incorporate once in the fall and a second time in the spring before planting. For spring applications, incorporate twice for best results.

Application With Dry Bulk Fertilizers

Dry bulk fertilizers impregnated or coated with Treflan may be applied as preplant incorporated treatments. See instructions for "Application with Dry Bulk Fertilizer" in the "General Information" section of the Treflan label. Under reduced or conservation tillage conditions, uniformly applied dry bulk fertilizers impregnated with Treflan provide weed and grass control equal to or better than Treflan applied in liquid sprays. Two incorporation passes are required when Treflan is applied with dry bulk fertilizer. For best results with spring applications, incorporate once within 24 hours after application and a second time 3 to 5 days later.

Application Rates Per Acre (Fall Applications)

Soil Texture	Treflan E.C. or Treflan M.T.F. (pints)	Treflan S (pints)	Treflan 800G (ounces)
Coarse	1.5 - 2.0	1.2 - 1.6	15 - 20
Medium	2.0 - 2.5	1.6 - 2.0	20 - 25
Fine	2.5 - 3.0	2.0 - 2.4	25 - 30



BEST AVAILABLE COPY

continued on back

Application Rates Per Acre (Spring Applications)

Soil Texture	Treflan E.C. or Treflan M.T.F.	Treflan 5	Treflan 100G
	(pints)	(pints)	(ounces)
Coarse	1.0 - 1.5	10.8 - 1.2	10 - 15
Medium	1.5 - 2.0	1.2 - 1.6	15 - 20
Fine	2.0 - 2.5	1.6 - 2.0	20 - 25

Use the higher rate in the rate range where higher crop residues are present or where dense weed populations are anticipated.

Precautions

To be effective, Treflan must be mixed thoroughly in the top 2-3 inches of soil in the final seedbed. Weed control may be poor or erratic where soil conditions or heavy crop residues do not permit thorough soil mixing.

*Trademark of DowElanco

(Label Code) Approved / /
Initial printing.

Amendment(s):
1) Added use on soybeans under
reduced/conservation tillage conditions.