



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
62719-106

Date of Issuance:
NOV 30 1999

NOTICE OF PESTICIDE:
Registration
 Reregistration

Term of Issuance:

Name of Pesticide Product:
Surflan* 75WSP

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Dow AgroSciences LLC
9330 Zionville Road
Indianapolis, IN 46268

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Based on your response to the Reregistration Eligibility Document, EPA has reregistered the product listed above. The following revisions must be incorporated into the labeling before the product is released for shipment:

1. Under the headings entitled "Personal Protective Equipment" on page (5) five of the product label, remove the reference to "protective eyewear".

Signature of Approving Official:

Date:

NOV 30 1999

M. HOWARD

page 2

EPA Reg. No. 62719-106

Enclosed is a copy of your stamped "Accepted with Comments". This action is taken under the authority of section 4(g)(2)(C) of the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended. Reregistration under this section does not eliminate the need for data at any time to maintain registration of your product.

Joanne I. Miller
Product Manger (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

(Base Label)

NOV 30 1999

(logo) Dow AgroSciences LLC

Surflan* 75WSP

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
62719-106

A preemergence surface-applied herbicide for control of most annual grasses and certain broadleaf weeds.

Active Ingredient:

oryzalin: 3,5-dinitro- N ⁴ , N ⁴ -dipropylsulfanilamide	75%
Inert Ingredients	25%
Total	100%

Contains 1.0 pound active ingredient per 1.3 pound water soluble packet.

This resealable package contains Surflan 75WSP herbicide in water soluble packets. Do not remove water soluble packet except for immediate use.

Keep Out of Reach of Children

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Moderate Eye Irritation • Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Avoid contact with skin, eyes, or clothing.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 15 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Handling Precautions for Water Soluble Packets

Do not remove packet from foil overpack except for immediate use. Do not allow packet to come in contact with water prior to use. Do not handle with wet hands. Carefully reseal package after use and protect unopened packets from moisture.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. **Exception:** If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Workers may enter treated areas without required PPE during the reentry interval following 1/2 to 1 inch of rainfall or irrigation, otherwise, PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves
- Shoes plus socks

STORAGE AND DISPOSAL

Storage: Store in original container only. In case of leak or spill, contain material with absorbent material and dispose as waste.

Pesticide Disposal: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Dispose of overpack in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General Information

Surflan 75WSP herbicide is a preemergence, surface applied herbicide for the control of most annual grasses and certain broadleaf weeds. Surflan 75WSP controls susceptible annual weeds by disrupting plant growth processes during germination. Surflan 75WSP may be applied in liquid sprays of water or liquid fertilizer (See special mixing instruction if used with liquid fertilizer). Surflan 75WSP may be tank mixed with other herbicides to control existing vegetation or improve the spectrum of weeds controlled. Surflan 75WSP alone does not control established weeds.

General Use Precautions and Restrictions

Do not graze or feed forage from treated fields or orchards to livestock.

Poor weed control may result if directions are not carefully followed.

Do not over-apply Surflan 75WSP. Over-application may result in crop injury or soil residue.

Do not plant any root crop for 12 months following a Surflan 75WSP application.

Do not use Surflan 75WSP on soils containing more than 5% organic matter. Apply Surflan 75WSP directly to the soil surface in orchards or vineyards.

Carefully follow label directions to avoid poor weed control or crop injury.

Chemigation: See instructions for chemigation in "Application Methods".

Do not aerially apply this product.

Soil Preparation

Optimum herbicidal activity occurs when Surflan 75WSP is applied to the soil surface. Prior to application, existing unwanted vegetation should be destroyed by tillage or use of contact or translocated herbicides. Crop or weed residues, prunings and trash should be thoroughly mixed into the soil by tillage equipment or removed before treatment. Soil should be in good tilth and free of clods at the time of application.

Activation and Cultivation

A single 1/2 to 1 inch rainfall or sprinkler irrigation is required to activate Surflan 75WSP and move the herbicide into the zone of weed germination. Rainfall or irrigation of 1 inch or more is needed to activate Surflan 75WSP on fine-textured, high organic matter soils. If weeds begin to emerge, a shallow cultivation to a depth of 1 to 2 inches will destroy existing weeds and place Surflan 75WSP in the zone of weed seed germination.

Mixing Directions

Surflan 75WSP - Alone

Surflan 75WSP may be applied in water or most liquid fertilizer materials. Prior to mixing Surflan 75WSP in liquid fertilizer, refer to "Testing for Compatibility in Liquid Fertilizers" for test procedures to determine compatibility with the fertilizer product to be used. The combination of Surflan 75WSP with solution and suspension-type fertilizers provides annual weed control equal to Surflan 75WSP applied in water. Individual state regulations relating to liquid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company offering the fertilizer and chemical mixture for sale.

Start with a clean spray tank. Fill the sprayer to 1/3 to 1/2 of the required spray volume. Start agitation. Add the required number of water soluble packets to the spray tank. Allow time for complete mixing of Surflan 75WSP after packets have disintegrated. Continue agitation and fill spray tank to required spray volume. Maintain continuous agitation from mixing through application.

Special Instruction: Do not add Surflan 75WSP water soluble packets directly to liquid fertilizers. Surflan 75WSP in water soluble packets should first be premixed with water (See instruction for premixing or slurring below). Add premixed Surflan 75WSP to the spray tank through a 20-35 mesh wetting screen to insure good initial dispersion.

Precaution: Do not allow the mixture to siphon back into the water source.

Surflan 75WSP - Tank Mix Combinations

Surflan 75WSP may be tank mixed with other products and applied with water or most liquid fertilizer materials. Prior to tank mixing Surflan 75WSP with other herbicides, mix the products together in correct proportions in a small container of water. After mixing, let stand for 5 minutes and check for compatibility. If the combination remains mixed or can be resuspended easily, the mixture is compatible.

Tank Mixing in Liquid Fertilizer: Prior to mixing Surflan 75WSP with other products in liquid fertilizer, refer to the tank mix product manufacturer's label to determine if application in liquid fertilizer is recommended. Also refer to "Testing for Compatibility in Liquid Fertilizers" for testing procedures to determine compatibility with the fertilizer product to be used. The combination of Surflan 75WSP with solution and suspension-type fertilizers provides annual weed control equal to Surflan 75WSP applied in water. Individual state regulations relating to liquid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company offering the fertilizer and chemical mixture for sale. Read and follow all label instructions for each material to be added to the spray tank.

Vigorous continuous agitation is required for all tank mixes. Sparger pipe agitators generally provide the best agitation in spray tanks. To prevent foaming, keep the end of the fill pipe below the surface of the water in the spray tank during filling to prevent air from being stirred or splashed into the mixture.

Mixing Order: Fill the spray tank to 3/4 of the total spray volume required. Start agitation. Add different formulation types in the order indicated below, allowing time for complete mixing and dispersion after

addition of each product. Allow extra mixing and dispersion time for dry flowable products. Allow time for complete mixing after Surflan 75WSP water soluble packets have disintegrated.

Add different formulation types in the following order: dry flowables (DF); Surflan 75WSP and or other wettable powders (WP); flowables (F), liquids (L) or aqueous suspensions (AS); solutions (S); and emulsifiable concentrates (EC). Fill tank to final spray volume. Maintain continuous agitation during mixing, final filling and throughout application. If spraying and agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be resuspended before spraying is resumed. A sparger agitator is particularly useful for this purpose. Settled materials may be more difficult to resuspend than when originally mixed.

Premixing: When tank mixing or mixing with liquid fertilizers, initial mixing and dispersion of dry flowable or wettable powder products may be improved by premixing with water (slurrying). Surflan 75WSP in water soluble packets should be slurried prior to mixing with liquid fertilizer materials. Where recommended for other products, follow product label instructions for each material. Adding the slurried material to the spray tank through a 20 to 35 mesh wetting screen will help assure good initial dispersion. Line screens in the tank should be no finer than 50 mesh (100 mesh is finer than 50 mesh).

Testing for Compatibility in Liquid Fertilizers

Surflan 75WSP alone or in combination with dry flowable (DF), wettable powder (WP), aqueous suspension (AS), flowable (F), liquid (L), solution (S) or emulsifiable concentrate (EC) formulations may not combine properly with some liquid fertilizer materials. Small quantities of such mixtures should always be tested before full-scale mixing. Follow the testing procedure below to determine if a compatibility agent is needed or which compatibility agent works best in your liquid fertilizer plus herbicide mixture. Nine compatibility agents listed at the end of this section have been thoroughly tested. Other surfactant materials available may not be formulated for use with liquid fertilizers.

Testing Procedure

1. Add 1 pint of liquid fertilizer to 1 quart glass jar.
2. Add 1 to 4 teaspoonfuls of DF, Surflan 75WSP, other WP formulations, AS, F, or L formulations, depending on mixing ratio required, to the liquid fertilizer. Close the jar and shake until evenly dispersed after addition of each formulation. If dry flowable or wettable powder formulations do not disperse well, it may be necessary to slurry the materials in a small amount of water before addition to the liquid fertilizer.
3. After dispersing the materials in step 2, add any S formulations to the jar and shake well. Finally, add EC formulations to the mixture and shake well. Observe the jar for about 10 minutes. If materials rise to the surface and form a thick layer that will not redisperse when agitated, a compatibility agent is needed. If the mixture is easily redispersed with slight agitation, a compatibility agent is not required. Good agitation, however, must be provided to maintain dispersion in the spray tank from mixing through application.
4. If the need for a compatibility agent is demonstrated in step 3, the following procedure is recommended: Using a clean clear plastic or glass container, repeat step 1 above and add 1/2 teaspoon of the compatibility agent to the liquid fertilizer mixture. Shake well and then repeat steps 2 and 3.

An effective compatibility agent will cause the mixture to remain uniformly mixed with little or no separation for 1/2 hour or longer. If slight separation occurs, 2 to 3 inversions of container should be sufficient to uniformly redisperse the mixture. If layers form which will not disperse, try adding additional compatibility agent or use an alternative compatibility agent to achieve a uniform mixture.

Use a clean jar in each test. A compatible mixture will have a uniform appearance and will be relatively easy to redisperse with gentle agitation of the jar.

Compatibility Agents

The phosphate ester-type surfactants listed below are designed for use with liquid fertilizers and can be mixed at rates as low as 1 1/2 to 2 pints per ton of liquid fertilizer. They usually do not work well as compatibility agents for tank mixes in plain water. Add the compatibility agent just before adding herbicides. Read and follow label directions for the compatibility agent.

1. Sponto 168D (Witco Chemicals Co., Chicago, IL)
2. Compat (Farm Chemicals, Inc., Aberdeen, NC)
Not for use in California.
3. Unite (Hopkins Ag Chemical, Madison, WI)
4. T-Mulz 734-2 (Thompson-Hayward Chemical Co., Kansas City, KS)

- Not for use in California.
5. Rigo Compatibility Agent (Rigo Company, Buckner, KY)
Not for use in California.
 6. Amoco Spray Mate™ (Amoco Oil Co., Chicago, IL)
Not for use in California
 7. Kem-Link (Universal Coop, Minneapolis, MN)
Not for use in California
 8. Blendex™ (Helena Chemical Co., Cayce - West, Columbia, SC)
Not for use in California.
 9. Spray-Aide™ (Miller Chemical and Fertilizer Corp., Hanover, PA)

Application Methods

Ground Broadcast Application

Apply Surflan 75WSP in a total spray volume of 20 to 40 gallons per acre (broadcast basis), using any properly calibrated low pressure herbicide sprayer that will apply the spray uniformly. Use herbicide nozzle tips and screens no finer than 50 mesh for nozzle and in-line strainers. As the amount of spray volume per acre decreases, the importance of accurate calibration and uniform application increases. Check the sprayer daily to insure proper calibration and uniform application. Avoid boom overlaps that will increase rates above those recommended.

Band Application

For band application, use the following formula to calculate the required amount of Surflan 75WSP per acre.

$$\frac{\text{Band Width (inches)}}{\text{Row Width (inches)}} \times \text{Broadcast rate per acre} = \text{Amount required per acre}$$

Chemigation

Surflan 75WSP may be applied through properly equipped chemigation systems for weed control in fruit and nut orchards or vineyards. Read and follow all label instructions outlined below concerning chemigation before applying Surflan 75WSP by this method.

Chemigation Use Precautions: Make **non-dormant** chemigation applications of Surflan 75WSP only through solid set or hand move systems designed to distribute sprinkler irrigation beneath the tree canopy. Solid set systems utilizing tall risers for overhead application may be used only for **dormant** season chemigation applications. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for chemigation application to a public water system.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Sprinkler Chemigation Directions

The following directions must be followed for all recommended sprinkler irrigation systems (solid set and hand move systems):


1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain, appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, spring-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

13 71

Brachiaria (Signalgrass)	<i>Brachiaria</i> spp.
Browntop panicum	<i>Panicum fasciculatum</i>
Crabgrass (Large crabgrass) (Smooth crabgrass)	<i>Digitaria</i> spp.
Crowfootgrass	<i>Dactyloctenium aegyptium</i>
Cupgrass	<i>Eriochloa gracilis</i>
Downy brome	<i>Bromus tectorum</i>
Fall panicum (Spreading panicgrass)	<i>Panicum dichotomiflorum</i>
Foxtails	<i>Setaria</i> spp.
(Bottlegrass)	
(Bristlegrass)	
(Giant foxtail)	
(Green foxtail)	
(Pigeongrass)	
(Robust foxtail)	
(Yellow foxtail)	
Guineagrass (Narrowleaf panicum)	<i>Panicum maximum</i>
Johnsongrass (Seedling only)	<i>Sorghum halepense</i>
Junglerice	<i>Echinochloa colonum</i>
Lovegrass, Mexican	<i>Eragrostis mexicana</i>
Lovegrass, orcutt	<i>Eragrostis orcuttiana</i>
Oat, wild	<i>Avena fatua</i>
Ryegrass, annual (Italian)	<i>Lolium multiflorum</i>
Sandbur, field	<i>Cenchrus incertus</i>
Sprangletop, red	<i>Leptochloa filiformis</i>
Texas panicum (Buffalograss) (Coloradograss)	<i>Panicum texanum</i>
Witchgrass	<i>Panicum capillare</i>

Annual Broadleaf Weeds

Common Name	Scientific Name
Carpetweed	<i>Mollugo verticillata</i>
Chickweed	<i>Stellaria media</i>
Cudweed	<i>Gnaphalium chilense</i>
Fiddleneck, coast	<i>Amsinckia intermedia</i>
Florida pusley (Florida purslane) (Mexican clover) (Pusley)	<i>Richardia scabra</i>
Henbit	<i>Lamium amplexicaule</i>
Knotweed, prostrate	<i>Polygonum aviculare</i>
Lambsquarters	<i>Chenopodium album</i>
Pigweeds	<i>Amaranthus</i> spp.
(Carelessweed)	
(Prostrate pigweed)	
(Redroot pigweed)	
(Rough pigweed)	
(Smooth pigweed)	
(Spiny pigweed)	
(Tumble pigweed)	
Puncturevine	<i>Tribulus terrestris</i>
Purslane, common	<i>Portulaca oleracea</i>
Rockpurslane, redmaids	<i>Calandrinia caulescens</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>



14 2 1

Surflan 75WSP provides partial control or suppression of:

Common Name	Scientific Name
Filaree, redstem	<i>Erodium cicutarium</i>
Filaree, whitestem	<i>Erodium moschatum</i>
Groundsel, common	<i>Senecio vulgaris</i>
Ladysthumb	<i>Polygonum persicaria</i>
Lettuce, prickly	<i>Lactuca serriola</i>
Mallow, common	<i>Malva neglecta</i>
Milkweed, climbing	<i>Sarcostemma cynanchoides</i>
Morningglory, annual	<i>Ipomoea spp.</i>
Mustard, black	<i>Brassica nigra</i>
Mustard, wild	<i>Sinapis arvensis</i>
Nightshade, black	<i>Solanum nigrum</i>
Prickly sida (teaweed)	<i>Sida spinosa</i>
Ragweed, common	<i>Ambrosia artemisiifolia</i>
Ragweed, giant	<i>Ambrosia trifida</i>
Rocket, London	<i>Sisymbrium irio</i>
Smartweed, annual	<i>Polygonum spp.</i>
Sowthistle, annual	<i>Sonchus oleraceus</i>
Spurge, prostrate	<i>Euphorbia humistrata</i>
Spurge, spotted	<i>Euphorbia maculata</i>
Velvetleaf	<i>Abutilon theophrasti</i>

Approved Crops

**Fruit and Nut Crops and Vineyards
(Non-bearing and Bearing)**

Surflan 75WSP - Alone

Apply Surflan 75WSP to fruit and nut crops and vineyards as a preemergence treatment to control annual grasses and broadleaf weeds listed in the "General Information" section of this label.

Surflan 75WSP is approved for use on the following crops:

Almond	Grapefruit	Pear
Apple	Kiwi	Pecan
Apricot	Lemon	Pistachio
Avocado	Macadamia Nut	Plum
Cherry	Nectarine	Pomegranate
Fig	Olive	Prune
Filbert Orange	Orange	Walnut, English
Grape	Peach	

Small Fruits:

Blackberry	Currant	Gooseberry
Blueberry ^{††}	Dewberry	Loganberry
Boysenberry	Elderberry	Raspberry

^{††}Do not apply Surflan 75WSP to lowbush blueberries.

Broadcast Application Rates per acre

Soil Texture	Length of Control	Surflan 75WSP (lb/acre)	Minimum Time Between Applications (months)	Total Amount Allowed Per Year (lb/acre)
All Soil Textures	Short Term (2 - 4 months)	2.6	2.5	16
	Long Term (6 - 8 months)	5.2	2.5	16
	(8 - 12 months)	7.8	2.5	16

Chemigation Application of Surflan 75WSP

Apply Surflan 75WSP in 1/2 to 1 inch of sprinkler irrigation. Apply in at least 1 inch of sprinkler irrigation on medium to fine textured or high organic matter soils. Surflan 75WSP does not control established weeds. Surflan 75WSP must be applied prior to weed germination or immediately after existing weeds are controlled. Control existing unwanted vegetation with tillage or with a contact or translocated herbicide.

Cultivation

If weeds begin to emerge after application from below the zone of herbicidal activity, a shallow cultivation of 1 to 2 inches will destroy existing weeds and mix Surflan 75WSP into the zone of weed germination.

Chemigation Use Precautions

Overtop applications with sprinkler systems utilizing tall risers may be made only to **dormant** tree fruit and nut orchards or vineyards. Make chemigation applications to **non-dormant** tree fruit and nut orchards only through solid set or hand move sprinkler systems with risers and sprinkler heads designed to distribute water uniformly beneath the tree canopy.

Surflan 75WSP - Tank Mix Recommendations

The products listed below are compatible with Surflan 75WSP when used in tank mix combination. Performance and risk of carryover from products listed below when used in combination with Surflan 75WSP at recommended rates is the same as when each product is used separately.

Follow tank mixing instructions provided in the "General Information" section of this label when mixing Surflan 75WSP with other products.

Users should always consult the manufacturers label of product(s) to be tank mixed with Surflan 75WSP for specific information on use rates, additional weeds controlled, rotational crop restrictions or risk of carryover, special tank mix instructions, additional use directions, precautions and limitations before use. **Note:** Some crop uses of tank mix products are limited to or excluded from certain states. Such limitations are specified in the label of the product to be tank mixed with Surflan 75WSP

Recommended Surflan 75WSP Tank Mix Combinations:

Surflan 75WSP plus Goal
 Surflan 75WSP plus Goal plus Gramoxone Super*
 Surflan 75WSP plus Gramoxone Super
 Surflan 75WSP plus Princep (Simazine)
 Surflan 75WSP plus Roundup
 Surflan 75WSP plus Solicam

*Use of an agriculturally registered non-ionic surfactant is recommended with this 3-way tank mix combination to improve the compatibility of these herbicides when mixed together. Fill the spray tank one-third to one-half (1/3 to 1/2) full with clean water. With agitation running, add either one (1) quart of X-77 or two (2) quarts of Triton AG-98 per 100 gallons of spray mixture. Add the recommended amount of Surflan and mix thoroughly. Continue filling the spray tank and add Goal. Then add Gramoxone Super. If a nonionic spreader is being used, add it to the tank last. Apply in at least 50 gallons of water per acre.

Surflan 75WSP - Tank Mix Reference Table:

The following table lists approved crops for Surflan 75WSP and products for tank mixes with Surflan 75WSP

(* = Approved Crop). Refer to the table to determine if the herbicides recommended for tank mixing can be applied to your crop. For use on a given crop, two asterisks must be indicated for Surflan 75WSP and the product with which it is to be tank mixed.

	Surflan 75WSP	Goal	Gramoxone Super	Princep (Simazine)	Roundup	Solicam
Almond	**	**	**	**	**	**
Apple	**	**	**	**	**	**
Apricot	**	**	**	**	**	**
Avocado	**	**	**	**	**	**
Cherry	**	**	**	**	**	**
Fig	**	**	**	**	**	**
Filbert	**	**	**	**	**	**
Grape	**	**	**	**	**	**

16 2 1

Grapefruit	"	-	"	"	"	"
Kiwi	"	"	"	"	"	"
Lemon	"	-	"	"	"	"
Macadamia nut	"	"	"	"	"	"
Nectarine	"	"	"	"	"	"
Olive	"	"	"	"	"	"
Orange	"	-	"	"	"	"
Peach	"	"	"	"	"	"
Pear	"	"	"	"	"	"
Pecan	"	"	"	"	"	"
Pistachio	"	"	"	"	"	"
Plum	"	"	"	"	"	"
Pomegranate	"	"	"	"	"	"
Prune	"	"	"	"	"	"
Walnut, English	"	"	"	"	"	"

Small Fruits:

Blackberry	"	-	"	"	"	"
Blueberry	"	-	"	"	"	"
Boysenberry	"	-	"	"	"	"
Currants	"	-	"	"	"	"
Dewberry	"	-	"	"	"	"
Elderberry	"	-	"	"	"	"
Gooseberry	"	-	"	"	"	"
Loganberry	"	-	"	"	"	"
Raspberry	"	-	"	"	"	"

Christmas Tree Plantations**Surflan 75WSP Alone**

Apply Surflan 75WSP as a directed spray to the soil surface or as an otop spray to established plantings of field grown Christmas tree species, including fir (*Abies spp.*), pine (*Pinus spp.*), and spruce (*Picea spp.*). Do not apply to Douglas-fir (*Pseudotsuga menziesii*). Do not apply to seedbeds or seedling transplant beds. Apply only to established plantings. Established plants are defined as those that have been transplanted into their final growing location for a sufficient period of time to allow the soil to be firmly settled around the roots from packing and rainfall or irrigation. Follow all instructions provided in the "General Information" section of this label.

Broadcast Application Rates

Length of Control	Surflan 75 WSP	Minimum Time Between Applications (months)	Total Amount Allowed Per Year (lb/acre)
	(lb/acre)		
Short term (2 - 4 months)	2.6	2	10.7
Long term (4 - 8 months)	5.2	2	10.7

Tank-Mix Combinations

Tank-mix combinations of Surflan 75WSP plus other labeled herbicides may be used as directed or otop sprays in established Christmas tree plantings. When applied according to use directions, these tank-mixes will provide control of susceptible weed species listed on the respective product labels. Refer to label of the product to be tank mixed with Surflan 75WSP for specific use directions, precautions and limitations before use.

Surflan 75WSP/Roundup: Apply tank-mix combinations of Surflan 75WSP plus Roundup only as directed sprays in Christmas tree plantings. When applied according to use directions, Surflan 75WSP plus Roundup will provide postemergence control of susceptible weed species listed on the Roundup label and residual preemergence control of susceptible weed species listed on the Surflan 75WSP label. Refer to the Roundup label for specific use directions, precautions and limitations before use.

Precautions:

Do not apply sprays containing Roundup over the top of Christmas tree plantings.

