

(Base Label which remains on container after booklet is removed.)

(logo) DowElanco

Surflan* A.S.


A selective preemergence surface-applied herbicide for control of annual grasses and many broadleaf weeds in:

- Landscape Ornamentals
- Container Grown Ornamentals
- Field Grown Ornamentals
- Drainage Areas Under Greenhouse Shadehouse Benches
- Ornamental Bulbs
- Ground Covers / Perennials
- Christmas Tree Plantations
- Noncropland and Industrial Sites - Non-bearing fruit and nut trees and non-bearing vineyards
- Industrial Sites
- Established Warm Season Turf (including Bahiagrass, Bermudagrass, Buffalograss, Centipedegrass, St. Augustinegrass, Tall Fescue and Zoysiagrass)
- Tall Fescue (warm season areas)

Active Ingredient:

oryzalin: 3,5-dinitro-N,N'-dipropylsulfanilamide	40.4%
Inert Ingredients	59.6%
Total	100.0%

Contains 4.0 pounds of active ingredient per gallon.



Shake Well Before Using

Precautionary Statements

Hazards to Humans and Domestic Animals

Keep Out Of Reach Of Children

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Causes Eye Irritation • Harmful If Swallowed, Inhaled Or Absorbed Through The Skin

Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Surflan A.S. may cause skin sensitization reactions in certain individuals.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid**If in eyes:** Flush with plenty of water. Get medical attention if irritation develops.**If on skin:** Wash with plenty of soap and water. Get medical attention if irritation develops.**If swallowed:** Do not induce vomiting. Call a physician or Poison Control Center. If available, administer activated charcoal (6-8 heaping teaspoonfuls) with a large quantity of water. Do not give anything by mouth to an unconscious person. Immediately transport to a medical care facility and see a physician.**If inhaled:** Remove individual to fresh air. Get medical attention if breathing difficulty occurs. If not breathing, give artificial respiration, preferably cardiopulmonary resuscitation and get medical attention immediately.**Environmental Hazards**

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Cover or incorporate spills.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read All Directions for Use carefully before applying.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.**Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.**Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to label booklet for additional precautionary information including Personal Protective Equipment (PPE) and User Safety Recommendations, and Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.

In case of emergency endangering health or the environment involving this product, call collect 517-636-4400.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-113

EPA Est. 37429-GA-01

page 3

printed April 12, 1995

*Trademark of DowElanco
DowElanco • Indianapolis, IN 46268, U.S.A.

Specialty Herbicide

Net Contents XX gal

[Laminated Booklet - Surflan AS (Specialty)]
[Front Cover of Booklet]

(logo) DowElanco

Surflan* A.S.

A selective preemergence surface-applied herbicide for control of annual grasses and many broadleaf weeds in:

- Landscape Ornamentals
- Container Grown Ornamentals
- Field Grown Ornamentals
- Drainage Areas Under Greenhouse Shadehouse Benches
- Ornamental Bulbs
- Ground Covers / Perennials
- Christmas Tree Plantations
- Noncropland and Industrial Sites
- Non-bearing fruit and nut trees and non-bearing vineyards
- Industrial Sites
- Established Warm Season Turf (Including Bahiagrass, Bermudagrass, Buffalograss, Centipedegrass, St. Augustinegrass, Tall Fescue and Zoysiagrass)
- Tall Fescue (warm season areas)

Active Ingredient:

oryzalin: 3,5-dinitro- <i>N,N</i> -dipropylsulfanilamide	40.4%
Inert Ingredients	59.6%
Total	100.0%

Contains 4.0 pounds of active ingredient per gallon.

Keep Out of Reach of Children

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

Shake Well Before Using

Agricultural Use Requirements
Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Personal Protective Equipment (PPE) and User Safety Recommendations, and Directions for Use including Storage and Disposal.

Shake Well Before Using:

Notice: Read the entire label. Use only according to label directions. Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.

In case of emergency endangering health or the environment involving this product, call collect 517-636-4400.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-113

EPA Est. 37429-GA-01

*Trademark of DowElanco
DowElanco • Indianapolis, IN 46268, U.S.A.

Specialty Herbicide

Net Contents XX gal

Table of Contents	Page
Precautionary Statements	
Hazards to Humans and Domestic Animals	
First Aid	
Environmental Hazards	
Directions for Use	
Storage and Disposal	
General Information	
General Use Precautions	
Soil Preparation	
Mixing Directions	
Surflan A.S. Alone	
Surflan A.S. Tank Mix Combinations	
Mixing Order	
Premixing	
Application Methods	
Ground Application	
Aerial Application	
Equipment Cleaning	
Activation and Cultivation	
Weeds Controlled by Surflan A.S.	
Weeds Suppressed by Surflan A.S.	
Approved Uses	
Ornamental Plantings	
Special Use Precautions	
Broadcast Application Rates	
Recommended Species Including Fruit-Plant-Nursery-Liners	
Field- and Liner-Container-Grown Species	
Container-Grown Species	
Field-Grown Fruit-Plant-Nursery-Liners <u>Non-bearing Trees and Vines</u>	
Tank Mix Combinations	
Ornamental Bulbs	
Special Use Precautions	
Broadcast Application Rates	
Greenhouse Areas	
Christmas Tree Plantations	
Surflan A.S. Alone	
Broadcast Application Rates	
Tank Mix Combinations	
Noncropland Areas and Industrial Sites	
Noncropland Areas-Tank Mix Combinations	
Industrial Sites-Tank Mix Combinations	
Warm Season Turfgrasses	
Special Use Precautions	
Annual Grasses Controlled by Surflan A.S.	
Annual Broadleaf Weeds Controlled by Surflan A.S.	
Broadleaf Weeds Suppressed by Surflan A.S.	
Application Rates, Frequency and Timing of Application	
Summer Annual Grasses and Broadleaf Weeds	
Single Application Program	
Split Application Program	
Annual Bluegrass (<i>Poa annua</i>) and Winter Annual Broadleaf Weeds	
Weed Control in Florida	

Application Equipment
Reseeding
Warranty Disclaimer
Inherent Risks of Use
Limitation of Remedies

7

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

Causes Eye Irritation • Harmful If Swallowed, Inhaled Or Absorbed Through The Skin

Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Surflan A.S. may cause skin sensitization reactions in certain individuals.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If in eyes: Flush with plenty of water. Get medical attention if irritation develops.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation develops.

If swallowed: Do not induce vomiting. Call a physician or Poison Control Center. If available, administer activated charcoal (6-8 heaping teaspoons) with a large quantity of water. Do not give anything by mouth to an unconscious person. Immediately transport to a medical care facility and see a physician.

If inhaled: Remove individual to fresh air. Get medical attention if breathing difficulty occurs. If not breathing, give artificial respiration, preferably cardiopulmonary resuscitation and get medical attention immediately.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Cover or incorporate spills.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Read All Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General Information

Surflan® A.S. herbicide is a preemergence surface-applied herbicide product for the control of many annual grasses and many broadleaf weeds in ornamental plantings, bulbs, ground covers/perennials, establishe ! warm-season turfgrass, Christmas tree plantations, non-bearing trees and vines, non-cropland and noncropland and industrial sites.

Shake Well Before Using:

—General Use Precautions

Surflan A.S. will not control emerged weeds. Poor weed control may result if directions are not followed. Over application may result in crop injury or excessive soil residue.

Surflan A.S. is orange in color and may cause temporary discoloration of sprayed surfaces. If this discoloration is undesirable, it may be altered by using a commercially available colorant such as Blazon or removed by spraying surface with water or washing with an industrial cleaner immediately after application. Surflan A.S. may also be applied with mulch colorants, such as Mulch Magic or Nu-Mulch.

Treatment of Plant Species Not Listed on the Label for Surflan A.S.

Users who wish to use Surflan A.S. on plant species not recommended on this label may determine the suitability for such uses use by treating a small number of such plants at a recommended rate. Prior to treatment of larger areas, the treated plants should be observed for any sign of herbicidal injury for during

30 to 60 days of normal growing conditions to determine if the treatment is non-injurious to the target plant species. The user assumes responsibility for any plant damage or other liability resulting from use of Surflan A.S. on plant species not recommended on this label.

~~Chemigation: Do not apply this product through any type of irrigation system.~~
(Ed. note: Chemigation statement added back under "Application"; see below)

Application

Soil Preparation

Surflan A.S. controls weeds growing from seed. Surflan A.S. will not control emerged weeds. Surflan A.S. does not control established weeds, weeds growing from stolens, rhizomes, or root pieces. Therefore, areas to be treated should be free of emerged weeds. Weed residues, prunings and trash should be thoroughly mixed into the soil or removed prior to treatment. In field applications, the soil should be in good tilth and free of clods at the time of application.

[Ed. Note: Sections which follow were rearranged and edited as indicated.]

Ground Application: Apply Surflan A.S. as a directed spray to the soil surface or over the top of plants using a vehicle-mounted, pull-type, or backpack sprayer. Use only a properly calibrated, low-pressure herbicide sprayer that will apply the spray uniformly. Use screens no finer than 50 mesh in nozzles and in-line strainers. Apply the appropriate rate of Surflan A.S. as outlined in "Approved Uses" section of this label. In all cases, use sufficient water volume to obtain uniform coverage and deliver the desired rate of Surflan A.S. to the treated area. The volume of water used is not critical, as long as the desired rate of Surflan A.S. is delivered uniformly across the area treated. When calibrating, determine the volume of water delivered by the sprayer to a given area (1,000 sq ft, 1 acre, etc.). Then mix the desired rate of Surflan A.S. in the amount of water required to cover the entire area to be treated. ~~Use only a properly calibrated low-pressure herbicide sprayer that will apply the spray uniformly. Use herbicide tips with screens no finer than 50 mesh in nozzles and in-line strainers.~~ As the amount of water used (spray volume) decreases, the importance of accurate calibration and uniform application increases. Check the sprayer daily to ensure proper calibration and uniform application. Maintain continuous agitation from mixing through application. Avoid spray pattern skips and overlaps that may result in incomplete coverage or over-application.

Hand Held or Backpack Sprayer Application: The amount of water used to apply Surflan A.S. herbicide is not critical, but should be sufficient for uniform coverage of the target area. Calibrate by determining the volume of water required to treat 1,000 square feet. Use this calibration volume to determine the amount of water and Surflan A.S. herbicide needed to treat the target area (see the following calibration example). Note: Sprayer calibration (volume of spray needed to treat 1,000 square feet) will vary with each individual operator.

Steps in Calibration:

1. Mark an area of 1,000 square feet (i.e. 20 by 50 feet or 25 by 40 feet).
2. Place the sprayer on a level surface and add water noting the final level of water in the spray tank.
3. Spray the marked area with a sufficient volume of water to provide uniform coverage. Refill the sprayer to the same level as before measuring the amount of water added. The measured water added to the sprayer is the volume needed to cover 1,000 square feet.
4. Determine the application rate (fl oz/1,000 sq ft) for Surflan A.S. from the Approved Uses section of this label.
5. To each volume of water used as measured in step 3, add the amount of Surflan A.S. as determined in step 4.

Example: If the sprayer used 2 gallons of water to cover 1,000 square feet and the desired application rate of Surflan A.S. is 3 fluid oz/1,000 square feet, then you would add 3 fluid ounces of Surflan A.S. to every 2 gallons of water to be used.

Aerial Application: Do not aerially apply this product. Use a standard aerial herbicide boom sprayer. Aerial spray equipment should be calibrated to apply the proper amount of Surflan A.S. alone or in tank mix combinations in 2 to 10 gallons of spray mixture per acre. Nozzle screens and in-line strainers should be no finer than 50 mesh. Surflan A.S. mixes readily with water for concentrate aerial sprays; however, constant vigorous agitation that sweeps the contents from the bottom of the spray tank up into the main body of the liquid is required to maintain a uniform suspension until the spray tank is empty. Avoid spray pattern skips and overlaps that may result in incomplete coverage or over-application. Do not apply when wind conditions favor drift from the target area.

Chemigation: Do not apply this product through any type of irrigation system.

Precaution: Avoid spray drift to non-target areas when applying Surflan A.S. Spray drift may result in reduced emergence of non-target plants adjacent to the treated area. Poor weed control may result if directions are not followed. Over-application may result in crop injury or excessive soil residue.

Mixing Directions

Shake Well Before Using

Precaution: Do not allow the spray mixture to siphon back into water source.

Surflan A.S. - Alone

Make sure spray tank is clean and use only clean water. Fill spray tank 1/2 to 3/4 full. Start agitation and add the required amount of Surflan A.S. Continue agitation and finish filling the spray tank. Maintain continuous agitation until application is completed.

Surflan A.S. - Tank Mix Combinations

Prior to mixing, read and carefully follow all label instructions and precautions for each product added to the tank mixture. Vigorous, continuous agitation is required for all Surflan A.S. tank mixes of Surflan A.S. Sparger pipe agitators generally provide the best agitation in spray tanks.

Precaution: Do not allow the spray mixture to siphon back into the water source.

Mixing Order: Fill the tank 3/4 full with clean water. Start agitation and add different formulation types in the order indicated below, allowing time for complete mixing and dispersion after addition of each product. Allow extra mixing and dispersion time for dry flowable products.

Add different formulation types in the following order: dry flowables (DF); wettable powders (WP); Surflan A.S. and other aqueous suspensions (AS), flowables (F) and liquids (L); solutions (S); and emulsifiable concentrates (EC).

Continue agitation and finish filling the spray tank with clean water. Maintain agitation until application is completed. If spraying and agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be completely resuspended before spraying is continued. A sparger agitator is particularly useful for this purpose.

Premixing: When tank mixing, initial mixing and dispersion of certain dry flowable or wettable powder products may be improved by premixing with water (slurrying). Adding the slurried material to the spray tank through a 20 or 35 mesh wetting screen will help assure good initial dispersion. Fine screens in the tank should be no finer than 50 mesh (100 mesh is finer than 50 mesh).

Application Methods

Ground Application: Apply Surflan A.S. as a directed spray to the soil surface or over the top of plants using a vehicle-mounted, pull-type, or backpack sprayer. Apply the appropriate rate of Surflan A.S. as outlined in "Approved Uses" section of this label. In all cases, use sufficient water volume to obtain uniform coverage and deliver the desired rate of Surflan A.S. to the treated area. The volume of water used is not critical, as long as the desired rate of Surflan A.S. is delivered uniformly across the area treated. When calibrating, determine the volume of water delivered by the sprayer to a given area (1,000 sq ft, 1 acre, etc.). Then mix the desired rate of Surflan A.S. in the amount of water required to cover the entire area to be treated. Use only a properly calibrated low-pressure herbicide sprayer that will apply the spray uniformly. Use herbicide tips with screens no finer than 50 mesh in nozzles and in-line strainers. As the amount of water used (spray volume) decreases, the importance of accurate calibration and uniform application increases. Check the sprayer daily to ensure proper calibration and uniform application. Maintain continuous agitation from mixing through application. Avoid spray pattern skips and overlaps that may result in incomplete coverage or over-application.

Aerial Application: Use a standard aerial herbicide boom sprayer. Aerial spray equipment should be calibrated to apply the proper amount of Surflan A.S. alone or in tank mix combinations in 2 to 10 gallons of spray mixture per acre. Nozzle screens and in-line strainers should be no finer than 50 mesh. Surflan A.S. mixes readily with water for concentrate aerial sprays; however, constant vigorous agitation that sweeps the contents from the bottom of the spray tank up into the main body of the liquid is required to maintain a uniform suspension until the spray tank is empty. Avoid spray pattern skips and overlaps that may result in incomplete coverage or over-application. Do not apply when wind conditions favor drift from the target area.

Equipment Cleaning

If a buildup of material occurs on the walk of the spray tank, it should be removed between fillings by washing with soap and water and rinsing thoroughly. Tanks, lines, screens and nozzles should be cleaned thoroughly after each use.

Activation and Cultivation

Surflan A.S. will remain stable on the soil surface up to 21 days following application. In the absence of timely rainfall, irrigation can be used to activate Surflan A.S. A minimum of one-half (1/2) inch of rain or its equivalent in sprinkler irrigation is necessary to activate Surflan A.S.. If weeds begin to emerge due to lack of rainfall or irrigation, shallow cultivate 1 to 2 inches deep to destroy existing weeds or remove them by hand. Shallow cultivation to a depth of 1 to 2 inches will enhance herbicidal effectiveness. If Surflan A.S. is not activated by rainfall, irrigation or cultivation within 21 days of application or existing weeds have not been removed, erratic weed control may result.

Weeds Controlled by Surflan A.S.

Annual Grasses:

Common Name	Scientific Name
barley, little	<i>Hordeum pusillum</i>
barnyardgrass (watergrass)	<i>Echinochloa crus-galli</i>
bluegrass, annual	<i>Poa annua</i>
crabgrass, large	<i>Digitaria sanguinalis</i>
crabgrass, smooth	<i>Digitaria ischaemum</i>
crowfootgrass	<i>Dactyloctenium aegyptium</i>
cupgrass, southwestern	<i>Eriochloa gracilis</i>
foxtail, bristleglass	<i>Setaria magna</i>
foxtail, giant	<i>Setaria faberi</i>

foxtail, green (pigeongrass)	<i>Setaria viridis</i>
foxtail, robust	<i>Setaria robusta</i>
foxtail, yellow	<i>Setaria glauca</i>
goosegrass (silver crabgrass)	<i>Eleusine indica</i>
Johnsongrass (seedling only)	<i>Sorghum halepense</i>
jungle rice	<i>Echinochloa colonum</i>
lovegrass, Mexican	<i>Eragrostis mexicana</i>
lovegrass, orcutt	<i>Eragrostis orcuttiana</i>
oat, wild	<i>Avena fatua</i>
panicum, browntop	<i>Panicum fasciculatum</i>
panicum, fall (spreading panicgrass)	<i>Panicum dichotomiflorum</i>
panicum, Texas (buffalograss) (Coloradograss)	<i>Panicum texanum</i>
ryegrass, Italian	<i>Lolium multiflorum</i>
sandbur, field	<i>Cenchrus incertus</i>
signalgrass (Brachiaria)	<i>Brachiaria</i> spp.
sprangletop, red	<i>Leptochloa filiformis</i>
witchgrass	<i>Panicum capillare</i>
Broadleaf Weeds:	Scientific Name
Common Name	
bittercress	<i>Cardamine oligosperma</i>
carpetweed	<i>Mollugo verticillata</i>
chickweed, common	<i>Stellaria media</i>
fiddleneck, coast	<i>Amsinckia intermedia</i>
filaree, redstem	<i>Erodium cicutarium</i>
filaree, whitestem	<i>Erodium moschatum</i>
groundsel, common	<i>Senecio vulgaris</i>
henbit	<i>Lamium amplexicaule</i>
knotweed, prostrate	<i>Polygonum aviculare</i>
lambquarters	<i>Chenopodium album</i>
pigweed, prostrate	<i>Amaranthus blitoides</i>
pigweed, redroot	<i>Amaranthus retroflexus</i>
pigweed, spring	<i>Amaranthus hybridus</i>
pigweed, tumble	<i>Amaranthus albus</i>
puncturevine	<i>Tribulus terrestris</i>
purslane, common	<i>Portulaca oleracea</i>
pusley, Florida (Florida purslane) (Mexican clover) (pusley)	<i>Richardia scabra</i>
rocket, London	<i>Sisymbrium irio</i>
rockpurslane, desert	<i>Calandrinia ciliata</i>
shepherdspurse	<i>Capsella bursa-pastoris</i>
spurge, prostrate	<i>Euphorbia humistrata</i>
woodsorrel, yellow	<i>Oxalis stricta</i>

Weeds Suppressed by Surflan A.S.

Control of the following weeds may be erratic, ranging from poor to excellent, depending upon soil temperature, time of germination, depth of seed in the soil, and amount and timing of soil moisture:

Common Name	Scientific Name
horseweed	<i>Coryza canadensis</i>
ladysthumb	<i>Polygonum persicaria</i>
lettuce, prickly	<i>Lactuca scariola</i>
mallow, common	<i>Malva neglecta</i>
milkweed, climbing	<i>Sarcostemma cynanchoides</i>
momingglory	<i>Ipomoea</i> spp.
mustard, black	<i>Brassica nigra</i>
mustard, wild	<i>Brassica kaber</i>
nightshade, black	<i>Solanum nigrum</i>
ragweed, common	<i>Ambrosia artemisiifolia</i>
smartweed	<i>Polygonum pennsylvanicum</i>
sowthistle, annual	<i>Sonchus oleraceus</i>
spurge, spotted	<i>Euphorbia maculata</i>
teaweed (prickly sida)	<i>Sida spinosa</i>
velvetleaf	<i>Abutilon theophrasti</i>
wheat, volunteer	<i>Triticum</i> spp.

Approved Uses

Ornamental Plantings

[Ed. Note: sections rearranged and edited as shown.]

Recommended Species Including Fruit-Plant-Nursery Liners

Surflan A.S. is recommended for use on certain landscape container- and field-grown established ornamental plants, including: trees, and shrubs, established ground covers/perennials, flowers, non-bearing fruit and nut trees, nonbearing vineyards, field-grown fruit tree and shrub nursery liners; and in the production of ornamental bulbs (See "Ornamental Bulbs" section for special use directions).

Broadcast Application Rates

Length of Control	Surflan A.S.	
	quarts/acre	fl oz/1000 sq ft
2 to 4 months	2	1.5
4 to 8 months	4	3

[Editors note: Revised section on backpack sprayers previously at this location and moved to General Information section.]

Tank Mix Combinations

Tank mix combinations of Surflan A.S. plus Roundup, and many other labeled herbicides may be used to control undesirable vegetation in ornamental areas. Surflan A.S. may also be tank mixed with Gallery® herbicide and applied preemergence to broaden the spectrum of broadleaf weed control in ornamental areas. Applied as directed, these tank mixes of Surflan A.S. will provide control of susceptible weed species listed on the respective labels. Refer to tank mix product labels for specific use directions, precautions and limitations before use.

Surflan A.S. Plus Roundup: Tank mix combinations of Surflan A.S. plus Roundup are recommended to control existing undesirable vegetation. Applied as directed, Surflan A.S. plus Roundup will provide postemergence control of susceptible weed species listed on the label for Roundup and residual preemergence control of susceptible weed species listed on the label for Surflan A.S. Refer to the label for Roundup for specific use directions, precautions and limitations before use.

Precautions: Do not apply sprays containing Roundup over the top of ornamental plants. Extreme care must be exercised to prevent sprays containing Roundup from coming in contact with foliage and stems of turfgrasses, trees, shrubs, or other desirable vegetation since severe damage or death may result. If spraying with Roundup in areas adjacent to desirable plants, use a shield to prevent spray from contacting foliage and stems of desirable plants.

Special Use Precautions:

Apply only to established plantings.—Established plants are defined as those that have been transplanted into their growing location for a sufficient period of time to allow the soil to be firmly settled around the roots from packing and rainfall or irrigation.

Rooted liners should be removed from their original growing containers and placed in new containers at least two weeks prior to treatment or injury may occur.

To avoid possible injury, do not apply Surflan A.S. to:

- Nursery, forest, or Christmas tree seedling beds, cutting beds, or transplant beds.
- ~~Either nursery seedbeds or forest or Christmas tree seedling-transplant beds.~~
- Unrooted liners or cuttings that have been planted in pots for the first time.
- Pots less than four inches wide.
- Ground covers until they are established and well rooted.
- Ornamental plantings where there is likelihood of runoff onto lawn areas.
- Areas containing dichondra or cool season turfgrass species.

On container grown ornamentals where weed seed germination continues for extended periods of time, do not make repeat applications of Surflan A.S. for at least 90 days or crop injury may occur.

Applications of Surflan A.S. over the top of plants with newly forming buds may cause injury. In this situation a directed spray is recommended.

For soils treated with Surflan A.S. during the previous season, plant only the ornamental species listed on this label or injury may occur.

Ice Plant: When establishing unrooted ice plant (*Mesembryanthemum crystallinum* and *Carpobrotus edulis*) on coarse textured soils in landscape plantings, use only do not exceed the 2 quart per acre rate of Surflan A.S. or crop injury may occur. After the ice plant is well established, a second application may be made.

[Ed. note: This section moved to General Information section.]

~~Hand Held or Backpack Sprayer Application~~

~~Apply Surflan A.S. at a rate of 1.5 to 3 ounces per 1000 sq. m. feet. The amount of water used to apply Surflan A.S. is not critical, but should be sufficient for uniform treatment of the target area. Calibrate by determining the volume of water required to treat 1000 square feet. Use this calibration volume to determine the amount of water and Surflan A.S. needed to treat the target area (see following table). Note: Sprayer calibration (volume of spray needed to treat 1,000 square feet) will vary with each individual operator.~~

Length of Control	Application Rate (fl-oz/1000 sq-ft)	Quantity of Water Needed
2-to-4 months	1.5	The amount required by your sprayer to cover 1000 sq-ft of area
4-to-8 months	2.0	

Sample Calculation:

Size of target area \times 1000 \times Application rate = Amount of Surflan A.S. required

Size of target area \times 1000 \times Calibration volume per 1000 sq-ft = Amount of water required

Do not apply Surflan A.S. to the following plant species when container grown or field grown or injury may occur:

Note: Injury on the following plant species has been observed following applications of Surflan A.S. and use is not recommended:

- Deutzia gracilis* (slender deutzia)
- Pseudotsuga manziesii* (Douglas-fir)
- Thuja occidentalis* 'Techny' (Techny arborvitae)
- Tsuga canadensis* (eastern hemlock)
- Beconia* spp. (beconia)
- Coleus hybridus* (coleus)

Surflan A.S. May be Used on the Following **Established Plant Species: Field and Liner†-Grown Plants and Plants in Landscape Plantings:**

(Note: Limitations on recommended treatment methods.)

†Plants transplanted for additional growth before transplanting to final growing location.

[Editor's note: The plant species grouped as follows: Trees, Shrubs, Groundcovers / Perennials, Flowers, Nonbearing fruit and nut trees and non-bearing vineyards.]

Trees

Recommended Treatment Method:

F = Field Grown

C = Container Grown

Scientific Name	Common Name	
<i>Abies balsamea</i>	Fir, balsam	F
<i>Abies concolor</i>	Fir, white	F
<i>Abies fraseri</i>	Fir, fraser	F
<i>Abies grandis</i>	Fir, grand	F
<i>Abies veitchi</i>	Fir, "etch	F
<i>Abies lasiocarpa</i>	Fir, alpine	F
<i>Abutilon hybridum</i>	Albus-flowering maple	F
	Luteus-flowering maple	F
	Roseus-flowering maple	F
	Tangerine-flowering maple	F
	Vesuvius red-flowering maple	F
<i>Acer ginnala</i>	Flame maple	F
<i>Acer rubrum</i>	Red sunset maple	F
<i>Acer saccharinum</i>	Silver maple	F
<i>Acer</i> spp.	Maple	F
<i>Arecastrum romanzoffianum</i>	Queen palm	F
<i>Betula nigra</i>	Birch, river	F
<i>Betula papyrifera</i>	Paper birch	F
<i>Betula pendula</i>	Birch, white	F
<i>Bucida buceras</i>	Black olive	F

<i>Carya</i> spp.	Pecan, ornamental	C,F
<i>Ceratonia siliqua</i>	Carob	F
<i>Cercidium floridum</i>	Palo Verde, blue	F
<i>Cercis canadensis</i>	Redbud	C,F
<i>Chamaecyparis lawsoniana</i>	Falsecypress, Lawson	F
<i>Chamaecyparis obtusa</i>	Filicoides-fernspray cypress	F
	Gracilis-slender Hinoki cypress	F
	Sawara-false cypress	F
<i>Chamaecyparis pisifera</i>	Squarrosa-moss cypress	F
	Cat Palm	F
<i>Chamaedorea cataractarum</i>	Palm	F
<i>Chamaedorea costaricana</i>	Parlor palm	F
<i>Chamaedorea elegans</i>	Citrus, ornamental	C,F
<i>Citrus</i> spp.	Dogwood, flowering	F
<i>Cornus florida</i>	Cryptomeria, Japanese	C,F
<i>Cryptomeria japonica</i>	Carrot wood	F
<i>Cupaniopsis anacardioides</i>	Cypress, Arizona	C,F
<i>Cupressus arizonica (glabra)</i>	Arizona cypress	C,F
<i>Cupressus glabra</i>	Cypress, Italian	C,F
<i>Cupressus sempervirens</i>	Russian olive	C,F
<i>Elaeagnus angustifolia</i>	Red gum eucalyptus	F
<i>Eucalyptus camaldulensis</i>	Eucalyptus, mealy	F
<i>Eucalyptus cinerea</i>	Silver dollar eucalyptus	F
	Eucalyptus, narrow-leaved	F
<i>Eucalyptus nicholii</i>	Eucalyptus, red ironbark	F
<i>Eucalyptus sideroxylon</i>	Ficus	F
<i>Ficus benjamina</i>	Ash	F
<i>Fraxinus</i> spp.	Ginkgo (Maidenhair tree)	C,F
<i>Ginkgo biloba</i>	Honey locust	F
<i>Gleditsia triacanthos</i>	Toyon	F
<i>Heteromeles arbutiflora</i>	Redcedar, Eastern	F
<i>Juniperus virginiana</i>	Goldenrain tree	F
<i>Koelreuteria paniculata</i>	Sweetgum, American	C,F
<i>Liquidambar styraciflua</i>	Magnolia, Southern	F
<i>Magnolia grandiflora</i>	Crabapple	F
<i>Malus</i> spp.	White mulberry	F
<i>Morus alba</i>	Pendula-weeping Norway spruce	F
<i>Picea abies</i>	Repens-spreading Norway spruce	F
	Spruce, Norway	F
<i>Picea engelmannii</i>	Spruce, Englemann	F
<i>Picea glauca</i>	Spruce, white	F
	Conica-dwarf Alberta spruce	F
<i>Picea glauca conica</i>	Dwarf Alberta spruce	F
<i>Picea mariana</i>	Spruce, black	F
<i>Picea pungens</i>	Glauca-Colorado blue spruce	F
	Hoopsii-Horp's blue spruce	F
	Koster-Koster blue spruce	F
	Spruce, Colorado	C,F
<i>Pinus aristata</i>	Bristlecone pine	F
<i>Pinus canariensis</i>	Canary Island pine	F
<i>Pinus contorta</i>	Shore pine, beach pine	F
<i>Pinus edularica</i>	Eldarica pine	F
<i>Pinus radiata</i>	Monterey pine	F
<i>Pinus</i> spp.	Pine	C,F

<i>Pinus strobus</i>	Eastern white pine	F
<i>Pinus sylvestris</i>	Scotch pine	F
<i>Pinus thunbergiana</i>	Japanese black pine	F
<i>Platanus occidentalis</i>	American sycamore	F
<i>Platanus racemosa</i>	California sycamore	F
<i>Podocarpus spp.</i>	Podocarpus	F
<i>Populus deltoides</i>	Cottonwood	F
	Cottonwood (grown for pulp)	F
<i>Prunus caroliniana</i>	Laurelcherry, Carolina	F
<i>Prunus laurocerasus</i>	Laurelcherry, English	F
<i>Prunus mahaleb</i>	Cherry, Mahaleb	F
<i>Prunus yedoensis</i>	Yoshino flowering cherry	F
<i>Pyrus communis</i>	Pear	F
<i>Quercus ilicifolia</i>	Bear oak	F
<i>Quercus palustris</i>	Pin oak	F
<i>Quercus phellos</i>	Willow oak	F
<i>Quercus rubra</i>	Red oak	C,F
<i>Quercus spp.</i>	Oak	C,F
<i>Salix babylonica</i>	Babylon weeping willow	F
	Corkscrew willow	F
<i>Schinus molle</i>	California pepper tree	F
<i>Sequoia sempervirens</i>	Redwood, coast	F
<i>Sequoiadendron giganteum</i>	Giant sequoia	F
<i>Swietenia mahogani</i>	Mahogany	F
<i>Tabebuia caraiba</i>	Yellow tab	F
<i>Tilia cordata</i>	Linden, little leaf	C,F
<i>Ulmus parvifolia</i>	Chinese elm	F
<i>Umbellularia californica</i>	California laurel	F
<i>Washingtonia robusta</i>	Mexican fan palm	F

Shrubs**Recommended Treatment Method:**

F = Field Grown

C = Container Grown

Scientific Name	Common Name	
<i>Abelia grandiflora</i>	Glossy abelia	F
<i>Acacia redolens</i>	Acacia, prostrate	F
<i>Agave americana</i>	Century plant	F
<i>Agave macroculmis</i>	Agave	F
<i>Arctostaphylos stanfordiana</i>	Manzanita, Stanford	F
<i>Astilbe chinensis</i>	Astilbe/false spirea	C,F
<i>Baccharis pilularis</i>	coyotebush	F
<i>Berberis thunbergii</i>	Aurea-golden Japanese barberry	C,F
	Crimson pygmy barberry	C,F
	Atropurea-redleaf Japanese barberry	C,F
	Barberry, Japanese	C,F
<i>Bougainvillea spp.</i>	Barbara Karst	F
	California gold	F
	Scarlet O'Hara	F
	Texas dawn	F
<i>Buxus microphylla</i>	Littleleaf boxwood	F
<i>Buxus microphylla japonica</i>	Boxwood, Japanese	C,F
<i>Buxus sempervirens</i>	Boxwood, common	C,F
<i>Callistemon citrinus</i>	Bottlebrush, lemon	C,F

<i>Cassia artemisioides</i>	Cassia, feathery	F
<i>Ceanothus americanus</i>	Jerseytea, redroot	C,F
<i>Ceanothus</i> spp.	Wild lilac	C,F
<i>Chamaecyparis obtusa</i>	Kosteri cypress	F
	Nana-dwarf Hinoki cypress	F
	Torulosa cypress	F
<i>Chamaecyparis pisifera</i>	Squarrosa Minima cypress	F
<i>Chamaecyparis pisifera</i> spp.	Filifera-thread cypress	F
<i>Chrysalidocarpus lutescens</i>	Areca palm	F
<i>Cleyera japonica</i>	Cleyera, Japanese	C,F
<i>Cornus alba</i>	Sibirica-Siberian dogwood	F
<i>Cornus kousa</i>	Dogwood, kousa	C,F
<i>Cornus stolonifera</i>	Flaviramea-yellowtwig dogwood	F
<i>Cotoneaster adpressus</i>	Praecox-early cotoneaster	F
<i>Cotoneaster apiculatus</i>	Cotoneaster, cranberry	C,F
<i>Cotoneaster buxifolius</i>	Cotoneaster, brightbead	F
<i>Cotoneaster congestus</i>	Cotoneaster, Pyrenees	F
<i>Cotoneaster dammeri</i>	Cotoneaster, bearberry	C, F
<i>Cotoneaster himalayan</i>	Himalayan cotoneaster	F
<i>Cotoneaster horizontalis</i>	Cotoneaster, rock	C,F
<i>Cotoneaster lacteus</i>	Cotoneaster, parney	C,F
<i>Cotoneaster microphyllus</i>	Cotoneaster, rockspray	F
<i>Cytisus praecox</i>	Hollandia-warminster broom	F
<i>Cytisus scoparius</i>	Lena-Scotch broom	F
<i>Dasylinon wheeleri</i>	Sotol, desert spoon	F
<i>Deutzia crenata</i>	Nakiana-dwarf deutzia	F
<i>Dodonaea viscosa</i>	Hopseedbush, clammy	F
	Hopseed bush	F
<i>Escallonia axoniensis</i>	Escallonia	C,F
<i>Euonymus alata</i>	Euonymus, winged	F
<i>Euonymus fortunei</i>	Canadale gold euonymus	C,F
	Emerald'n gold euonymus	C,F
	Euonymus, stringybark	C,F
	Wintercreeper	C,F
<i>Euonymus japonica</i>	Euonymus, evergreen	C,F
	Silver king euonymus	F
	Spreading euonymus	F
<i>Euonymus kiatschovica</i>	Fatsyhedera	C,F
<i>Fatsyhedera lizei</i>	Forsythia, border	F
<i>Forsythia intermedia</i>	Gardenia	C,F
<i>Gardenia jasminoides</i>	Woadwaxen	F
<i>Genista pilosa</i>	Ross Estey-hibiscus	F
<i>Hibiscus rosa-sinensis</i>	Hibiscus, Chinese	F
	Rose of Sharon, Red Bird	F
	Rose of Sharon, Red Heart	F
	Rose of Sharon, Woodbridge	F
	Rose-of-Sharon (Shrubalthea)	F
<i>Ilex aquifolium</i>	Balkans holly	F
	Gold coast holly	F
	Holly, English	F
<i>Ilex aquipernyi</i>	San Jose holly	C,F
<i>Ilex cornuta</i>	Burford holly	C,F
	Holly, Chinese	C,F
<i>Ilex crenata</i>	Compacta-dwarf Japanese holly	C,F

	Convexa holly	C,F
	Dwarf Chinese holly	
	Helleri-Heller's Japanese holly	C,F
	Holly, Japanese	C,F
<i>Ilex glabra</i>	Nordica-inkberry holly	F
<i>Ilex meserveae</i>	Blue boy holly	F
	Blue girl holly	F
	Ebony magic holly	F
<i>Ilex vomitoria</i>	Nana-dwarf yaupon holly	C,F
	Pendula-weeping yaupon holly	C,F
	yaupon holly	C,F
<i>Juniperus conferta</i>	Emerald sea shore juniper	F
<i>Juniperus horizontalis</i>	Huntington blue juniper	C,F
	Wiltonii-blue carpet juniper	C,F
<i>Juniperus procumbens</i>	Nana-dwarf Japanese garden juniper	C,F
<i>Juniperus prostrata</i>	Prostrata juniper	C,F
<i>Juniperus sabina</i>	Broadmoor juniper	F
	Foemina-Hicks juniper	F
	Tamariscifolia-Tam juniper	F
<i>Juniperus scopulorum</i>	Emerald green juniper	F
<i>Juniperus spp.</i>	Juniper	C,F
<i>Juniperus squamata</i>	Blue juniper	F
	Blue star juniper	F
	Parsonii juniper	F
<i>Justicia brandegeana</i>	Shrimp plant	C,F
<i>Justicia spicigera</i>	Honeysuckle, Mexican	F
<i>Kalmia latifolia</i>	Laurel, mountain	F
<i>Lagerstroemia indica</i>	Crape myrtle	C,F
<i>Leucothoe axillaris</i>	Leucothoe, coast	F
<i>Leucothoe fontanesiana</i>	Leucothoe, drooping	F
<i>Ligustrum amurense</i>	Privet, amur	C,F
<i>Ligustrum japonicum</i>	Privet, Japanese	C,F
	yellow tip ligustrum	C,F
<i>Ligustrum lucidum</i>	Privet, glossy	C,F
<i>Ligustrum ovalifolium</i>	California privet	F
<i>Ligustrum texanum</i>	Howardi privet	F
	wax leaf privet	F
<i>Ligustrum vicaryi</i>	Privet, golden	C,F
	Vicary golden privet	C,F
<i>Livistona chinensis</i>	Chinese fountain palm	F
<i>Lonicera fragrantissima</i>	Winter honeysuckle	F
<i>Lonicera perichlymenum</i>	Flowering woodbine	F
	Serotina woodbine	F
<i>Lonicera sempervirens</i>	Trumpet honeysuckle	F
<i>Mahonia aquifolium</i>	Oregon grape	F
<i>Myoporum parvifolium</i>	Myoporum, prostrate	F
<i>Myrtus communis</i>	Myrtle, true	C,F
<i>Nandina domestica</i>	Compacta-dwarf heavenly bamboo	C,F
	Harbour dwarf-heavenly bamboo	C,F
	Heavenly bamboo (Nandina)	C,F
	Nana compacta-heavenly bamboo	C,F
	Nana purpurea-heavenly bamboo	C,F
	Woods dwarf-heavenly bamboo	C,F
<i>Nerium oleander</i>	Hardy red oleander	C,F

	Oleander	C,F
	Ruby lace oleander	C,F
<i>Osmanthus heterophyllus</i>	Osmanthus, holly-leaf	F
<i>Philadelphus</i> spp.	Mockorange	C,F
<i>Phoenix roebelenii</i>	Pigmy date palm	F
<i>Photinia fraseri</i>	Fraser's photinia	C,F
	Photinia	C,F
<i>Pieris japonica</i>	Lily-of-the-valley	F
	Snowdrift lily-of-the-valley	F
	Temple bells lily-of-the-valley	F
	Valley rose lily-of-the-valley	F
	Andromeda	C,F
<i>Pittosporum</i> spp.	Pittosporum	C,F
<i>Pittosporum tobira</i>	Green pittosporum	F
	Japanese pittosporum	
	Tobira	F
	Wheeler's dwarf pittosporum	F
<i>Platycladus orientalis</i>	Arborvitae, Oriental	C,F
<i>Plumbago ariculata</i>	Blue cape plumbago	F
<i>Podocarpus macrophyllus</i>	Yewpine	C,F
<i>Potentilla fragiformis</i>	Cinquefoil	F
<i>Potentilla fruticosa</i>	Cinquefoil	C,F
<i>Protea neriifolia</i>	Protea	F
<i>Pyracantha coccinea</i>	Firethorn, scarlet	C,F
<i>Pyracantha fortuneana</i>	Lolendei Monrovia pyracantha	C,F
<i>Pyracantha fortuneana</i>	Monon pyracantha	C,F
	Red elf hybrid pyracantha	C,F
	Rutgers hybrid pyracantha	C,F
	Santa Cruz pyracantha	C,F
	Victory pyracantha	C,F
<i>Pyracantha skoidzumi</i>	Firethorn, formosa	C,F
<i>Pyracantha, fortuneana</i>	Firethorn	C,F
<i>Raphiolepis indica</i>	Enchantress-Moness raphiolepis	F
	Raphiolepis (India hawthorn)	C,F
	Springtime-Monme raphiolepis	F
	Roundleaf raphiolepis	F
<i>Raphiolepis ovata</i>	Eastercactus	C,F
<i>Rhipsalidopsis gaertneri</i>	Flame azalea	F
<i>Rhododendron calendulaceum</i>	Butterfly rhododendron	F
<i>Rhododendron campylocarpum</i>	PJM rhododendron	F
<i>Rhododendron carolinianum x daurium</i>	Catawba album rhododendron	C,F
<i>Rhododendron catawbiense</i>	Catawba rhododendron	C,F
	Lord Roberts rhododendron	C,F
	Rocket rhododendron	C,F
<i>Rhododendron caucasicum x ponticum</i>	Cunningham White rhododendron	F
<i>Rhododendron forrestii x griersonianum</i>	Elizabeth rhododendron	F
<i>Rhododendron griffithianum</i>	Jean Marie rhododendron	F
<i>Rhododendron hybrid</i> spp.	America rhododendron	F
	English Roseum rhododendron	F
	Nova Zembla rhododendron	F
	Scintillation rhododendron	F
<i>Rhododendron impeditum</i>	Rhododendron	F
<i>Rhododendron indica</i>	Formosa azalea	C,F
	Waucabusa azalea	C,F

<i>Rhododendron kerume</i>	Coral bells azalea	C,F
	Hino crimson azalea	C,F
	Hino pink azalea	C,F
	Snow azalea	C,F
<i>Rhododendron maximum</i>	Rhodie max (rosebay)	C,F
<i>Rhododendron mucronulatum</i>	Rhododendron	F
<i>Rhododendron ponticum</i>	Chionoides rhododendron	F
<i>Rhododendron sassthigiatim x carolinianum</i>	Ramapo rhododendron	F
<i>Rhododendron satuski</i>	Gumpo pink azalea	F
	Higasa azalea	F
<i>Rhododendron spp.</i>	Azalea	C,F
	Rhododendron	C,F
<i>Rhododendron spp. hybrids</i>	Carror azalea	C,F
	Girard Roberta azalea	F
	Golden flare exbury azalea	F
	Sumac, African	C,F
<i>Rhus lancea</i>	Ramanas rose	F
<i>Rosa rugosa</i>	Rosemary	F
<i>Rosmarinus officinalis</i>	Bridal wreath	F
<i>Spiraea vanhouttei</i>	Lilac, common	C,F
<i>Syringa vulgaris</i>	Brush cherry	C,F
<i>Syzygium paniculata</i>	Yew, Japanese	F
<i>Taxus cuspidata</i>	Yew	F
<i>Taxus media</i>	Arborvitae, American	C,F
<i>Thuja occidentalis</i>	Emerald arborvitae	F
	Globosa-globe arborvitae	F
	Little giant-dwarf arborvitae	F
	Nigra-dark American arborvitae	F
	Pyramidalis arborvitae	F
	Rheingold arborvitae	F
	Woodwardii arborvitae	F
<i>Thuja orientalis</i>	Aureus nana-dwarf golden arborvitae	F
	Minima glauca-dwarf arborvitae	F
<i>Thuja plicata</i>	Red Cedar, Western	F
<i>Trachelospermum jasminoides</i>	Star jasmine, Chinese	F
<i>Veitchia merrilli</i>	Christmas palm	F
<i>Viburnum davidi</i>	David viburnum	F
<i>Viburnum japonicum</i>	Viburnum	F
<i>Viburnum opulus sterile</i>	Common snowball viburnum	F
<i>Viburnum plicatum tomentosum</i>	Doublefile viburnum	F
<i>Viburnum setigerum</i>	Tea viburnum	F
<i>Viburnum suspensum</i>	Viburnum, Sandankwa	F
<i>Viburnum tinus</i>	Viburnum, Laurustinus	C,F
	Compactum-spring bouquet viburnum	F
<i>Viburnum tinus compactum</i>	Spring bouquet viburnum	F
<i>Viburnum trilobum compactum</i>	Dwarf cranberry bush	F
<i>Viburnum x pragense</i>	Viburnum	F
<i>Weigela florida</i>	Bristol ruby weigela	F
	Java red weigela	F
	Minuet weigela	F
	Weigela, oldfashioned	F
<i>Xylosma congestum</i>	Xylosma	F
<i>Yucca elata</i>	Yucca, soaptree	C,F
<i>Yucca recurvifolia</i>	Yucca, pendulous	F

Groundcovers/Perennials

Recommended Treatment Method:
F = Field Grown
C = Container Grown

Scientific Name	Common Name	
<i>Agapanthus africanus</i>	Lily-of-the-Nile	C,F
<i>Ajuga</i> spp.	Carpet bugle	F
<i>Arctotheca calendula</i>	Cape weed	F
<i>Campanula elatines</i>	Bellflower	C,F
<i>Carpobrotus edulis</i>	Ice plant, largeleaf (see label)	F
<i>Clytostoma callistegioides</i>	Trumpet vine, violet	C,F
<i>Cortaderia selloana</i>	Pampas grass	F
<i>Delosperma alba</i>	White iceplant	F
<i>Drosanthemum floribundum</i>	Trailing rosea iceplant	F
<i>Festuca ovina glauca</i>	Blue fescue	F
<i>Gazania</i> spp.	Gazania	F
<i>Hedera canariensis</i>	Ivy, Algerian	F
<i>Hedera helix</i>	Ivy, English	F
<i>Hemerocallis</i> spp.	Daylily	C,F
<i>Hosta lancifolia</i>	Albo-marginata hosta	C,F
<i>Hosta</i> spp.	Lily, plantain	C,F
<i>Hypericum</i> spp.	St. Johnswort	C,F
<i>Lampranthus spectabilis</i>	Trailing iceplant	F
<i>Liriope gigantea</i>	White lily turf	F
<i>Liriope muscari</i>	Lilac beauty lily turf	C,F
	Majestic lily turf	C,F
	Monroe white lily turf	C,F
	Silvery sunproof lily turf	C,F
	Variogated liriope lily turf	C,F
	Big blue lily turf	C,F
<i>Lonicera japonica</i>	Honeysuckle, Japanese	F
<i>Mesembryanthemum crystallinum</i>	Ice plant (see label)	F
<i>Ophiopogon japonicus</i>	Mondo grass	F
<i>Osteospermum fruticosum</i>	Daisy, trailing African	F
<i>Pachysandra terminalis</i>	Japanese spurge	F
<i>Sedum brevifolium</i>	Stoncrop	F
<i>Vinca major</i>	Periwinkle, bigleaf	F
<i>Vinca minor</i>	Periwinkle, dwarf	F

Flowers

Recommended Treatment Method:
F = Field Grown
C = Container Grown

Scientific Name	Common Name	
<i>Achillea</i> spp.	Yarrow	F
<i>Antirrhinum majus</i>	Snapdragon	F
<i>Caladium bicolor</i>	Caladium, fancy leafed	F
<i>Chrysanthemum coccineum</i>	Daisy, painted	F
<i>Chrysanthemum maximum</i>	Daisy, Shasta	F
<i>Chrysanthemum morifolium</i>	Chrysanthemum	F
<i>Cladium bicolor</i>	Fancy-leafed caladium	F
<i>Coreopsis lanceolata</i>	Coreopsis	F
<i>Dianthus barbatus</i>	Sweet William	F
<i>Dicentra spectabilis</i>	Bleeding heart	C,F
<i>Dimorphotheca</i> spp.	Marigold, cape	F

<i>Echinacea purpurea</i>	Coneflower, purple	F
<i>Geum quellyon</i>	Geum	F
<i>Gladiolus hortulanus</i>	Gladiolus	F
<i>Gypsophila paniculata</i>	Baby's breath	F
<i>Impatiens wallerana</i>	Impatiens (Busy lizzie)	F
<i>Iris spp.</i>	Iris, bearded	F
<i>Liatris spicata</i>	Blazing star	C,F
<i>Pelargonium hortorum</i>	Geranium	F
<i>Petunia spp.</i>	Petunia	F
<i>Portulaca grandiflora</i>	Moss, rose	F
<i>Ranunculus asiaticus</i>	Ranunculus, Persian	F
<i>Rosa spp.</i>	Rose	F
<i>Rudbeckia hirta</i>	Daisy, gloriosa (black-eyed Susan)	F
<i>Salvia spp.</i>	Salvia (Sage)	F
<i>Stokesia laevis</i>	Aster, stokes	F
<i>Strelitzia reginae</i>	Bird of paradise	F
<i>Tagetes spp.</i>	Marigold	F
<i>Viola wittrockiana</i>	Pansy	F
<i>Zinnia elegans</i>	Zinnia, common	F

Non-bearing trees and vines

Recommended Treatment Method:
F = Field Grown
C = Container Grown

almond	F
apple	F
apricot	F
avocado	F
blackberry	F
blueberry	F
boysenberry	F
cherry, sour	F
cherry, sweet	F
currant	F
dewberry	F
elderberry	F
fig	F
filbert	F
gooseberry	F
grape, American	F
grape, European	F
grapefruit	F
kiwi	F
Kumquat	C,F
lemon	F
loganberry	F
macadamia nut	F
nectarine	F
olive	F
orange	C,F
peach	F
pear	F
pecan	C,F
pistachio	F
plum	F

pomegranate	F
prune	F
raspberry	F
walnut, black	F
walnut, English	F

*Nonbearing plants are defined as those which will not bear fruit for at least one year after treatment.

Ornamental Bulbs

Surflan A.S. may be applied for control of susceptible annual weeds in ornamental bulbs, e.g., bulbous iris, daffodil (narcissus), hyacinth and tulip. Apply Surflan A.S. to the soil surface 2 to 4 weeks after planting, but prior to the emergence of annual weeds. For fall planted bulbs, apply Surflan A.S. again in late winter or early spring to weed-free soil surfaces.

Special Use Precautions:

Do not apply to tulip plants that have emerged to a height greater than 3/4 inch.
Do not apply to gladioli corms prior to emergence or less than one inch in diameter.

Broadcast Application Rates

Time of Application	Soil Texture	Surflan A.S.	
		quarts/acre	fl oz/1000 sq ft
Fall	Coarse	0.75	0.5
Fall	Medium and Fine	1.5	1.0
Feb. - March	All Soil Textures	0.75	0.5

Special Use Precautions:

Do not apply to tulip plants that have emerged to a height greater than 3/4 inch.
Do not apply to gladioli corms prior to emergence or less than one inch in diameter.

Shadehouse Greenhouse Areas

Surflan A.S. may be applied to drainage areas under benches in open greenhouse shadehouse-type structures where the natural flow of air is unimpeded. Do not apply in enclosed greenhouses or in enclosed shadehouse-type structures. Do not apply within three weeks prior to enclosure of in greenhouse or poly-type structures.

Christmas Tree Plantations

Surflan A.S. Alone

Apply Surflan A.S. as a directed spray to the soil surface or as an overtop spray to established plantings of field grown Christmas tree species, including fir (*Abies* spp.), pine (*Pinus* spp.), and spruce (*Picea* spp.). Do not apply to Douglas fir (*Pseudotsuga menziesii*). Do not apply to seedbeds or seedling transplant beds. Apply only to established plantings. Established plants are defined as those that have been transplanted into their final growing location for a sufficient period of time to allow the soil to be firmly settled around the roots from packing and rainfall or irrigation. Follow all instructions provided in the "General Information" section of this label.

Broadcast Application Rates

Length of Control	Surflan A.S.	
	quarts/acre	fl oz/1000 sq ft
2 to 4 months	2	1.5
4 to 8 months	4	3

Tank Mix Combinations

~~Tank mix~~ Tank mix combinations of Surflan A.S. plus other labeled herbicides may be used as directed or overtop sprays in established Christmas tree plantings. When applied according to use directions, these tank mixes will provide control of susceptible weed species listed on the respective product labels. Refer to tank mix product labels for specific use directions, precautions and limitations before use.

Surflan A.S. Plus Roundup: Apply tank mix combinations of Surflan A.S. plus Roundup only as directed sprays in Christmas tree plantings. When applied according to use directions, Surflan A.S. plus Roundup will provide postemergence control of susceptible weed species listed on the Roundup label and residual preemergence control of susceptible weed species listed on the Surflan A.S. label. Refer to the Roundup label for specific use directions, precautions and limitations before use.

Special Use Precautions:

Do not apply to Douglas-fir (*Pseudotsuga menziesii*). Do not apply to seedbeds or seedling transplant beds. Apply only to established plantings. ~~Established plants are defined as those that have been transplanted into their final growing location for a sufficient period of time to allow the soil to be firmly settled around the roots from packing and rainfall or irrigation.~~

[Ed. note: The following paragraph moved to the beginning of the Approved Uses section.]

~~Do not apply sprays containing Roundup over the top of Christmas tree plantings.~~

~~Extreme care must be exercised to avoid contact of spray containing Roundup with foliage and stems of Christmas trees or severe damage or death may result.~~

Noncropland Areas and Industrial Sites**Noncropland Areas – Tank Mix Combinations**

Tank mix combinations of Surflan A.S. plus Roundup and many other labeled herbicides may be used to control undesirable vegetation in noncropland areas such as roadsides, rights-of-way, etc. When applied according to use directions, these tank mixes will provide control of susceptible weed species listed on the respective product labels. Refer to tank mix product labels for specific use directions, precautions and limitations before use.

Broadcast Application Rates

Length of Control	Surflan A.S.	
	quarts/acre	fl oz/1000 sq ft
2 to 4 months	2	1.5
4 to 8 months	4	3
8 to 12 months	6	4.5

Industrial Sites – Tank Mix Combinations

Tank mix combinations of Surflan A.S. plus Roundup, Spike and many other labeled herbicides may be used as overtop sprays to control existing vegetation on industrial sites such as utility substations, highway guard rails, sign posts and delineators. When applied according to use directions, these tank mixes will provide control of susceptible weed species listed on the respective product labels. Refer to tank mix product labels for specific use directions, precautions and limitation before use.

Warm Season Turfgrasses

Surflan A.S. may be applied as a preemergence treatment for control of annual grasses and certain broadleaf weeds in established warm season turf including bahiagrass, bermudagrass, buffalograss, centipedegrass, St. Augustinegrass, and zoysiagrass and or established tall fescue growing in warm season areas. Established turf is defined as a dense turf having a well-anchored root system and healthy, vigorous top growth. Use Surflan A.S. only as a part of a total turf management program that includes good fertilization practices.

Surflan A.S. may be tank mixed with Gallery[®] herbicide (California registration pending) and applied preemergence to broaden the spectrum of broadleaf weed control in warm season turf. Refer to the label for Gallery label for specific use directions, precautions, and limitations before use.

Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of Surflan A.S.

Surflan A.S. will not control emerged weeds. Successful preemergence control of weeds listed on this label requires that Surflan A.S. be applied prior to weed germination and be activated by at least one-half (1/2) inch of rainfall or irrigation within 21 days of application.

Special Use Precautions:

To avoid possible injury, do not apply Surflan A.S. to:

- Cool season turfgrass species.
- Golf course putting greens or tees or lawns containing dichondra or cool season turfgrass species.
- Newly sprigged or sodded areas of bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until these turfs are well established and have well anchored root systems.
- Newly hydromulched areas of bermudagrass until such areas are well established.
- Bermudagrass variety "Sun-Turf" when tank mixed with atrazine.

~~Surflan A.S. will not control emerged weeds.~~

~~Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of Surflan A.S.~~

Surflan A.S. may injure turf that is not well-established or is stressed or weakened due to unfavorable winter climatic conditions, drought, nematodes, or other factors which damage or weaken turf root systems. Apply Surflan A.S. only to healthy, well-established turf that has a well-anchored root system.

~~Use Surflan A.S. only as a part of a total turf management program that includes good fertilization practices.~~

Do not apply Surflan A.S. in the spring or early summer to tall fescue turfgrass reseeded the previous fall. In such cases, apply Balan[®] 2.5G granular herbicide at 60 to 80 pounds per acre in early summer (Round 1) and Surflan A.S. at 1.5 quarts per acre approximately eight weeks later (Round 2). Do not apply Surflan A.S. at the single application rate (2 quarts per acre) to established tall fescue; in such cases, apply 1.5 quarts per acre of Surflan A.S. in an initial application, followed by a second application of 1.5 quarts per acre 8 to 10 weeks later.

In bermudagrass areas that have been overseeded with winter grasses, a spring application of Surflan A.S. will thin the overseeded grasses.

Annual Grasses Controlled by Surlan A.S.

Summer Annuals:

Common Name	Scientific Name
barnyardgrass (watergrass)	<i>Echinochloa crus-galli</i>
crabgrass, large	<i>Digitaria sanguinalis</i>
crabgrass, smooth	<i>Digitaria ischaemum</i>
crabgrass	<i>Digitaria</i> spp.
crowfootgrass	<i>Dactyloctenium aegyptium</i>
foxtail, bristlegrass	<i>Setaria magna</i>
foxtail, giant	<i>Setaria faberi</i>
foxtail, green (pigeongrass)	<i>Setaria viridis</i>
foxtail, robust	<i>Setaria robusta</i>
foxtail, yellow	<i>Setaria glauca</i>
goosegrass (silver crabgrass)	<i>Eleusine indica</i>
Johnsongrass (seedling only)	<i>Sorghum halepense</i>
ryegrass, Italian	<i>Lolium multiflorum</i>
sandbur, field	<i>Cenchrus incertus</i>

Winter Annuals:

Common Name	Scientific Name
bluegrass, annual	<i>Poa annua</i>

Annual Broadleaf Weeds Controlled by Surlan A.S.

Summer Annuals:

Common Name	Scientific Name
carpetweed	<i>Mollugo verticillata</i>
knotweed, prostrate	<i>Polygonum aviculare</i>
purslane, common	<i>Portulaca oleracea</i>

Winter Annuals:

Common Name	Scientific Name
chickweed, common	<i>Stellaria media</i>
henbit	<i>Lamium amplexicaule</i>

Broadleaf Weeds Suppressed by Surlan A.S.

Common Name	Scientific Name
groundsel, common	<i>Senecio vulgaris</i>
spurge, prostrate	<i>Euphorbia humistrata</i>
woodsorrel, yellow	<i>Oxalis stricta</i>

Application Rates, Frequency and Timing of Application

Surflan A.S. can be applied in the spring for summer annual grass and broadleaf weed control, and in the fall for annual bluegrass (*Poa annua*) and winter annual broadleaf weed control.

1. Summer Annual Grasses and Broadleaf Weeds

Single Application Program: Apply 2 quarts per acre of Surflan A.S. per acre in late winter or early spring, prior to the onset of conditions favorable for annual weed germination.

Split Application Program: As an alternative to a single application program, Surflan A.S. may be applied in a split application. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time. Apply 1.5 quarts per acre of Surflan A.S. in an initial application, followed by a second application of 1.5 quarts per acre 8 to 10 weeks later.

The second treatment of the split application may follow application of a different preemergence grass herbicide in place of the initial application of Surflan A.S. application.

2. Annual Bluegrass (*Poa annua*) and Winter Annual Broadleaf Weeds

In areas of heavy annual bluegrass infestation, its elimination will result in temporary thinning of turfgrass cover. Proper fertilization, irrigation and soil incorporated reseeding should be employed to speed the restoration of desirable turfgrass cover in areas previously occupied by annual bluegrass (see section on reseeding).

Apply Surflan A.S. as a preemergence treatment in late summer or early fall, prior to the expected germination period for annual bluegrass and winter annual broadleaf weeds. If annual bluegrass infestation is severe and its elimination will result in thinning of turfgrass cover, apply Surflan A.S. at 1.5 quarts per acre. If thinning of turfgrass cover is not a potential problem, Surflan A.S. may be applied at 2 quarts per acre.

Broadcast Application Rates

Surflan A.S.	
quarts/acre	fl oz/1000 sq ft
1.5	1
2	1.5

Weed Control In Florida

In Florida, apply 1.5 quarts per acre of Surflan A.S. three times per year, or every 90 to 100 days, in the fall, early spring, and early summer. Do not apply more than 1.5 quarts per acre of Surflan A.S. in any single application.

Application Equipment

Apply Surflan A.S. evenly over the turfgrass area. Avoid spray pattern skips and overlaps that may result in incomplete coverage or over-application. For best results use application equipment designed to uniformly broadcast liquid herbicides. Calibrate application equipment prior to use, according to manufacturer's directions. Check equipment frequently to make sure it is working properly and distributing spray uniformly.

Reseeding

Herbicides that control annual weeds may also affect establishment of desirable turfgrass seedlings. Reseeding should be delayed for at least 90-120 days following application of Surflan A.S. With reseeding, it is essential that proper cultural practices such as soil cultivation and seedbed preparation,

irrigation and fertilization be followed. For satisfactory reseeding results following use of Surlan A.S. use, the seeding rate should be increased and equipment designed to place seed in full contact with soil (such as the Rogers Aero Seeder) should be employed.

[Ed. note: The following moved to this location from the beginning of this section.]

Special Use Precautions:

To avoid possible injury, do not apply Surlan A.S. to:

- Cool season turfgrass species.
- Golf course putting greens or tees or lawns containing dichondra or cool season turfgrass species.
- Newly sodded or sodded areas of bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until these turfgrasses are well-established and have well-anchored root systems.
- Newly hydromulched areas of bermudagrass until such areas are well-established.
- Bermudagrass variety "Sun Turf" when tank mixed with atrazine.

Warranty Disclaimer

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. DowElanco MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of DowElanco or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at DowElanco's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

DowElanco shall not be liable for losses or damages resulting from handling or use of this product unless DowElanco is promptly notified of such loss or damage in writing. In no case shall DowElanco be liable for consequential or incidental damages or losses.

The terms of the "Warranty Disclaimer" above and this "Limitation of Remedies" cannot be varied by any written or verbal statements or agreements. No employee or sales agent of DowElanco or the seller is authorized to vary or exceed the terms of the "Warranty Disclaimer" or this "Limitation of Remedies" in any manner.

*Trademark of DowElanco