

Supplemental Labeling



DowElanco

Quad IV, 9002 Purdue Road

P.O. Box 681428

Indianapolis, Indiana 46268-1189 USA

~~Treflan[®] E.C. (EPA Reg. No. 62710-03)~~
~~Treflan M.T.F. (EPA Reg. No. 62710-116)~~
 Treflan 5 (EPA Reg. No. 62719-118)
~~Treflan 80DG (EPA Reg. No. 62710-216)~~

Treflan Fall Applied Preplant Soil Incorporated for Foxtail (Pigeongrass) Control in Spring Seeded Wheat, Durum and Barley

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan before applying. Carefully follow use precautions and applicable use directions.
- Use of Treflan according to this supplemental labeling is subject to all precautions and limitations imposed by the label affixed to the container for Treflan.

Directions for Use

Apply Treflan herbicide in the fall for foxtail (pigeongrass) control during the following growing season. Incorporate one time within 24 hours. Incorporate a second time before planting to destroy existing weeds and insure a uniform distribution of Treflan in treated soil. Treflan may be applied to ground that has a manageable trash level, has been fallowed or pre-tilled.

Broadcast Application Rates per Acre:

Soil Texture	Treflan E.C. or Treflan M.T.F.	Treflan 5	Treflan 80DG
	(pints)	(pints)	(ounces)
Coarse and Medium	1	0.8	10
Fine	1.5	1.2	15

ACCEPTED
 24 FEB 1992
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 62719-118

Incorporation Directions

Any of the implements described below may be used for the first incorporation of Treflan. Use only the disc or field cultivator for the spring incorporation pass and incorporate in a different direction. Poor weed control may result if untreated soil is moved to the surface during the second incorporation pass. To avoid this problem, the second incorporation should not be deeper than the first.

1. **Chisel Plow:** May be used for the first incorporation pass only. Operate at 4 to 6 inches deep and at 4 to 6 mph. A chisel plow is defined as having 3 rows of 14 to 18 inch sweeps on no greater than 12 inch centers. Stagger sweeps so that no soil is left unturned. Use of chisel points may result in inadequate soil mixing.
2. **Tandem Disc:** Operate at 3 to 4 inches deep and at 4 to 6 mph.
3. **Field Cultivator:** Operate at 3 to 4 inches deep and at 5 or more mph. A field cultivator is defined as having 3 to 4 rows of sweeps with "c" or "s" shaped shanks on no greater than 7 inch centers. Stagger sweeps so that no soil is left unturned.

Planting Directions

Set equipment to place seed approximately 2 inches deep.

continued on back

Precautions

Carefully follow use directions to minimize potential crop stress. Under certain conditions, delayed crop emergence and or stand reduction may occur when Treflan E.C. is applied to wheat, durum or barley. The combined effect of certain cultural practices and unfavorable soil or environmental conditions may cause excessive crop seedling stress resulting in retarded crop growth, stand reduction and possibly reduced yield. For best results, observe the following cultural practices or precautions: ✓

- Use tillage methods that provide a uniformly firm seedbed and time tillage operations to conserve moisture.
- Do not exceed recommended application rates for Treflan. This is particularly important on coarse textured or low organic matter soils.
- Use only high quality seed where Treflan is to be applied (avoid use of small seed with low starch reserves).
- If seed treatments are used, apply at the correct rate and uniformly across all seeds. Misapplication may result in reduced germination and/or seedling vigor.
- Avoid use of seed varieties known to have poor seedling (emergence) vigor.
- Do not fall apply Treflan E.C. in combination with any other preplant incorporated herbicide.

Soil characteristics and environmental conditions which may contribute to crop seedling stress that may be accentuated by use of Treflan E.C. include:

- Soil related - High salinity, eroded knolls/hilltops, loose dry soils and compaction.
- Weather related - Cold and/or wet soils, excessively hot soils, excessive moisture, drought, and soil crusting from heavy rainfall.

Note: Do not apply Treflan on small grains where a dinitroaniline herbicide such as Treflan was applied at a rate recommended for row crops (oil seeds) during the previous growing season. ✓

*Trademark of DowElanco

T2EAGXXX Approved / / /
Replaces EA 2251

Amendment(s)

- 1) Label edited to reflect DowElanco transfer.
- 2) Revised use precautions to be consistent with Treflan product labels.