

01116

62719-121

log 13


(Container Label)

(logo) DowElanco

Spike* 20P

A surface applied herbicide for woody plant control in such areas as:

- Along highways
- Ditchbanks
- Fence rows
- Firebreaks
- Industrial sites
- Pastureland
- Pipelines
- Railroad rights-of-way
- Rangeland
- Utility rights-of-way
- Other noncropland areas for control of woody plant species
- Clearings for Wildlife Habitat



CAREFULLY READ THE ENTIRE LABEL, INCLUDING ALL USE PRECAUTIONS, BEFORE APPLYING TO AVOID INAPPROPRIATE USE OF THIS PRODUCT.

Active Ingredient:

tebuthiuron: N-[5-(1,1-dimethylethyl)-1,3,4-thiadiazol-2-yl]-N,N'-dimethyl urea	20.0%
Inert Ingredients	80.0%
Total	100.0%

Contains 0.2 pounds active ingredient per pound of product.

Keep Out of Reach of Children

CAUTION

See additional precautionary statements on side panel. See directions for use inside booklet.

NOT FOR RESIDENTIAL USE

NOTICE

Read the entire label before using. Use only according to label directions.

Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" elsewhere on this label.

IN CASE OF EMERGENCY endangering life or property involving this product, call collect 517-336-4400

AGRICULTURAL CHEMICAL --Keep away from food, feedstuffs and water supplies. Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-121
EPA Est. No. 00000-XX-00

*Trademark of DowElanco
DowElanco • Indianapolis, IN 46268, U.S.A.

Specialty Herbicide

(insert graphic of tree in circle with slash through it. Copy below the tree is: Spike* 20P will kill trees and shrubs. Carefully read the precautions before using.)

2-13

Precautionary Statements

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

CAUTION

PRECAUCION:

Precaucion al usuario: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

Causes Eye Irritation - Harmful If Swallowed, Inhaled Or Absorbed Through The Skin

Avoid breathing dust and contact with eyes, skin or clothing. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

First Aid

If in eyes: Flush eyes with plenty of water. Call a physician if irritation develops.

If swallowed: Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger or, if available, by administering one to two teaspoons of syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled: Remove individual to fresh air. If breathing difficulty occurs, get medical attention. If not breathing, provide cardiopulmonary resuscitation assistance (mouth-to-mouth) and get medical attention.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation develops.

Environmental Hazards

Do not apply Spike near desirable trees or other woody species. Exposure of even a small part of a plant root system to Spike may cause severe plant injury or death. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning equipment, disposing of equipment washwaters, or disposing of wastes. Do not apply to any portion of a ditchbank that will come into contact with water. Do not apply to ditches used to transport irrigation or potable water.

Spills: To prevent unintended damage to nontarget vegetation, carefully collect and dispose of spilled pellets, whether in storage areas, vehicles or on the soil surface.

Special Use Precautions:

Spike 20P is intended for control of unwanted woody vegetation such as trees, shrubs and vines. Spike 20P will also control herbaceous broadleaf plants such as clover or lespedeza. Grasses in the area immediately adjacent to pellets may be temporarily damaged. Dormant season application is recommended to minimize herbicidal effects on grasses and other herbaceous plants. The herbicidal activity of Spike in soil may prevent the growth of trees, shrubs and other broadleaf vegetation for several years after treatment.

Safe use of Spike 20P requires the following guidelines to be carefully followed:

Treatment Setback

Do not apply Spike 20P in the vicinity of desirable plants. Exposure of even a small part of a plant root system to Spike may cause severe plant injury or death.

Note: Plant roots usually occupy an area much larger than the aerial portion of the plant. Treatment setback distance should be 1 to 2 times the height or width of adjacent nontarget vegetation, whichever is greater. For example, if adjacent nontarget vegetation is 25 feet tall, the treatment setback should be 25 to 50 feet.

If there is a question about appropriate setback distance, contact an Arboriculturist (tree expert) to determine if the proposed treatment area is free of roots of desirable vegetation.

Potential Product Movement:

Spike 20P or Spike-containing soil may be moved from treated areas by flowing water, wind or mechanical means. Do not apply Spike in areas where overland flow of water might move Spike 20P or Spike-containing soil from the treated area. Do not apply where wind erosion may cause movement of Spike-containing soil from the treated area unless the surface has been stabilized with a gravel mulch or some other means. Do not apply in areas where soil may be redistributed by mechanical means to non-treated areas.

Cleaning of Equipment:

Thoroughly clean all traces of Spike 20P from application equipment after use. Do not empty residues cleaned from application equipment on areas where contact with roots of desirable vegetation may occur.

Remedial Action After Accidental Application or Spill:

Take action to minimize the effects of an accidental application or spill immediately. Once rainfall has moved Spike into the plant root zone, the effect on woody plants is irreversible. Damage from accidental application or spill may be prevented only if soil containing Spike is carefully removed before rainfall has moved Spike into the root zone. Apply a waterproof covering to the affected area until cleanup is accomplished. Carefully collect Spike pellets and/or Spike-containing soil with appropriate equipment and dispose at an approved landfill site. If rainfall has occurred, remove surface soil in the affected area to the depth of Spike penetration.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: The herbicidal properties of Spike 20P require caution in handling, storage and transportation of this product. Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Open dumping is prohibited. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Do not reuse. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Spike 20P Container Label
Revised 07/30/91

404 13

(Label Booklet)

(logo) DowElanco

Spike* 20P

A surface applied herbicide for woody plant control in such areas as:

- Along highways
- Ditchbanks
- Fence rows
- Firebreaks
- Industrial sites
- Pastureland
- Pipelines
- Railroad rights-of-way
- Rangeland
- Utility rights-of-way
- Other noncropland areas for control of woody Plant species
- Clearings for Wildlife Habitat

CAREFULLY READ THE ENTIRE LABEL, INCLUDING ALL USE PRECAUTIONS, BEFORE APPLYING TO AVOID INAPPROPRIATE USE OF THIS PRODUCT.

Active Ingredient:

telcuthiuron: N-[5-(1,1-dimethylethyl)-1,3,4-thiadiazol-2-yl]-N,N'-dimethylurea	20.0%
Inert Ingredients	80.0%
Total	100.0%

Contains 0.2 pounds active ingredient per pound of product.

Keep Out of Reach of Children

CAUTION

See additional precautionary statements on side panel. See directions for use inside booklet.

NOT FOR RESIDENTIAL USE

NOTICE

Read the entire label before using. Use only according to label directions.

Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" elsewhere on this label.

IN CASE OF EMERGENCY endangering life or property involving this product, call collect 517-636-4400

AGRICULTURAL CHEMICAL --Keep away from food, feedstuffs and water supplies. Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-121
EPA Est. No. 00000-XX-00

*Trademark of DowElanco
DowElanco • Indianapolis, IN 46268, U.S.A.

Specialty Herbicide

(insert graphic of tree in circle with slash through it. Copy below the tree is: Spike* 20P will kill trees and shrubs. Carefully read the precautions before using.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

CAUTION

PRECAUCION:

Precaucion al usuario: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

Causes Eye Irritation • Harmful If Swallowed, Inhaled Or Absorbed Through The Skin

Avoid breathing dust and contact with eyes, skin or clothing. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

First Aid

If in eyes: Flush eyes with plenty of water. Call a physician if irritation develops.

If swallowed: Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger or, if available, by administering one to two teaspoons of syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled: Remove individual to fresh air. If breathing difficulty occurs, get medical attention. If not breathing, provide cardiopulmonary resuscitation assistance (mouth-to-mouth) and get medical attention.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation develops.

Environmental Hazards

Do not apply Spike near desirable trees or other woody species. Exposure of even a small part of a plant root system to Spike may cause severe plant injury or death. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning equipment, disposing of equipment washwaters, or disposing of wastes. Do not apply to any portion of a ditchbank that will come into contact with water. Do not apply to ditches used to transport irrigation or potable water.

Spills: To prevent unintended damage to nontarget vegetation, carefully collect and dispose of spilled pellets, whether in storage areas, vehicles or on the soil surface.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: The herbicidal properties of Spike 20P require caution in handling, storage and transportation of this product. Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Open dumping is prohibited. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Do not reuse. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

621

Special Use Precautions:

Spike 20P is intended for control of unwanted woody vegetation such as trees, shrubs and vines. Spike 20P will also control herbaceous broadleaf plants such as clover or lespedeza. Grasses in the area immediately adjacent to pellets may be temporarily damaged. Dormant season application is recommended to minimize herbicidal effects on grasses and other herbaceous plants. The herbicidal activity of Spike in soil may prevent the growth of trees, shrubs and other broadleaf vegetation for several years after treatment.

Safe use of Spike 20P requires the following guidelines to be carefully followed:

Treatment Setback:

Do not apply Spike 20P in the vicinity of desirable plants. Exposure of even a small part of a plant root system to Spike may cause severe plant injury or death.

Note: Plant roots usually occupy an area much larger than the aerial portion of the plant. Treatment setback distance should be 1 to 2 times the height or width of adjacent nontarget vegetation, whichever is greater. For example, if adjacent nontarget vegetation is 25 feet tall, the treatment setback should be 25 to 50 feet.

If there is a question about appropriate setback distance, contact an Arboriculturist (tree expert) to determine if the proposed treatment area is free of roots of desirable vegetation.

Potential Product Movement:

Spike 20P or Spike-containing soil may be moved from treated areas by flowing water, wind or mechanical means. Do not apply Spike in areas where overland flow of water might move Spike 20P or Spike-containing soil from the treated area. Do not apply where wind erosion may cause movement of Spike-containing soil from the treated area unless the surface has been stabilized with a gravel mulch or some other means. Do not apply in areas where soil may be redistributed by mechanical means to non-treated areas.

Cleaning of Equipment:

Thoroughly clean all traces of Spike 20P from application equipment after use. Do not empty residues cleaned from application equipment on areas where contact with roots of desirable vegetation may occur.

Remedial Action After Accidental Application or Spill:

Take action to minimize the effects of an accidental application or spill immediately. Once rainfall has moved Spike into the plant root zone, the effect on woody plants is irreversible. Damage from accidental application or spill may be prevented only if soil containing Spike is carefully removed before rainfall has moved Spike into the root zone. Apply a waterproof covering to the affected area until cleanup is accomplished. Carefully collect Spike pellets and/or Spike-containing soil with appropriate equipment and dispose at an approved landfill site. If rainfall has occurred, remove surface soil in the affected area to the depth of Spike penetration.

General Information

Spike 20P is a surface applied soil-active herbicide for control of woody plants (trees, shrubs and vines). Treatments become effective after sufficient rainfall has occurred to move the active ingredient in Spike 20P into the root zone. Herbicidal symptoms appear most rapidly when applied just before seasonal rainfall. Treated trees and shrubs (brush) exhibit leaf chlorosis and browning followed by defoliation. Woody plants may go through several defoliation cycles before death occurs. Time required to achieve control of woody vegetation may vary from one to several years. Increased application rates and additional time is required to achieve consistent woody plant control under the following conditions: (1) the treated area contains deep, medium-to-fine textured or high organic matter soils; (2) the target species are deep-rooted; or (3) the vegetation consists of Spike-tolerant species.

7.3.13

For best brush control results, do not disturb intact plants by such practices as wood cutting, chaining or burning for two years after Spike 20P application. Resprouting is more likely to occur if plants are disturbed before complete woody plant control occurs.

General Use Precautions

- **Maximum Application Rate for Grazing or Haying:** If treated area is to be used for grazing or haying, do not apply more than 10 pounds per acre of Spike 20P in areas receiving 20 inches or less average annual rainfall, or more than 20 pounds per acre of Spike 20P in areas receiving more than 20 inches average annual rainfall.
- **Haying restriction:** Do not cut hay for livestock feed for one year after a Spike treatment.
- **Effects on Herbaceous Vegetation:** Spike 20P may injure or suppress certain herbaceous vegetation in the treated area. Therefore, do not apply where such injury cannot be tolerated. Injury to most herbaceous perennial plants is reduced if Spike is applied when vegetation is dormant.
- **Ditchbank Usage:** Do not apply Spike 20P to any portion of a ditchbank that will come into contact with water as movement of Spike in water may result in injury or death to nontarget vegetation. Do not apply on ditches used to transport irrigation or potable water.
- Do not apply Spike 20P more than once per year.

Factors in Herbicidal Response

Soil Texture, Soil Depth and Organic Matter: Poor control or erratic results are likely to occur when Spike 20P is applied to soils containing more than 5% organic matter or more than 30% clay. Do not apply to "blackland" or other heavy clay soils which crack extensively upon drying. Other deep, medium and fine-textured soils supporting deep-rooted woody plant species require increased application rates for consistent control. The susceptibility of woody plants occurring on shallow, coarse or rocky soils with low organic matter is normally increased due to increased soil availability of Spike and shallow rooting depth. Therefore, use application rates at the low end of the rate range in these situations.

Woody Plant Size and Density: The height and density of woody vegetation is a reliable indicator of soil conditions. Woody vegetation is generally taller and more dense where soils are deeper and/or of medium to fine texture and where soil moisture conditions are more favorable. Higher rates in the rate range are required on such sites. On sites with coarse, shallow or rocky conditions with less favorable soil moisture conditions, woody vegetation will be smaller and less dense. Lower rates in the rate range are required for control on such sites. Where a high level of woody plant control is required and application rates cannot be adjusted for changes in soils, plant size or density, apply Spike at a rate sufficient to control the tallest and most dense woody vegetation in the treatment area.

Application Timing: Spike 20P may be applied anytime except when the soil is frozen or is saturated with moisture. For optimum results, applications should be made prior to the resumption of active seasonal growth in the spring or before expected seasonal rainfall. In areas receiving greater than 25 inches of annual rainfall, late summer and fall applications may require a higher application rate in the indicated rate range to achieve consistent control.

Spike 20P is recommended for control of brush regrowth after dozing or shredding, provided the regrowth has reached an average height of five feet or more prior to application. Spike works best when there is an abundance of active leaf area to stimulate water and herbicide uptake during the season following application. Taller regrowth will tend to provide faster and more consistent brush control.

Spike 20P may cause temporary herbicidal symptoms to appear on perennial grasses. Dormant season application is recommended to minimize herbicidal effects on desirable forage grasses.

Rangeland and Pasture Management Information

There are no label restrictions which require livestock grazing to be delayed following an application of Spike 20P. Spike 20P is a pelleted formulation and does not adhere to plants. Therefore, significant plant residues do not occur as a result of application. However, at the time of application forage species may be sparsely distributed and in a low state of vigor due to competition from woody plants. Under such circumstances, the density and vigor of forage species may be enhanced by deferment of grazing following application of Spike 20P.

Grazing Management

For optimum perennial forage grass response, desirable species should be present in the area to be treated at a minimum of 10% of normal plant density (density = plants per unit area) compared to similar rangeland or pasture sites not dominated by woody plants. To encourage forage grass response, grazing should be deferred during the entire active growing season following application. Poor vegetative vigor or inadequate rainfall, may necessitate additional grazing deferment during periods of active forage growth. Light grazing of mature forage after seed maturity will not harm grass recovery and can aid in seed dispersal. Forage grass production usually increases as woody plant competition for water and nutrients is reduced. However, increased forage production is also dependent on adequate rainfall and a sound grazing management program.

Precaution: The density of cool season grass stands such as fescue and crested wheatgrass may be reduced after Spike 20P application. Factors which may contribute to the possibility of stand reduction include excessive application rates, areas of shallow or rocky soil and low brush density.

Rangeland and Pasture Overseeding: Apply Spike 20P at recommended rates. Overseeding involving burning or chaining of treated brush should not be attempted for at least two growing seasons after application. Apply seed and fertilizer at recommended rates into ash as soon as possible after burning or just prior to chaining. Cool season grasses are normally seeded in early fall and warm season grasses in the spring after the expected frost-free date. Aerial seeding without burning or chaining may be attempted in the fall or spring following a Spike application, but natural seedbed conditions must be relied upon for seedling establishment. Consult local range management specialists for recommendations on locally adapted species, seeding time and grazing management.

Application of Spike 20P

Individual Plant Treatments

Individual plants, multistem clumps or small stands of woody vegetation may be hand treated. For individual plant treatments apply Spike 20P evenly over the area occupied by the target plant(s).

Pasture and Rangeland:

- In areas receiving 20 inches or less average annual rainfall: Apply Spike 20P at a rate of 3/8 oz. per 100 sq ft (equivalent broadcast rate = 10 lb/A).
- In areas receiving more than 20 inches average annual rainfall: Apply Spike 20P at a rate of 3/4 oz. per 100 sq ft (equivalent broadcast rate = 20 lb/A).

Noncropland:

- Apply Spike 20P at a rate of 3/8 oz. to 1 1/8 oz. per 100 sq ft (equivalent broadcast rate = 10 to 30 lb/A).

Broadcast Treatments

Spike 20P must be applied with ground or aerial application equipment capable of accurate calibration and able to provide a uniform distribution of pellets on the soil surface. Use of equipment not capable of confining the spread of pellets to the target area may result in injury or

death of vegetation outside the intended treatment area. Contact a DowElanco Sales Representative for recommendations on application equipment for different use situations.

Spike 20P may be applied by ground or air broadcast by/or under the supervision of U.S. Government Agencies.

Broadcast Application Rates:

Pasture and Rangeland: Do not apply more than 10 pounds per acre Spike 20P in areas receiving 20 inches or less average annual rainfall. Do not apply more than 20 pounds per acre in areas receiving 20 inches or more average annual rainfall.

Noncropland: Spike 20P may be applied at up to 30 pounds per acre on noncropland.

Woody Plant Controlled by Spike 20P

Rangeland, Pastureland and Noncropland:

Apply Spike 20P at 3.75 to 5 lb per acre on the following woody plant species:

Note: On rangeland and pastureland, apply 3.75 to 5 lb/A of Spike 20P where a higher degree of control is required (See "Factors in Herbicidal Response" in "General Information" section of this label). Spike 20P may be applied at rates as low as 2.5 per acre on sites with shallow, rocky and coarse textured soils having low organic matter content, or where partial control is desired.

<u>Common Name</u>	<u>Scientific Name</u>
Burroweed (Density less than 1/sq. ft.)	Haplopappus tenuisectus
Ceniza	Leucophyllum frutescens
Creosotebush	Larrea tridentata
Mimosa, Catclaw (Wait-a-minute-bush)	Mimosa pigra
Paloverde	Cercidium spp.
Sagebrush, Big	Artemisia tridentata
Sagebrush, Sand	Artemisia filifolia
Snakeweed, Broom (Density less than 1/sq. ft.)	Gutierrezia sarothrae
arbush	Flourensia cernua
Whitethorn	Acacia constricta

Rangeland, Pastureland and Noncropland:

Apply Spike 20P at 2.5 to 10.0 lb per acre on the following woody plant species:

<u>Common Name</u>	<u>Scientific Name</u>
Oak, Sand Shinnery*	Quercus havardii

***Note:** A wide rate range is provided to accommodate the broad range of soil and climatic variation which occurs in areas occupied by sand shinnery. Use the lowest application rate only on shallow sands in southern part of species range or where partial control is desired. Use a higher dose in indicated rate range for deeper sands and dunes, and on shinnery varieties with tall and dense growth habit which become more prevalent in the mid-to-northern part of the species range (See "Factors in Herbicidal Response" in "GENERAL INFORMATION" section of this label).

10 of 13

Rangeland, Pastureland and Noncropland:**Apply Spike 20P at 5 to 10 lb per acre on the following woody plant species:**

<u>Common Name</u>	<u>Scientific Name</u>
Oak, Bigelow* (Parital control)	Quercus durandi
Oak, Mohr* (Parital control)	Quercus mohriana
Oak, Running Live* (Parital control)	Quercus virginiana
Whitebrush	Aloysia lycoides
Wolfberry, Berlandier	Lycium berlanderi

* Note: Use a higher dosage in indicated rate range on tall and dense stands.

Rangeland, Pastureland and Noncropland:**Apply Spike 20P at 10 to 20 lb per acre on the following woody plant species:**

<u>Common Name</u>	<u>Scientific Name</u>
Acacia, Blackbrush	Acacia rigidula
Acacia, Catclaw	Acacia greggii
Acacia, Twisted	Acacia tortuosa
Alder, Speckled*	Alnus rugosa
Apple-of-Sodom	Solanum sodomaeum
Birch, Gray	Betula populifolia
Blueberry	Vaccinium spp.
Bluewood (Brazil)	Condalia obovata
Buckbrush	Symphoricarpos orbiculatus
Cherry, Bitter	Prunus emarginata
Cottonwood, Eastern*	Populus deltoides
Dogwood, Roughleaf	Cornus drummondii
Elm, American	Ulmus americana
Elm, Winged	Ulmus alata
Fir, Balsam*	Abies balsamea
Guajillo	Acacia berlanderi
Guava	Psidium guajava
Hackberry, Spiny (Granjeno)	Celtis pallida
Hackberry, Western	Celtis occidentalis
Hardhat *	Spiraea tomentosa
Hawthorn	Crataegus spp.
Huckleberry	Gaylussacia spp.
Huisache*	Acacia farnesiana
Koa haole	Leucaena leucocephala
Locust, Black	Robinia pseudoacacia
Manzanita	Arctostaphylos spp.
Maple, Sugar*	Acer saccharum
Mulberry, Red	Morus rubra
Oak, Black	Quercus velutina
Oak, Blackjack	Quercus marilandica
Oak, Blue	Quercus douglasii
Oak, Bur	Quercus macrocarpa
Oak, Post	Quercus stellata
Oak, Shrub live	Quercus turbinella
Oak, Southern red	Quercus falcata
Oak, White*	Quercus alba
Pinus*	Pinus spp.
Poplar, Balsam*	Populus balsamifera
Rose, McCartney*	Rosa bracteata
Rose, Multiflora	Rosa multiflora
Sage, black	Salvia melifera

Spruce, White*	<i>Picea glauca</i>
Sumac, Dwarf	<i>Rhus copallina</i>
Sumac, Littleleaf	<i>Rhus microphylla</i>
Sumac, Skunkbush	<i>Rhus trilobata</i>
Sumac, Smooth	<i>Rhus glabra</i>
Sumac, Staghorn	<i>Rhus typhina</i>
Tamarack*	<i>Larix laricina</i>
Thornapple, Desert	<i>Datura discolor</i>
Yaupon	<i>Ilex vomitoria</i>
Yaupon, Desert	<i>Schaefferia cummolinia</i>

* Note: Use a higher dosage in indicated rate range on all sites.

Noncropland only:

Apply Spike 20P at 20 to 30 lb per acre on the following woody plant species:

<u>Common Name</u>	<u>Scientific Name</u>
Alder, Red	<i>Alnus rubra</i>
Ash, Green*	<i>Fraxinus pennsylvanica</i>
Aspen, Bigtooth	<i>Populus grandidentata</i>
Beech, American	<i>Fagus grandifolia</i>
Blackberry	<i>Rubus spp.</i>
Blackberry, Cutleaf*	<i>Rubus laciniatus</i>
Boxelder	<i>Acer negundo</i>
Chamise	<i>Adenostoma fasciculatum</i>
Cherry, Black	<i>Prunus serotina</i>
Chokecherry, Common	<i>Prunus virginiana</i>
Colubrina, Texas	<i>Colubrina texensis</i>
Coyotebush*	<i>Baccharis pilularis</i>
Creeper, Virginia	<i>Parthenocissus quinquefolia</i>
Dogweed, Flowering	<i>Cornus florida</i>
Douglasfir	<i>Pseudotsuga menziesii</i>
Guayacan	<i>Portiera angustifolia</i>
Hickory, Bitternut	<i>Caraya cordiformis</i>
Hickory, Black	<i>Caraya texana</i>
Hickory, Pignut	<i>Caraya glabra</i>
Hickory, Shagbark	<i>Caraya uvata</i>
Huisache	<i>Acacia farnesiana</i>
Kidneywood, Texas	<i>Eysenhardtia texana</i>
Kudzu	<i>Pueraria lobata</i>
Lantana*	<i>Lantana camara</i>
Leatherstem	<i>Jatropha dioica</i>
Lotebust (Condalia)	<i>Ziziphus obtusifolia</i>
Maple, Bigleaf	<i>Acer macrophyllum</i>
Maple, Norway*	<i>Acer platanoides</i>
Maple, Silver*	<i>Acer saccharinum</i>
Melaleuca*	<i>Melaleuca quinquenervia</i>
Mountainmahogany, birchleaf	<i>Cercocarpus betuloides</i>
Oak, California Scrub	<i>Quercus dumosa</i>
Oak, Live	<i>Quercus virginiana</i>
Oak, Pin	<i>Quercus palustris</i>
Oak, Red	<i>Quercus rubra</i>
Oak, White	<i>Quercus alba</i>
Peppertree, Brazilian*	<i>Schinus terebinthifolius</i>
Pine	<i>Pinus spp.</i>
Raspberry, Black	<i>Rubus occidentalis</i>
Sala*	<i>Gaultheria shallon</i>
Sweetgum	<i>Liquidambar styraciflua</i>
Sycamore, American*	<i>Platanus occidentalis</i>
Trumpetcreeper	<i>Campsis radicans</i>
Tuliptree*	<i>Liriodendron tulipifera</i>

12813

Willow

Salix spp.

* Note: Use a higher dosage in indicated rate range on all sites.

Noncropland only:

Apply Spike 20P at 30 lb per acre on the following woody plant species:

<u>Common Name</u>	<u>Scientific Name</u>
Ash, White	<i>Fraxinus americana</i>
Ceanothus, Wedgeleaf	<i>Ceanothus cuneatus</i>
Chaparral, Whitethorn	<i>Ceanothus leucodermis</i>
Elm, Chinese	<i>Ulmus parvifolia</i>
Elm, Slippery	<i>Ulmus rubra</i>
Greenbriar, Roundleaf	<i>Smitax rotundifolia</i>
Hawthorn, Cockspur	<i>Crataegus crus-calli</i>
Manzanita, Greenleaf	<i>Arctostaphylos patula</i>
Maple, Vine	<i>Acer circinatum</i>
Privet	<i>Ligustrum spp.</i>
Redcedar, Eastern	<i>Juniperus virginiana</i>
Russianolive	<i>Eleagnus angustifolia</i>
Sumac, Laurel	<i>Rhus laurina</i>
Tallowtree, Chinese	<i>Sapium sebiferum</i>

WARRANTY DISCLAIMER

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. DowElanco makes no other express or implied warranty of merchantability or fitness for a particular purpose or any other express or implied warranty.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to the label instructions (including adverse conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of DowElanco or the seller. All such risks shall be assumed by the buyer.

LIMITATION OF REMEDIES

The exclusive remedy for losses or damages resulting from the use of this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at DowElanco's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

DowElanco shall not be liable for losses or damages resulting from handling or use of this product unless DowElanco is promptly notified of such loss or damage in writing. In no case shall DowElanco be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of DowElanco or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

DowElanco
 Quad IV, 9002 Purdue Road
 Indianapolis, IN 46268, U.S.A.

Spike 20P Booklet
Revised 07/30/91