

Supplemental Labeling



DowElanco

Quad IV, 9002 Purdue Road

P.O. Box 681428

Indianapolis, Indiana 46268-1428 USA

Treflan* TR-10* Herbicide

(EPA Reg. No. 62719-131)

(For Distribution and Use in the State of North Dakota)

Green Foxtail (Pigeongrass) Resistance to Treflan TR-10 and Other Dinitroaniline Herbicides Identified in the Spring Cereal Grain Production Area of North Dakota

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan TR-10 before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan TR-10 according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Treflan TR-10, except as described below.

Crop Production Precaution

Populations of green foxtail (pigeongrass) resistant to the dinitroaniline (DNA) class of herbicides have been identified in Bottineau, Mc Henry, Montrail, Ramsey, Renville, Rolette, Towner and Ward counties of North Dakota in fields which have a long history of dinitroaniline herbicide use. Treflan TR-10 will not control green foxtail which has developed DNA resistance. **Therefore, the grower assumes the risk of nonperformance due to DNA resistance if Treflan TR-10 is used to control green foxtail in spring cereal grain production areas of North Dakota.** Alternative green foxtail control practices should be utilized in these fields.

DowElanco strongly recommends utilizing the following management practices to prevent or delay the development or spread of DNA-resistant green foxtail in spring cereal production areas:

1. Rotate herbicides so that the same product or same class of herbicide is not used repeatedly year after year. Treflan TR-10 and/or other dinitroaniline herbicides should not be applied in consecutive years. Use other herbicides with different modes of action such as Tiller™, Hoelon™, Dakota™, or Stampede™ in alternate years.
2. Rotate crops and use alternative weed control methods, including tillage, fallow periods and/or other herbicides with different modes of action.
3. Thoroughly clean small grains harvested from fields with confirmed resistance before using as seed, or avoid using grain from DNA-resistant fields for seed.
4. Thoroughly clean all crop residues from tillage and harvesting equipment before moving out of fields with confirmed resistance.

For further information or consultation, contact your DowElanco representative.

*Trademark of DowElanco

122-33-057 Approved 04/23/92
Initial printing.

Amendment(s)

1) Warning about dinitroaniline resistant green
foxtail.

NOT REVIEWED

In Accordance with PR Notice 82-2.
Based on Draft Labeling Dated

15 MAY 1992