



Rubigan*

E.C.

Fungicide

A locally systemic fungicide for the control of scab, powdery mildew and rusts of apple; scab and powdery mildew of pear; leaf spot and powdery mildew of cherry; and powdery mildew of grape

Active Ingredient:

fenarimol: α -(2-chlorophenyl)- α -(4-chlorophenyl)-5-pyrimidinemethanol.....12.0%
 Inert Ingredients.....88.0%

Contains petroleum distillate.
 Contains 1 pound active ingredient per gallon.

EPA Reg. No. 62719-134
 E.P.A. Est. 5905-GA-01; 464-MI-1; 52379-MI-002
 Net Contents 1 qt

Keep Out of Reach of Children

WARNING AVISO:

Precaucion al usarlo: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

Precautionary Statements

Hazards to Humans and Domestic Animals

WARNING

Causes Substantial But Temporary Eye Injury • Causes Skin Irritation • Harmful If Swallowed, Inhaled Or Absorbed Through The Skin

Do not get in eyes, on skin or on clothing. Avoid breathing vapor or spray mist. Wear goggles, face shield or safety glasses, and protective

clothing such as coveralls, a long-sleeved shirt and impermeable gloves (i.e. butyl rubber) when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected workers. See additional farm worker safety requirements under the heading "Note to User - Reentry".

First Aid

If in eyes: Flush with plenty of water. Call a physician.

If swallowed: Drink promptly large quantities of milk, egg whites or gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Get medical attention.

If on skin: Immediately wash with plenty of soap and water. Get medical attention if irritation develops.

If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

Environmental Hazards

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply when weather conditions favor runoff or drift from the target area. Do not contaminate water when disposing of equipment washwaters.

Notice: Read the entire label before using. Use only according to label directions. Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" in attached label booklet.

In case of emergency endangering life or property involving this product, call collect (517) 636-4400.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

Rubigan* E.C.

ACCEPTED

NOV 02 1992

Under the Federal Insecticide, Fungicide, and Rodenticide Act, this product is registered under EPA Reg. No. 62719-134.

BEST AVAILABLE COPY

2015

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Note to User - Reentry

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your state Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. (Indicate specific crop warnings which inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to be taken in case of accidental exposure.) When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written warnings must include the following information:

WARNING. Area treated with Rubigan E. C. on (date of application). Do not enter without appropriate protective clothing until sprays have dried. In case of eye contact, immediately flush with plenty of water and call a physician. In case of skin contact, immediately wash with plenty of soap and water. Get medical attention if irritation develops. If inhaled and discomfort occurs, move to fresh air. Get medical attention if breathing difficulty occurs.

STORAGE AND DISPOSAL

Storage: Avoid freezing. Do not store near heat or open flame. Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on treatment site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General Information

Rubigan E.C. fungicide is an emulsifiable concentrate formulation designed to control various foliar and fruit diseases of apple, pear, cherry and grape listed in each respective crop section. Rubigan is a locally systemic fungicide that controls susceptible fungal diseases by preventing the formation of certain sterol components and other materials necessary for their growth. Rubigan is compatible with most insecticides, miticides and fungicides commonly recommended for the crops listed.

General Use Precautions

Do not graze cover crops treated with Rubigan or harvest forage for silage or hay.

Apply Rubigan with ground application equipment only.

Do not apply Rubigan through any type of irrigation system.

Observe rate restrictions and preharvest intervals outlined for each crop.

Mixing Directions

Fill the spray tank to at least one-half the required spray volume and start agitation. Add (in order) dry flowable, wettable powder, and flowable (aqueous suspension) products to the spray tank. Then add the recommended rate of Rubigan followed by other emulsifiable concentrate and solution products. Allow time for thorough mixing after adding each product. Finish filling spray tank and maintain continuous agitation in the spray tank during mixing, filling and spraying operations.

Use and Dosage Recommendations

Use Rubigan as a foliar spray to control the diseases listed in the apple, pear, cherry and grape crop sections. **Note:** Rubigan use recommendations for all crops are made according to geographic area: Eastern United States (East of the Rocky Mountain continental divide); and Western United States (West of the Rocky Mountain continental divide). Follow use recommendations in the appropriate section for your area.

Spray Volume and Concentration

Mix the recommended dosage of Rubigan (consult appropriate crop section for your area) in sufficient water to ensure adequate coverage and apply as a concentrate (less than 200 gallons per acre) or dilute (greater than 200 gallons per acre) spray using ground application equipment only. The final Rubigan tank mixture must have a minimum concentration of 3 fl oz (2 fl oz for grapes) of Rubigan per 100 gallons of water. Do not exceed the maximum use rate of Rubigan per acre per application for the respective crop. Attention should be given to sprayer speed and calibration, wind speed, and tree and vine foliar canopy to ensure adequate spray coverage with Rubigan.

Rate Ranges and Spray Intervals

Where a rate range is provided in use directions for Rubigan, use higher rates when moderate to severe infection conditions exist or are anticipated. Where a range in the spray interval is indicated (e.g., 7 to 10 days), use the shortest spray interval within the indicated spray interval range if moderate to severe infection conditions exist. Climatic conditions, varietal susceptibility, disease history and current disease status of the orchard or vineyard must be considered when choosing rates and spray schedules for Rubigan. Begin applications of Rubigan according to label directions or as recommended by your state agricultural experiment station or extension service specialist.

Tank Mixing

For control of each disease listed in the following tables Rubigan may be used alone or in combination (tankmix) with a protectant fungicide active against each respective disease. ~~Use the protectant fungicides at application rates recommended on their respective product labels.~~

Use the protectant fungicides at application rates and according to their respective product labels. **Note:** Tank mix combinations of Rubigan plus a copper-containing product may result in reduced effectiveness of Rubigan.

BEST AVAILABLE COPY

3045

Disease Control Recommendations for Rubigan:
 Eastern United States (East of the Rocky Mountain Continental Divide).

Crop Disease (Spray Schedule)	Rubigan ¹ Use Rate Per Acre (fl oz)	Maximum Spray Interval (days)	Use Directions	Use Restrictions
Apples and Pears Scab (Regular) (Extended ²) (Post-infection ³) 1st spray 2nd spray Powdery Mildew Rusts	8 - 12 (alone) 8 - 12 (tank mixed) 8 - 12 8 - 12 8 - 12 8 - 12	7 7 - 10 - 7 14 7 - 10	Begin applications at "green tip" or when environmental conditions become favorable for primary scab development and continue through the "second cover" spray. For "extended" spray schedule, tankmix with recommended rates of a protectant fungicide. Prior to switching from the "regular" spray schedule to a protectant fungicide (only) schedule, apply Rubigan in tankmix combination with recommended rates of a protectant fungicide. Apply "1st spray" within 96 hours from the beginning of the first "primary scab" infection period. Apply the "second spray" within 7 days of the "1st spray". Begin application at "tight cluster" and continue through the "cover sprays". Begin applications at "pink" and continue through the "second cover" spray.	Do not apply more than 12 fl oz of Rubigan per acre per application or more than 84 fl oz per acre per season. Do not apply within 30 days of harvest. <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;"> <p style="writing-mode: vertical-rl; transform: rotate(180deg);"> Note: Do not apply to pears until after petal fall. Such application(s) of Rubigan will only control scab resulting from infection periods occurring after petal fall. </p> </div>
Cherries Leaf Spot Powdery Mildew	6	7 - 10	Begin applications at "petal fall" and continue until "harvest". One or two "post-harvest" applications may be made within 45 days after "harvest". If an "alternate row" spray program is used, do not exceed a 7 day spray interval.	Do not apply more than 6 fl oz of Rubigan per acre per application or more than 36 fl oz per acre prior to harvest. Rubigan may be applied up to and after "harvest".
Grapes Powdery Mildew (Schedule) Pre-bloom Post-bloom 1st Summer spray 2nd Summer spray	2-3 4-5 5-6 5-6	14-18 14-18 14-18 14-18	Begin applications at 12 - 18 inches of shoot growth.	Do not apply more than 6 fl oz of Rubigan per acre per application or more than 19 fl oz per acre per season. Do not apply within 30 days of harvest.

¹ Rubigan may be applied alone or in tankmix combination with recommended rates of a protectant fungicide, insecticide and/or miticide. When used alone tankmix Rubigan with a surfactant, e.g.: R-88 Spreader Activator, Wilbur Ellis Co.; Bio-88 Spreader Activator, Kalo Laboratories; Regulard, Kalo Laboratories; Leafact 80A Activator, Purepro Co.; Induce, Helena Chemical Co.; and LI-700, Loveland Industries.
² When applied to dwarf or semi-dwarf trees using "tree row volume" calculations, do not apply less than 6 fl oz of Rubigan per acre in tank mix combination with recommended rates of a protectant fungicide.
³ Growers who do not have access to accurate on-farm scab infection period data cannot make effective use of the "post infection" spray schedule.

BEST AVAILABLE COPY

5015

Warranty Disclaimer

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. DOWELANCO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of DowElanco or the seller. All such risks shall be assumed by Buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at DowElanco's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

DowElanco shall not be liable for losses or damages resulting from handling or use of this product unless DowElanco is promptly notified of such loss or damage in writing. In no case shall DowElanco be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of DowElanco or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

* Trademark of DowElanco
DowElanco • Indianapolis, IN 46268

LABEL CODE ~~442-04-003~~
EPA APPROVAL ~~020402~~

DATE CODE ~~203~~
REPLACES ~~442-04-001~~

~~Revisions include:~~

- ~~1) Use directions placed in tabular format.~~
- ~~2) List recommended surfactants provided for use in Eastern U.S.~~
- ~~3) Use directions for grapes defined for Eastern and Western U.S.~~

Amendments:

- 1) Added precaution regarding tank mixes with copper-containing products.
- 2) Added recommendation to delay application of Rubigan until after petal fall on pears grown in California.

BEST AVAILABLE COPY