

112 35 37119 139

104/11

6/1/94

1

[Laminated Booklet -- Surflan AS, 2.5 gal]
[Front Cover of Booklet]

(logo) DowElanco

Surflan* Dry Flowable

A preemergence surface-applied herbicide for the control of most annual grasses and certain broadleaf weeds

Active Ingredient:

oryzalin: 3,5-dinitro-N ⁴ , N ⁴ -dipropylsulfanilamide.....	85.0%
Inert Ingredients	15.0%

Contains 0.85 pounds of active ingredient per pound of product.

Keep Out of Reach of Children

CAUTION

Agricultural Use Requirements
Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Personal Protective Equipment (PPE) and User Safety Recommendations, and Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. **Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.**

In case of emergency endangering health or the environment involving this product, call collect 517-636-4400.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.


EPA Reg. No. 62719-138

EPA Est. _____

*Trademark of DowElanco
DowElanco • Indianapolis, IN 46268 U.S.A.

Herbicide

Net Weight ___ lb



2/14

(next page of booklet):

Table of Contents	Page
Precautionary Statements	
Hazards to Humans and Domestic Animals	
Personal Protective Equipment	
User Safety Recommendations	
First Aid	
Environmental Hazards	
Directions for Use	
Agricultural Use Requirements	
Storage and Disposal	
General Information	
General Use Precautions	
Soil Preparation	
Activation and Cultivation	
Mixing Directions	
Surflan Dry Flowable - Alone	
Surflan Dry Flowable - Tank Mix Combinations	
Mixing Order	
Premixing	
Testing for Compatibility in Liquid Fertilizers	
Testing Procedure	
Compatibility Agents	
Application Methods	
Ground Broadcast Application	
Band Application	
Weeds and Grasses Controlled in Fruit and Nut Crops and Vineyards	
Approved Crops	
Fruit and Nut Crops and Vineyards (Non-bearing and Bearing)	
Surflan Dry Flowable - Alone	
Surflan Dry Flowable - Tank Mix Recommendations	
Recommended Surflan Dry Flowable Tank Mix Combinations	
Surflan Dry Flowable plus Goal plus Gramoxone Super	
Surflan Dry Flowable plus Gramoxone Super	
Surflan Dry Flowable plus Princep (simazine)	
Surflan Dry Flowable plus Roundup	
Surflan Dry Flowable plus Solicam	
Surflan Dry Flowable - Tank Mix Reference Table	
Warranty Disclaimer	
Inherent Risks of use	
Limitation of Remedies	

Precautionary Statements

Hazards to Humans and Domestic Animals**Keep Out of Reach of Children****CAUTION**

Causes Eye Irritation • Harmful If Swallowed, Inhaled Or Absorbed Through The Skin
Avoid breathing spray mist and contact with skin, eyes or clothing.

Personal Protective Equipment (PPE)**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If in eyes: Flush with plenty of water. Get medical attention if irritation develops.

If swallowed: Do not induce vomiting. Call a physician or Poison Control Center. If available, administer activated charcoal (6-8 heaping teaspoonfuls) with a large quantity of water. Do not give anything by mouth to an unconscious person. Immediately transport to a medical care facility and see a physician.

If inhaled: Remove individual to fresh air. Get medical attention if breathing difficulty occurs. If not breathing, give artificial respiration, preferably cardiopulmonary resuscitation and get medical attention immediately.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation develops.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Cover or incorporate spills.

BEST AVAILABLE COPY

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

BEST AVAILABLE COPY

General Information

Surflan Dry Flowable herbicide is a preemergence, surface applied herbicide for the control of most annual grasses and certain broadleaf weeds. Surflan Dry Flowable controls susceptible annual weeds by disrupting plant growth processes during germination. Surflan Dry Flowable may be applied in liquid sprays of water or liquid fertilizer. Surflan Dry Flowable may be tank mixed with other herbicides to control existing vegetation or improve the spectrum of weeds controlled. Surflan Dry Flowable alone does not control established weeds.

General Use Precautions

- Do not graze or feed forage from treated fields or orchards to livestock.
- Do not over-apply Surflan Dry Flowable. Over-application may result in crop injury or soil residue.
- Do not plant any root crop for 12 months following a Surflan Dry Flowable application.
- Do not use Surflan Dry Flowable on soils containing more than 5% organic matter.
- Apply Surflan Dry Flowable directly to the soil surface in orchards or vineyards.
- Carefully follow label directions to avoid poor weed control or crop injury.

Chemigation: Do not apply this product through any type of irrigation system.

Soil Preparation

Optimum herbicidal activity occurs when Surflan Dry Flowable is applied to the soil surface. Prior to application, existing unwanted vegetation should be destroyed by tillage or use of contact or translocated herbicides. Crop or weed residues, pruning and trash should be thoroughly mixed into the soil by tillage equipment or removed before treatment. Soil should be in good tilth and free of clods at the time of application.

Activation and Cultivation

A single 1/2 to 1 inch rainfall or sprinkler irrigation is required to activate Surflan Dry Flowable and move the herbicide into the zone of weed germination. Rainfall or irrigation of 1 inch or more is needed to activate Surflan Dry Flowable on fine-textured, high organic matter soils. If weeds begin to emerge, a shallow cultivation to a depth of 1 to 2 inches will destroy existing weeds and place Surflan Dry Flowable in the zone of weed germination.

Mixing Directions

BEST AVAILABLE COPY

Surflan Dry Flowable - Alone

Surflan Dry Flowable may be applied in water or most liquid fertilizer materials. Prior to mixing Surflan Dry Flowable in liquid fertilizer, refer to "Testing for Compatibility in Liquid Fertilizers" for test procedures to determine compatibility with the fertilizer product to be used. The combination of Surflan Dry Flowable with solution and suspension-type fertilizers provides annual weed control equal to Surflan Dry Flowable applied in water. Individual state regulations relating to liquid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company offering the fertilizer and chemical mixture for sale.

Start with a clean spray tank. Fill the sprayer to 1/3 to 1/2 of the required spray volume. Start agitation. Add the correct amount of Surflan Dry Flowable, continue agitation, and fill spray tank to required spray volume. Maintain continuous agitation from mixing through application.

Precaution: Do not allow the mixture to siphon back into the water source.

Surflan Dry Flowable - Tank Mix Combinations

Surflan Dry Flowable may be tank mixed with other products and applied with water or most liquid fertilizer materials. Prior to mixing Surflan and another product in liquid fertilizer, refer to the tank mix product manufacturer's label to determine if application in liquid fertilizer is recommended. Also refer to the section

60711

in this label entitled "Testing for Compatibility in Liquid Fertilizers" for testing procedures to determine tank-mix compatibility with the liquid fertilizer product to be used. The combination of Surflan with solution and suspension-type fertilizers provides annual weed control equal to Surflan applied in water. Individual state regulations relating to liquid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company offering the fertilizer and chemical mixture for sale. Read and follow all label instructions for each material to be added to the spray tank.

Vigorous and continuous agitation is required for all tank mixes. Sparger pipe agitators generally provide the best agitation in spray tanks. To prevent foaming, keep the end of the fill pipe below the surface of the water in the spray tank during filling to prevent air from being stirred or splashed into the mixture.

Mixing Order: Fill the spray tank to 3/4 of the total spray volume required. Start agitation. Add different formulation types in the order indicated below, allowing time for complete mixing and dispersion after addition of each product. Allow extra mixing and dispersion time for dry flowable products.

Add different formulation types in the following order: Surflan Dry Flowable and other dry flowables (DF); wettable powders (WP); flowables (F); liquids (L) or aqueous suspensions (S); solutions (S); and emulsifiable concentrates (EC).

Fill the spray tank to final volume. Maintain continuous agitation during mixing, final filling and throughout application. If spraying and agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be suspended before spraying is resumed. A sparger agitator is particularly useful for this purpose. Settled materials may be more difficult to resuspend than when originally mixed.

Premixing: When tank mixing, initial mixing and dispersion of Surflan Dry Flowable or other dry flowable or wettable powder products may be improved by premixing with water (slurrying). Where recommended, follow product label instructions for each material. Adding the slurried material to the spray tank through a 20 to 35 mesh wetting screen will help assure good initial dispersion. Line screens in the tank through a no finer than 50 mesh (100 mesh is finer than 50 mesh).

Testing for Compatibility in Liquid Fertilizers

Surflan Dry Flowable alone or in combination with other product formulations may not combine properly with some liquid fertilizer materials. Small quantities of such mixtures should always be tested before full-scale mixing. Follow the testing procedure below to determine if a compatibility agent is needed or which compatibility agent performs best in the mixture being applied. Nine compatibility agents listed at the end of this section have been thoroughly tested. Other surfactant materials available may not be formulated for use with liquid fertilizers.

Testing Procedure

1. Add 1 pint of liquid fertilizer to 1 quart glass jar.
2. Add 1 to 4 teaspoonfuls (depending on application rate) of the products to be tank mixed in the liquid fertilizer. Add the materials in the order recommended in the "Mixing Instructions" section of this label. Mix until evenly dispersed after addition of each formulation. If dry flowable or wettable powder formulations do not disperse well, it may be necessary to slurry the materials in a small amount of water before addition to the liquid fertilizer.
3. Observe the jar for about 10 minutes. If materials rise to the surface and form a thick layer (oil curds) which will not redispense when agitated, a compatibility agent is needed. If the mixture is easily redispersed with slight agitation, a compatibility agent is not required, however, good agitation must be provided to maintain dispersion in the spray tank from mixing through application.
4. If the need for a compatibility agent is demonstrated in step 3, it may be tested as follows: Using a clean clear plastic or glass container, start at step 1 above and add 1/2 teaspoon of the compatibility agent to the liquid fertilizer mixture. Shake well and then repeat steps 2 and 3.

BEST AVAILABLE COPY

An effective compatibility agent will cause the mixture to remain uniformly mixed with little or no separation for 1/2 hour or longer. If slight separation occurs, 2 to 3 inversions of container should be sufficient to uniformly redisperse the mixture. If oily curds form which will not disperse, try adding additional compatibility agent or use an alternative compatibility agent to achieve a uniform mixture.

A clean jar should be used for each test.

Compatibility Agents

The phosphate ester-type surfactants listed below are designed for use with liquid fertilizers and can be mixed at rates as low as 1 1/2 to 2 pints per ton of liquid fertilizer. They usually do not work well as compatibility agents for tank mixes in plain water. Add the compatibility agent just before adding herbicides. Read and follow label directions for the compatibility agent.

1. Sponto 168D (Witco Chemicals Co., Chicago, IL)
2. Compat (Farm Chemicals, Inc., Aberdeen, NC)
Not registered for use in California.
3. Unite (Hopkins Ag Chemical, Madison, WI)
4. T-Mulz 734-2 (Thompson-Hayward Chemical Co., Kansas City, KS)
Not registered for use in California.
5. Rigo Compatibility Agent (Rigo Company, Buckner, KY)
Not registered for use in California.
6. Amoco Spray Mate (Amoco Oil Co., Chicago, IL)
Not registered for use in California.
7. Kem-Link (Universal Coop, Minneapolis, MN)
Not registered for use in California.
8. Blendex (Helena Chemical, Cayce-West, Columbia, SC)
Not registered for use in California.
9. Spray-Aide (Miller Chemical and Fertilizer Corp., Hanover, PA)

Application Methods

Ground Broadcast Application

Apply Surlan Dry Flowable in a total spray volume of 20 to 40 gallons per acre (broadcast basis), using any properly calibrated low pressure herbicide sprayer that will apply the spray uniformly. Use herbicide nozzle tips and screens no finer than 50 mesh for nozzle and in-line strainers. As the amount of spray volume per acre decreases, the importance of accurate calibration and uniform application increases. Check the sprayer daily to insure proper calibration and uniform application. Avoid boom overlaps that will increase rates above those recommended.

BEST AVAILABLE COPY

Band Application

For band application, use the following formula to calculate the required amount of product per acre

Band Width (inches)

Row Width (inches)

X Broadcast rate per acre = Amount required per acre

Weeds and Grasses Controlled in Fruit and Nut Crops and Vineyards

Annual Grasses**Common Name****Scientific Name**

barley, little

Hordeum pusillum

barnyardgrass (watergrass)

Echinochloa crus-galli

bluegrass, annual (poa)

Poa annua

brachiaria (signalgrass)

Brachiaria spp.

browntop panicum

Panicum fasciculatum

crabgrass

Digitaria spp.

(large crabgrass)

(smooth crabgrass)

crowfootgrass

Dactyloctenium aegyptium

cupgrass

Eriochloa gracilis

downy brome

Bromus tectorum

fall panicum

Panicum dichotomiflorum

(spreading panicgrass)

foxtails

Setaria spp.

(bottlegrass)

(bristlegrass)

(giant foxtail)

(green foxtail)

(pigeongrass)

(robust foxtail)

(yellow foxtail)

guineagrass

Panicum maximum

(narrowleaf panicum)

johnsongrass (seedling only)

Sorghum halepense

junglerice

Echinochloa colonum

lovegrass, Mexican

Eragrostis mexicana

lovegrass, orcutt

Eragrostis orcuttiana

oat, wild

Avena fatua

ryegrass, annual (Italian)

Lolium multiflorum

sandbur, field

Cenchrus incertus

sprangletop, red

Leptochloa filiformis

Texas panicum

Panicum texanum

(buffalograss)

(Coloradograss)

witchgrass

Panicum capillare

BEST AVAILABLE COPY

Annual Broadleaf Weeds**Common Name**

carpet weed
 chickweed
 cudweed
 fiddleneck, coast
 Florida pusley
 (Florida purslane)
 (Mexican clover)
 (pusley)
 henbit
 knotweed, prostrate
 lambsquarters
 pigweeds
 (carelessweed)
 (prostrate pigweed)
 (redroot pigweed)
 (rough pigweed)
 (smooth pigweed)
 (spiny pigweed)
 (tumble pigweed)
 puncturevine
 purslane, common
 rockpurslane, redmaids
 shepherdspurse

Scientific Name

Mollugo verticillata
Stellaria media
Gnaphalium chilense
Amsinckia intermedia
Richardia scabra

Lamium amplexicaule
Polygonum aviculare
Chenopodium album
Amaranthus spp.

Tribulus terrestris
Portulaca oleracea
Calandrinia caulescens
Capsella bursa-pastoris

Surflan Dry Flowable provides partial control or suppression of:**Common Name**

filaree, redstem
 filaree, whitestem
 groundsel, common
 ladythumb
 lettuce, prickly
 mallow, common
 milkweed, climbing
 morningglory, annual
 mustard, black
 mustard, wild
 nightshade, black
 prickly sida (teaweed)
 ragweed, common
 ragweed, giant
 rocket, London
 smartweed, annual
 sowthistle, annual
 spurge, prostrate
 spurge, spotted
 velvetleaf

Scientific Name

Erodium cicutarium
Erodium moschatum
Senecio vulgaris
Polygonum persicaria
Lactuca serriola
Malva neglecta
Sarcostemma cynanchoides
Ipomoea spp.
Brassica nigra
Sinapis arvensis
Solanum nigrum
Sida spinosa
Ambrosia artemisiifolia
Ambrosia trifida
Sisymbrium irio
Polygonum spp.
Sonchus oleraceus
Euphorbia humistrata
Euphorbia maculata
Abutilon theophrasti

BEST AVAILABLE COPY

10244

Approved Crops

Fruit and Nut Crops and Vineyards (Non-bearing and Bearing)

Surflan Dry Flowable - Alone

Apply Surflan Dry Flowable to fruit and nut crops and vineyards as a preemergence treatment to control annual grasses and broadleaf weeds listed in "General Information". Surflan Dry Flowable is approved for use on the following crops:

almond	grapefruit	pear
apple	kiwi	pecan
apricot	lemon	pistachio
avocado	macadamia nut	plum
cherry	nectarine	pomegranate
fig	olive	prune
filbert	orange	walnut, English
grape	peach	

Small Fruits:

blackberry	currant	gooseberry
blueberry†	dewberry	loganberry
boysenberry	elderberry	raspberry

†Do not apply Surflan Dry Flowable to lowbush blueberries.

General Use Directions

Existing vegetation should be destroyed by tillage or use of a contact or translocated herbicide in tank mix combination with Surflan Dry Flowable. To insure tank mix compatibility of Surflan Dry Flowable with contact or translocated herbicides, mix the products together in correct proportions in a small container of water. After mixing, let stand for 5 minutes and check for compatibility. If the combination remains mixed or can be resuspended easily, the mixture is compatible.

Application Methods: Surflan Dry Flowable may be applied in liquid sprays of water or liquid fertilizer. Refer to the "General Information" section of this label for mixing directions and details on application methods and precautions before use.

Activation: A single 1/2 to 1 inch rainfall or sprinkler irrigation is required to activate Surflan Dry Flowable and move the herbicide into the zone of weed seed germination. If weeds begin to emerge, a shallow cultivation 1 to 2 inches deep will destroy existing weeds and place Surflan Dry Flowable in the zone of weed seed germination. Rainfall or sprinkler irrigation of 1 inch or more is required to activate Surflan Dry Flowable on fine-textured or high organic matter soil.

Broadcast Application Rates

Soil Texture	Length of Control	Surflan Dry Flowable (pounds/acre)
All Soil Textures	Short Term (2 - 4 months)	2.4
	Long Term (6 - 8 months)	4.7
	(8 - 12 months)	7.1

Precautions:

poor weed control may result if directions are not carefully followed. Over-application may result in crop injury or excessive soil residue. Apply spray directly to the soil surface of the orchard or vineyard.

BEST AVAILABLE COPY

Surflan Dry Flowable Tank Mix Recommendations

The following products are compatible with Surflan Dry Flowable when used in tank mix combination. Performance and risk of carryover from products listed below when used in combination with Surflan Dry Flowable at recommended rates is the same as when each product is used separately.

Follow tank mixing instructions provided in the "General Information" section of this label when mixing Surflan Dry Flowable with other products.

Users should always consult the manufacturer's label for the product(s) to be tank mixed with Surflan Dry Flowable for specific information on use rates, additional weeds controlled, rotational crop restrictions or risk of carryover, special tank mix instructions, additional use directions, precautions and limitations before use.

Note: Some crop uses of tank mix products are limited to or excluded from certain states.

Recommended Surflan Dry Flowable Tank Mix Combinations:

Surflan Dry Flowable plus Goal
 Surflan Dry Flowable plus Goal plus Gramoxone Super
 Surflan Dry Flowable plus Gramoxone Super
 Surflan Dry Flowable plus Princep (simazine)
 Surflan Dry Flowable plus Roundup
 Surflan Dry Flowable plus Solicam

Surflan Dry Flowable - Tank Mix Reference Table:

The following table lists approved crops for Surflan Dry Flowable and products for tank mixes with Surflan Dry Flowable

(* = Approved Crop). Refer to the table to determine if the herbicides recommended for tank mixing can be applied to your crop. For use on a given crop, two asterisks must be indicated for Surflan Dry Flowable and the product with which it is to be tank mixed.

	Surflan Dry Flowable	Goal	Gramoxone Super	Princep (simazine)	Roundup	Solicam
Almond	**	**	**	**	**	**
Apple	**	**	**	**	**	**
Apricot	**	**	**	-	**	**
Avocado	**	**	**	**	**	-
Cherry	**	**	**	**	**	-
Fig	**	**	**	-	**	-
Filbert	**	**	**	**	**	**
Grape	**	**	**	**	**	**
Grapefruit	**	-	**	**	**	**
Kiwi	**	**	**	-	**	-
Lemon	**	-	**	**	**	**
Macadamia nut	**	**	**	**	**	-
Nectarine	**	**	**	-	**	**
Olive	**	**	**	**	**	-
Orange	**	-	**	**	**	**
Peach	**	**	**	**	**	**
Pear	**	**	**	**	**	**
Pecan	**	**	**	**	**	**
Pistachio	**	**	**	-	**	-
Plum	**	**	**	**	**	**
Pomegranate	**	**	**	-	**	-
Prune	**	**	**	-	**	**
Walnut, English	**	**	**	-	**	**

BEST AVAILABLE COPY

120711

Small Fruits:

Blackberry	..	-
Blueberry	..	-
Boysenberry	..	-	-
Currants	..	-	-	-	..	-
Dewberry	..	-	-	-	..	-
Elderberry	..	-	-	-	..	-
Gooseberry	..	-	-	-	..	-
Loganberry	..	-	-	-
Raspberry	..	-

Warranty Disclaimer

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. DOWELANCO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of DowElanco or the seller. All such risks shall be assumed by Buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at DowElanco's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

DowElanco shall not be liable for losses or damages resulting from handling or use of this product unless DowElanco is promptly notified of such loss or damage in writing. In no case shall DowElanco be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of DowElanco or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

©Copyright 1992 by DowElanco

BEST AVAILABLE COPY

[Base Label of Laminated Booklet]

(logo) DowElanco

Surflan* Dry Flowable

A preemergence surface-applied herbicide for the control of most annual grasses and certain broadleaf weeds

oryzalin: 3,5-dinitro-*N,N*-dipropylsulfanilamide85.0%
 Inert Ingredients15.0%
 Contains 0.85 pounds of active ingredient per pound of product.

Precautionary Statements

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

CAUTION

Causes Eye Irritation • Harmful If Swallowed, Inhaled Or Absorbed Through The Skin
 Avoid breathing spray mist and contact with skin, eyes or clothing.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If in eyes: Flush with plenty of water. Get medical attention if irritation develops.

If swallowed: Do not induce vomiting. Call a physician or Poison Control Center. If available, administer activated charcoal (6-8 heaping teaspoonfuls) with a large quantity of water. Do not give anything by mouth to an unconscious person. Immediately transport to a medical care facility and see a physician.

If inhaled: Remove individual to fresh air. Get medical attention if breathing difficulty occurs. If not breathing, give artificial respiration, preferably cardiopulmonary resuscitation and get medical attention immediately.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation develops.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Cover or incorporate spills.

BEST AVAILABLE COPY

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to label booklet for additional precautionary information including Personal Protective Equipment (PPE) and User Safety Recommendations, and Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.

In case of emergency endangering health or the environment involving this product, call collect 517-636-4400.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-112

EPA Est. _____

*Trademark of DowElanco
DowElanco • Indianapolis, IN 46268 U.S.A.

Herbicide

Net Weight ___ lb

BEST AVAILABLE COPY