

JUN 1992

Dr. Marilyn Jones Ph.D.  
DOW 18000  
5002 Purdue Road  
Quad IV  
Indianapolis IN 46206

Dear Dr. Marilyn Jones:

Subject: Dow, Lactosulfone 75 Dry Flowable  
EPA Registration No. 148-100-01  
Re: Label Amendment  
Your submission dated 1/23/92.

The labeling referred to above submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is acceptable provided that under the Environmental Hazard Section of the label replace the statement "do not apply directly to water of wetlands" to read "do not apply directly to water, to areas where surface water is present or to subterranean water below the mean high water mark." Please submit five (5) copies of your final printed labels incorporating this amendment. A stamped copy of labeling is being sent for your records.

Sincerely,

Jeanne L. Butler  
Product Manager (20)  
Fungicide-herbicide Branch  
Registration Division (H-75660)

enclosure

**BEST AVAILABLE COPY**

CONCURRENCES							
SYMBOL	H-7505C						
SURNAME	Metcalf						
DATE	6/17/92						

2 9 22

[Container Label Front Panel]

(logo) DowElanco

# Gallery\*

## 75 Dry Flowable

A selective preemergence herbicide for control of certain broadleaf weeds in:

- Established turf
- Landscape ornamentals
- Container grown ornamentals
- Nursery stock
- Ground covers
- Ornamental bulbs
- Non-bearing fruit and nut trees and non-bearing vineyards
- Christmas tree plantations
- Non-cropland

**Active Ingredient:**

isoxaben: N-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers .....	75%
<b>Inert Ingredients</b> .....	25%
<b>Total</b> .....	100%

Contains 0.75 pounds active ingredient per pound.

**Keep Out of Reach of Children**

### CAUTION

Refer to back panel of container for precautionary information and Directions for Use including STORAGE AND DISPOSAL.

**Notice:** Read the entire label. Use only according to label directions. Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.

In case of emergency endangering health or the environment involving this product, call collect 517-636-4400.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-145  
20116

EPA Est. 464-MI-1  
900-000987

\*Trademark of DowElanco  
DowElanco • Indianapolis, IN 46208, U.S.A.

**Specialty Herbicide**

**Net Weight 1 lb**

REGISTERED  
EPA ESTABLISHED  
5/1/92

62719-145

(Back Panel of Container Label)

---

**Precautionary Statements**

---

**Hazards to Humans and Domestic Animals****CAUTION PRECAUCION:**

**Precaucion al usuario:** Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

**Causes Eye Irritation • Harmful If Inhaled**

**Avoid ingestion, breathing dust or spray mist, and contact with skin, eyes, or clothing. Use eye protection when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.**

**First Aid**

**If in eyes:** Flush eyes with plenty of water. Call a physician if irritation develops.

**If swallowed:** Call a physician or Poison Control Center.

**If on skin:** Wash with plenty of soap and water. Get medical attention if irritation develops.

**If inhaled:** Remove individual to fresh air. If not breathing, provide cardiopulmonary resuscitation assistance and get medical attention.

**Environmental Hazards**

Do not contaminate water when disposing of equipment washwaters. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift may result in reduced germination or emergence of non-target plants adjacent to treated area.

---

**Directions for Use**

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

**Chemigation:** Do not apply Gallery 75 Dry Flowable through any type of irrigation system.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**Storage:** Store in original container. Do not store in direct sunlight. Do not store at temperatures above 120°F. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Dispose of in a sanitary landfill, or incineration, or by burning if allowed by state and local authorities. If burned, stay out of smoke.

20116

\*Trademark of DowElanco

900-000988

Gallery 75 Dry Flowable

(Label Booklet)

(logo) **DowElanco**

# Gallery\*

## 75 Dry Flowable

A selective preemergence herbicide for control of certain broadleaf weeds in:

- Established turf
- Landscape ornamentals
- Container grown ornamentals
- Nursery stock
- Ground covers
- Ornamental bulbs
- Non-bearing fruit and nut trees and non-bearing vineyards
- Christmas tree plantations
- Non-cropland

**Active Ingredient:**

isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers .....	75%
<b>Inert Ingredients</b> .....	25%
<b>Total</b> .....	100%

Contains 0.75 pounds active ingredient per pound.

EPA Reg. No. 62719-145

**NOTICE:** The scoop provided may be used to measure Gallery 75 Dry Flowable herbicide for smaller treatment areas. One level scoopful is equivalent to approximately 1/2 oz and will treat a 1,000 sq ft area. Please refer to the label booklet for complete use directions and precautions.

\*Trademark of DowElanco

[Second Page of Booklet]

<b>Table of Contents</b>	<b>Page</b>
General Information	5
Mixing and Application Directions	6
Gallery 75 Dry Flowable - Alone	6
Gallery 75 Dry Flowable in Tank Mixes	6
Application	6
Weeds Controlled or Suppressed by Gallery 75 Dry Flowable	6
Approved Uses:	
Established Turfgrasses	9
Special Use Precautions	9
Landscape Ornamentals, Container Grown Ornamentals, Nursery Stock, Ground Covers, Ornamental Bulbs, Non-bearing Fruit and Nut Trees and Non-bearing Vineyards, and Christmas Tree Plantations	10
Special Use Precautions	10
Trees	10
Ornamental Shrubs	13
Ground Covers	23
Ornamental Bulbs	24
Christmas Tree Plantations	25
Non-cropland Areas	25
Warranty Disclaimer	26
Inherent Risks of Use	26
Limitation of Remedies	26

---

## Precautionary Statements

---

### Hazards to Humans and Domestic Animals

#### CAUTION PRECAUCION:

**Precaucion al usuario:** Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

#### Causes Eye Irritation - Harmful If Inhaled

**Avoid ingestion, breathing dust or spray mist, and contact with skin, eyes, or clothing. Use eye protection when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.**

#### First Aid

**If in eyes:** Flush eyes with plenty of water. Call a physician if irritation develops.

**If swallowed:** Call a physician or Poison Control Center.

**If on skin:** Wash with plenty of soap and water. Get medical attention if irritation develops.

**If inhaled:** Remove individual to fresh air. If not breathing, provide cardiopulmonary resuscitation assistance and get medical attention.

#### Environmental Hazards

Do not contaminate water when disposing of equipment washwaters. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift may result in reduced germination or emergence of non-target plants adjacent to treated area.

---

## Directions for Use

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

**Chemigation:** Do not apply Gallery 75 Dry Flowable through any type of irrigation system.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Storage:** Store in original container. Do not store in direct sunlight. Do not store at temperatures above 120°F. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Dispose of in a sanitary landfill, or incineration, or by burning if allowed by state and local authorities. If burned, stay out of smoke.

## General Information

Gallery 75 Dry Flowable herbicide is a preemergence herbicide for control of certain broadleaf weeds in established turf, landscape ornamentals, container grown ornamentals, nursery stock, ground covers, ornamental bulbs, non-bearing fruit and nut trees and non-bearing vineyards, Christmas tree plantations and non-cropland areas. Apply Gallery 75 Dry Flowable in late summer to early fall, in early spring, or anytime prior to germination of target weeds, or immediately after cultivation.

Gallery 75 Dry Flowable does not control established weeds. Existing weeds should be controlled by cultivation or with postemergence herbicides. Weed residues, prunings, and trash should be removed or thoroughly mixed into the soil prior to treatment. Soil in non-turf areas should be in good condition and free of clods at the time of application. Gallery 75 Dry Flowable is stable on the soil surface for up to 21 days, but must be activated to be effective. A single rainfall or sprinkler irrigation of 0.5 inches or more, or flood irrigation after application is necessary to activate Gallery 75 Dry Flowable. If Gallery 75 Dry Flowable is not activated by rainfall or irrigation within 21 days of application, erratic weed control may result. In non-

turf areas, if weeds emerge due to lack of rainfall or irrigation, shallow cultivate to a depth of 1 to 2 inches to destroy existing weeds or remove by hand.

**In Arizona:** The state of Arizona has not approved Gallery 75 Dry Flowable for use on plants grown for commercial production such as: turf sod farms, ornamentals and nursery stock grown for resale, and non-bearing fruit and nut trees and non-bearing vineyards.

### Mixing and Application Directions

#### Gallery 75 Dry Flowable - Alone

Check to be sure spray equipment is clean and not contaminated. Using clean water, fill the tank to 1/2 of the final volume required and start agitation. Slowly add the correct quantity of Gallery 75 Dry Flowable to the spray tank, continue agitation and complete filling the tank. Maintain agitation during filling and through application. Agitation should be sufficient to create a rippling or rolling action on the liquid surface. Sparger pipe agitation generally provides the best agitation. In-line screens and strainers should be no finer than 16 mesh.

If spraying and agitation is stopped, Gallery 75 Dry Flowable may settle to the bottom of the spray tank. If settling occurs, material must be re-suspended before continuing spray application. Clean the spray tank, lines and screens thoroughly after use.

#### Gallery 75 Dry Flowable in Tank Mixes

Add Gallery 75 Dry Flowable to the spray tank as described above. Fill the spray tank to 3/4 of the final volume required. Add other formulations in this order: (1) other dry flowables, (2) wettable powders, (3) aqueous suspensions, (4) flowables, (5) liquids, (6) solutions and emulsifiable or liquid concentrates. Allow two to three minutes between the addition of each product, and agitate continuously until the product(s) are completely dispersed in water. Maintain agitation during filling and through application. If a build-up of materials is observed on the walls of the spray tank, wash the tank with soapy water between fillings, rinse and then continue the spraying operation. Follow label directions for each material added to the tank.

#### Application

Apply Gallery 75 Dry Flowable with a properly calibrated low pressure herbicide sprayer that provides uniform distribution. Nozzle screens should be no finer than 50 mesh (50 mesh is finer than 16 mesh). In-line screens and strainers should be no finer than 16 mesh. Apply Gallery 75 Dry Flowable in 10 to 200 gallons of water carrier per acre. As the amount of spray volume decreases, the importance of accurate calibration and uniform application increases. Take precautions to avoid spray drift when applying Gallery 75 Dry Flowable. Drift may result in reduced germination or emergence of non-target plants adjacent to the treated area. Maintain agitation from mixing through application. Avoid boom overlaps that will increase rates above those recommended. Calibrate application equipment prior to use, according to manufacturer's directions. Check calibration frequently to be sure equipment is working properly and distributing spray uniformly.

### Weeds Controlled or Suppressed by Gallery 75 Dry Flowable

Weeds controlled when applied at 0.66 lb/acre (0.25 oz/1000 sq ft)

#### Common Name

aster, slender  
bursage, annual  
celery, wild  
chickweed, common  
clover, white  
cudweed, purple  
fiddleneck, coast  
filaree, redstem  
fleabane, blackleaved  
fleabane, dwarf

#### Scientific Name

*Aster exilis*  
*Ambrosia acanthicarpa*  
*Apium leptophyllum*  
*Stellaria media*  
*Trifolium repens*  
*Gnaphalium purpureum*  
*Amsinckia intermedia*  
*Erodium cicutanum*  
*Conyza bonariensis*  
*Conyza ramosissima*

groundcherry, lanceleaf  
henbit  
horseweed  
knotweed, prostrate  
lambsquarters, common  
mallow, little  
mustard, Indian  
mustard, wild  
nightshade, black  
pepperweed, Virginia  
pigweed  
pineappleweed  
plantain, slender  
purslane, common  
radish, wild  
ragweed, common  
rocket, London  
rockpurslane, desert  
shepherdspurse  
sibara  
smartweed, Pennsylvania  
sowthistle, annual  
speedwell, purslane  
spurge, spotted  
telegraphplant  
thistle, Russian  
velvetleaf

*Physalis lanceifolia*  
*Lamium amplexicaule*  
*Conyza canadensis*  
*Polygonum aviculare*  
*Chenopodium album*  
*Malva parviflora*  
*Brassica juncea*  
*Sinapis arvensis*  
*Solanum nigrum*  
*Lepidium virginicum*  
*Amaranthus* spp.  
*Matricaria matricarioides*  
*Plantago elongata*  
*Portulaca oleracea*  
*Raphanus raphanistrum*  
*Amorosa artemisiifolia*  
*Sisymbrium irio*  
*Calandrinia ciliata*  
*Capsella bursa-pastoris*  
*Sibara virginica*  
*Polygonum pennsylvanicum*  
*Sonchus oleraceus*  
*Veronica peregrina*  
*Euphorbia maculata*  
*Heterotheca grandiflora*  
*Salsola iberica*  
*Abutilon theophrasti*

In addition to the weeds listed at the 0.66 lb/acre rate, the following weeds will be controlled at 1.0 lb/acre (0.38 oz/1000 sq ft)

<u>Common Name</u>	<u>Scientific Name</u>
aster, heath	<i>Aster ericoides</i>
bittercress	<i>Cardamine oligosperma</i>
bittercress, hairy	<i>Cardamine hirsuta</i>
brassbuttons, southern	<i>Cotula australis</i>
carrot, wild	<i>Daucus carota</i>
chamberbitter	<i>Phyllanthus urinaria</i>
chickweed, mouseear	<i>Cerastium vulgatum</i>
dandelion	<i>Taraxacum officinale</i>
eclipta	<i>Eclipta prostrata</i>
galinsoga, hairy	<i>Galinsoga ciliata</i>
geranium, Carolina	<i>Geranium carolinianum</i>
groundsel, common	<i>Senecio vulgaris</i>
ladysthumb	<i>Polygonum persicaria</i>
lettuce, prickly	<i>Lactuca serriola</i>
mallow, dwarf	<i>Malva rotundifolia</i>
maretail	<i>Hippuris vulgaris</i>
mayweed	<i>Anthemis cotula</i>
morningglory, ivyleaf	<i>Ipomoea hederacea</i>
mustard, black	<i>Brassica nigra</i>
pennywort	<i>Hydrocotyle</i> spp.
phyllanthus, long-stalk	<i>Phyllanthus tenellus</i>
plantain, bracted	<i>Plantago aristata</i>
plantain, broadleaf	<i>Plantago major</i>
plantain, buckhorn	<i>Plantago lanceolata</i>
pokeweed, common	<i>Phytolacca americana</i>
rockpurslane, redmaids	<i>Calandrinia ciliata</i> var. <i>menziesii</i>

sida, prickly  
 sorrel, red  
 speedwell, thymeleaf  
 spurge, hyssop  
 sweetclover, yellow  
 tansymustard, green  
 woodsorrel, yellow

*Sida spinosa*  
*Rumex acetosella*  
*Veronica serpyllifolia*  
*Euphorbia hyssopifolia*  
*Melilotus officinalis*  
*Descurainia pinnata* ssp. *brachycarpa*  
*Oxalis stricta*

In addition to the weeds listed at the 0.66 lb/acre and 1.0 lb/acre rates, the following weeds will be controlled at 1.33 lb/acre (0.5 oz/1000 sq ft)

<u>Common Name</u>	<u>Scientific Name</u>
burclover, California	<i>Medicago polymorpha</i>
datura	<i>Datura</i> spp.
dogfennel	<i>Eupatorium capillifolium</i>
eveningprimrose	<i>Oenothera</i> spp.
fescue, rattail	<i>Vulpia myuros</i>
filaree, whitestem	<i>Erodium moschatum</i>
goosefoot, nettleleaf	<i>Chenopodium murale</i>
jimsonweed	<i>Datura stramonium</i>
knotweed, silversheath	<i>Polygonum argyrocoleon</i>
kochia	<i>Kochia scoparia</i>
medic, black	<i>Medicago lupulina</i>
mullein, turkey	<i>Eremocarpus setigerus</i>
nettle, burning	<i>Urtica urens</i>
oxtongue, bristly	<i>Picris echioides</i>
pimpernel, scarlet	<i>Anagallis arvensis</i>
sowthistle, spiny	<i>Sonchus asper</i>
spurge, petty	<i>Euphorbia peplus</i>
spurge, prostrate	<i>Euphorbia humistrata</i>
sunflower	<i>Helianthus</i> spp.
swinecress	<i>Coronopus didymus</i>
thistle, musk	<i>Carduus nutans</i>
willoweed, panicle	<i>Epilobium paniculatum</i>
woodsorrel, creeping	<i>Oxalis corniculata</i>

In addition to the weeds controlled, the following weeds will be partially controlled or suppressed at 1.33 lb/acre (0.5 oz/1000 sq ft)

<u>Common Name</u>	<u>Scientific Name</u>
bindweed, field	<i>Convolvulus arvensis</i>
carpetweed	<i>Mollugo verticillata</i>
dock, curly	<i>Rumex crispus</i>
mallow, Venice	<i>Hibiscus trionum</i>
milkweed, honeyvine	<i>Ampelamus albidus</i>
morningglory, tall	<i>Ipomoea purpurea</i>
pustley, Florida	<i>Richardia scabra</i>

## Approved Uses

### **Established Turfgrasses**

**Gallery 75 Dry Flowable is recommended as a preemergence treatment for control of certain broadleaf weeds in established cool season and warm season turfgrasses.**

Apply Gallery 75 Dry Flowable in late summer to early fall, in early spring, or anytime prior to germination of target weeds.

Refer to "General Information" section of this label prior to using this product on turf.

### **Special Use Precautions**

Apply Gallery 75 Dry Flowable only on established turf. Apply to newly seeded turf (including overseeded turf) only after seedlings are established (three leaf stage and tillering) and well rooted. Do not overseed established turf sooner than 60 days following an application of Gallery 75 Dry Flowable.

Do not use Gallery 75 Dry Flowable on golf course putting greens or tees.

Do not use Gallery 75 Dry Flowable on dichondra.

Do not use Gallery 75 Dry Flowable on grass grown for seed.

### **Treatment of Turf Species Not Listed on the Gallery 75 Dry Flowable Label**

Users who wish to use Gallery 75 Dry Flowable on turf species not recommended on this label may determine the suitability for such uses by treating a small area at a recommended rate. Prior to treatment of larger areas, the treated area should be observed for any sign of herbicidal injury during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious to the target turf species. **The user assumes the responsibility for any turf damage or other liability resulting from use of Gallery 75 Dry Flowable on turf species not recommended on this label.**

### **Tank Mixing**

Gallery 75 Dry Flowable may be tank mixed with Surflan® A S. and applied to warm season turfgrass only as a preemergence treatment to provide broad spectrum annual grass and broadleaf weed control. Gallery 75 Dry Flowable may also be applied as a separate treatment to supplement Balan 2.5G or Team 2G applications in cool and warm season turf. Gallery 75 Dry Flowable may be tank mixed with postemergence broadleaf herbicides registered for use on established turf to control existing broadleaf weeds and provide residual preemergence broadleaf weed control. Applied as directed, Gallery 75 Dry Flowable tank mixes will provide control of susceptible weed species listed on the respective labels. When using Gallery 75 Dry Flowable in tank mix combinations with other products read and follow all applicable use directions, precautions, and limitations on the respective product labels.

**Gallery 75 Dry Flowable is recommended for use on the following turf species:**

### **Established Cool Season Turf Grasses**

#### **Common Name**

bentgrass, creeping  
bentgrass, colonial  
bluegrass, Kentucky  
fescue, chewing  
fescue, creeping red  
fescue, sheeps  
fescue, tall  
ryegrass, perennial

#### **Scientific Name**

*Agrostis stolonifera*  
*Agrostis tenuis*  
*Poa pratensis*  
*Festuca rubra* var. *commutata*  
*Festuca rubra*  
*Festuca ovina*  
*Festuca arundinaceae*  
*Lolium perenne*

**Established Warm Season Turf Grasses****Common Name**

bahia grass

Bermudagrass

buffalograss

centipedegrass

St. Augustinegrass

zoysiagrass

zoysiagrass

**Scientific Name***Paspalum notatum*var. *Saurae parodi**Cynodon dactylon**Buchloe dactyloides**Eremochloa ophiuroides**Stenotaphrum secundatum**Zoysia japonica**Zoysia tenuifolia***Landscape Ornamentals, Container Grown Ornamentals, Nursery Stock, Ground Covers, Ornamental Bulbs, Non-bearing† Fruit and Nut Trees and Non-bearing† Vineyards, and Christmas Tree Plantations**

†Non-bearing fruit and nut trees and non-bearing vineyards are defined as plants which will not bear fruit for at least one year after treatment.

Gallery 75 Dry Flowable is recommended as a preemergence treatment for control of certain broadleaf weeds in landscape ornamentals, container grown ornamentals, nursery stock, ground covers, ornamental bulbs, non-bearing fruit and nut trees and non-bearing vineyards, and Christmas tree plantations.

Apply Gallery 75 Dry Flowable in late summer to early fall, in early spring, or anytime prior to germination of target weeds, or immediately after cultivation.

**Note:** Refer to the "General Information" section of this label for use precautions and information on mixing and application, application rates, and weeds controlled; refer to using this product.

**Special Use Precautions**

**To avoid possible injury, do not apply Gallery 75 Dry Flowable to:**

Nursery seedbeds or forest or Christmas tree seedling transplant beds.

Unrooted liners or cuttings that have been planted in pots for the first time.

Pots less than four inches wide.

Bedding plants or areas where bedding plants will be planted or transplanted within one year following application.

Ground covers until they are established and well rooted.

**Do not apply Gallery 75 Dry Flowable to newly transplanted ornamentals, nursery stock, ground covers, ornamental bulbs, non-bearing fruit and nut trees and non-bearing vineyards or Christmas trees until soil or potting media has been settled by packing and irrigation or rainfall and no cracks are present or plant injury may occur.**

When planting into a site treated with Gallery 75 Dry Flowable in the past 8 months, use untreated soil as fill around roots when replacing plants or injury may occur.

Repeat applications at 1.0 pound per acre or more of Gallery 75 Dry Flowable should not be made sooner than 60 days after a previous application of Gallery 75 Dry Flowable. Do not apply more than a total of 4.0 pounds per acre of Gallery 75 Dry Flowable within a 12 month period.

**Treatment of Plant Species Not Listed on the Gallery 75 Dry Flowable Label**

Users who wish to use Gallery 75 Dry Flowable on plant species not recommended on this label may determine the suitability for such uses by making a trial application of Gallery 75 Dry Flowable at a recommended rate to a small number of plants. Prior to using Gallery 75 Dry Flowable on a larger number

of plants, the treated plants should be observed for any sign of herbicidal injury during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious to the target plant species. The user assumes the responsibility for any plant damage or other liability resulting from use of Gallery 75 Dry Flowable on plant species not recommended on this label.

**Note: Do not apply Gallery 75 Dry Flowable to the following plant species or injury may occur:**

*Ajuga* spp. (bugle)  
*Brassica* spp. (mustard)  
*Euonymus alatus 'compacta'* (dwarf burning bush)  
*Euphorbia* spp. (spurge)  
*Hydrangea* spp. (hydrangea)  
*Iberi* spp. (candy cane)  
*Juniperus horizontalis 'Prince of Wales'* (Prince of Wales Juniper)  
*Melaleuca quinquenervia* (cajepout tree)  
*Rhododendron carolinianum* (Carolina rhododendron)  
*Rhododendron catawbiense 'Roseum elegans'* (Roseum elegans rhododendron)  
*Sedum* spp. (stonecrop)  
*Yucca recurvifolia* (green yucca)

### Tank Mixing

Gallery 75 Dry Flowable may be tank mixed with Roundup or other registered postemergence herbicides to control existing unwanted vegetation and provide residual preemergence broadleaf weed control in ornamental areas. Gallery 75 Dry Flowable may also be tank mixed with Surlan A.S. herbicide and applied preemergence to provide broad spectrum control of annual grasses and broadleaf weeds in ornamental areas. Applied as directed, Gallery 75 Dry Flowable tank mixes will provide control of susceptible weed species listed on the respective labels. When using Gallery 75 Dry Flowable in tank mix combination with other products, read and follow all applicable use directions, precautions, and limitations on the respective product labels.

**Note: Do not apply sprays containing Roundup over the top of ornamental plants. Extreme care must be exercised to prevent contact of sprays containing Roundup with foliage or stems of turfgrasses, trees, shrubs, or other desirable vegetation since severe damage or death may result. If spraying Roundup in areas adjacent to desirable plants, use a shield to prevent spray from contacting foliage or stems of desirable plants.**

Gallery 75 Dry Flowable may be used on the following established plant species:  
 (Note: Limitations on recommended treatment methods)

### Trees

<u>Scientific Name/Common Name</u>	<u>Recommended Treatment Method:</u> C = Container Grown F = Field Grown
<i>Abies concolor</i> White fir	F
<i>Abutilon hybridum</i> Albus-flowering maple	F
Luteus-flowering maple	F
Roseus-flowering maple	F
Tangerine-flowering maple	F
Vesuvius red-flowering maple	F
<i>Acer ginnala</i> Flame maple	F
<i>Acer rubrum</i> Red maple	F
Red sunset maple	F

<i>Acer saccharinum</i>	
Silver maple	C, F
<i>Areacastrum romanzoffianum</i>	
Queen palm	C, F
<i>Betula nigra</i>	
River birch	C, F
<i>Betula papyrifera</i>	
Paper birch	F
<i>Brachychiton populneus</i>	
Bottle tree	C, F
<i>Bucida buceras</i>	
Black olive	F
<i>Ceratonia siliqua</i>	
Carob	F
<i>Cercis canadensis</i>	
Redbud	C, F
<i>Chamaecyparis obtusa</i>	
Fillcoides-fernsprzy cypress	F
Gracilis-slender Hinoki cypress	F
<i>Chamaecyparis pisifera</i>	
Sawara-false cypress	F
Squarrosa-moss cypress	F
<i>Chamaedorea cataractarum</i>	
Palm	F
<i>Chamaedorea costaricana</i>	
Palm	F
<i>Chamaedorea elegans</i>	
Parlor palm	F
<i>Cornus florida</i>	
Cloud nine dogwood	C, F
Flowering dogwood	C, F
<i>Crataegus viridis</i>	
Green hawthorn	F
<i>Cupaniopsis anacardioides</i>	
Carrot wood	F
<i>Cupressus glabra</i>	
Arizona cypress	F
<i>Elaeagnus angustifolia</i>	
Russian olive	C, F
<i>Eucalyptus camaldulensis</i>	
Red gum eucalyptus	F
<i>Eucalyptus cinerea</i>	
Mealy eucalyptus	F
Silver dollar eucalyptus	F
<i>Eucalyptus microtheca</i>	
Coolibah tree	C, F
<i>Eucalyptus sideroxylon</i>	
Red ironbark eucalyptus	F
<i>Ficus benjamina</i>	
Ficus	C, F
Mini ficus	C, F
<i>Fraxinus udhei</i>	
Shamel ash	C, F
<i>Ginkgo biloba</i>	
Maidenhair tree	F
<i>Gleditsia triacanthos</i>	
Honey locust	F
Shademaster honey locust	F

<i>Heteromeles arbutiflora</i>	
Toyon	F
<i>Illicium floridanum</i>	
Florida anise-tree	C, F
<i>Juniperus virginiana</i>	
Eastern red cedar	C, F
<i>Liquidambar styraciflua</i>	
American sweet gum	F
<i>Magnolia grandiflora</i>	
Southern magnolia	C, F
<i>Morus alba</i>	
White mulberry	F
<i>Musa aluminata</i>	
Banana	C, F
<i>Oxydendrum arboreum</i>	
Sourwood	C, F
<i>Picea abies</i>	
Norway spruce	F
Pendula-weeping Norway spruce	F
Repens-spreading Norway spruce	F
<i>Picea glauca conica</i>	
Dwarf Alberta spruce	F
<i>Picea pungens</i>	
Dwarf globe blue spruce	C, F
Glauca-Colorado blue spruce	F
Hoopsii-Hoop's blue spruce	F
Koster-Koster blue spruce	F
<i>Pinus aristata</i>	
Bristlecone pine	F
<i>Pinus canariensis</i>	
Canary Island pine	F
<i>Pinus contorta</i>	
Shore pine, beach pine	F
<i>Pinus eldarica</i>	
Eldarica pine	C, F
<i>Pinus leucodermis</i>	
Bosnian pine	C, F
<i>Pinus mugo</i>	
Pumilo- shrubby swiss mountain pine	C, F
<i>Pinus nigra</i>	
Austrian black pine	C, F
<i>Pinus radiata</i>	
Monterey pine	F
<i>Pinus strobus</i>	
Eastern white pine	C, F
<i>Pinus sylvestris</i>	
Columnar Scotch pine	C, F
Scotch pine	C, F
<i>Pinus thunbergiana</i>	
Japanese black pine	F
<i>Platanus occidentalis</i>	
American sycamore	F
<i>Platanus racemosa</i>	
California sycamore	F
<i>Podocarpus spp.</i>	
Podocarpus	F
<i>Populus deltoides</i>	
Cottonwood	F

<i>Prosopis chilensis</i>	
Chilean mesquite	C, F
<i>Prunus yedoensis</i>	
Yoshino flowering cherry	F
<i>Quercus ilicifolia</i>	
Bear oak	F
<i>Quercus palustris</i>	
Pin oak	F
<i>Quercus phellos</i>	
Willow oak	C, F
<i>Quercus rubra</i>	
Red oak	C, F
<i>Quercus virginiana</i>	
Live oak	C, F
<i>Salix babylonica</i>	
Babylon weeping willow	F
Corkscrew willow	F
<i>Sequoiadendron giganteum</i>	
Giant sequoia	F
<i>Swietenia mahogani</i>	
Mahogany	F
<i>Tabebuia caraiba</i>	
Yellow tab	F
<i>Tsuga canadensis</i>	
Eastern hemlock	C, F
<i>Ulmus parvifolia</i>	
Chinese elm	F
<i>Washingtonia robusta</i>	
Mexican fan palm	F

**Ornamental Shrubs**

**Recommended  
Treatment Method:**  
C = Container Grown  
F = Field Grown

<u>Scientific Name/Common Name</u>	
<i>Abelia grandiflora</i>	
Edward Goucher abelia	C, F
Glossy abelia	C, F
<i>Acacia abyssinica</i>	
Abyssinica acacia	C, F
<i>Acacia redolens</i>	
Prostrate acacia	F
<i>Acacia stenophylla</i>	
Shoestring acacia	C, F
<i>Acalypha wilkesiana</i>	
Copper leaf	C, F
<i>Acer ginnala</i>	
Amur maple	C, F
<i>Acer palmatum</i>	
Coral bark Japanese maple	C, F
Dwarf Japanese maple	C, F
<i>Agave americana</i>	
Century plant	F
<i>Astilbe chinensis</i>	
False spiraea	C, F
<i>Athyrium nipponicum</i>	
Japanese painted fern	C, F

<i>Baccharis pilularis</i>	
Coyotebush	F
<i>Berberis gladwynensis</i>	
William Penn barberry	C, F
<i>Berberis mentorensis</i>	
Mentor barberry	C, F
<i>Berberis thunbergii</i>	
Atropurea-redleaf Japanese barberry	F
Aurea-golden Japanese barberry	F
Crimson pygmy barberry	C, F
Rose glow barberry	C, F
<i>Bougainvillea</i> spp.	
Barbara Karst	F
California gold	F
Pink pixie	C, F
Scarlet O'Hara	F
Temple fire	C, F
Texas dawn	F
<i>Buxus microphylla japonica</i>	
Japanese boxwood	C, F
<i>Buxus microphylla Koreana</i>	
Korean boxwood	F
<i>Callistemon citrinus</i>	
Lemon bottlebrush	F
<i>Callistemon viminalis</i>	
Weeping bottlebrush	C, F
<i>Calluna vulgaris</i>	
Spring torch scotch heather	C, F
<i>Camellia japonica</i>	
Camellia	C, F
<i>Cassia artemisioides</i>	
Feathery cassia	C, F
<i>Ceanothus</i> spp.	
Wild lilac	F
<i>Cephalotaxus drupacae</i>	
Plum yew	C, F
<i>Cerastium tomentosum</i>	
Snow-in-summer	C, F
<i>Chamaecyparis obtusa</i>	
Kosteri cypress	F
Nana-dwarf Hinoki cypress	F
Torulosa cypress	F
<i>Chamaecyparis pisifera</i>	
Filifera-thread cypress	F
<i>Chrysalidocarpus lutescens</i>	
Areca palm	F
<i>Clethra alnifolia</i>	
Summersweet	C, F
<i>Cleyera japonica</i>	
Japanese cleyera	C, F
<i>Cornus alba</i>	
Sibirica-Siberian dogwood	F
<i>Cornus stolonifera</i>	
Baileyi-red-osier dogwood	F
Flaviramea-yellowtwig dogwood	F
<i>Cotinus coggygia</i>	
Royal purple smoke tree	C, F

<i>Cotinus dammeri</i>	
Coral beauty smoke tree	C, F
Eichholz smoke tree	C, F
<i>Cotoneaster adpressus</i>	
Praecox-early cotoneaster	F
<i>Cotoneaster apiculatus</i>	
Cranberry cotoneaster	C, F
<i>Cotoneaster congestus</i>	
Pyrenees cotoneaster	F
<i>Cotoneaster dammeri</i>	
Bearberry cotoneaster	C, F
<i>Cotoneaster himalayan</i>	
Himalayan cotoneaster	F
<i>Cotoneaster horizontalis</i>	
Rock cotoneaster	C, F
<i>Cycas revoluta</i>	
Sago palm	C, F
<i>Cytisus praecox</i>	
Hollandia-warminster broom	F
<i>Cytisus scoparius</i>	
Lena-Scotch broom	F
<i>Daphne odora</i>	
Fragrant daphne	C, F
<i>Deutzia crenata</i>	
Nakiana-dwarf deutzia	F
<i>Deutzia gracilis</i>	
Slender gracilis	F
<i>Dodonea viscosa</i>	
Hopseed bush	F
<i>Elaeagnus pungens</i>	
Fruitland silver berry	C, F
<i>Erica cinerea</i>	
Purple bell heather	C, F
<i>Erica x darleyensis</i>	
Mediterranean pink heather	C, F
<i>Erica vagans</i>	
Cornish heather	C, F
<i>Euonymus fortunei</i>	
Canadale gold euonymus	F
Emerald'n gold euonymus	F
Sunspot euonymus	C, F
<i>Euonymus japonica</i>	
Silver king euonymus	F
Variegated evergreen euonymus	C, F
<i>Euonymus kiatschovica</i>	
Spreading euonymus	F
<i>Forsythia intermedia</i>	
Border forsythia	C, F
<i>Gardenia jasminoides</i>	
August beauty gardenia	C, F
Gardenia	C, F
Radican gardenia	C, F
<i>Gaultheria shallon</i>	
Salal/lemon leaf	C, F
<i>Gelsemium sempervirens</i>	
Carolina jessamine	C, F
<i>Genista pilosa</i>	
Woadwaxen	F

<i>Hibiscus rosa-sinensis</i>	
Ross Estey-hibiscus	C, F
<i>Hibiscus syriacus</i>	
Rose of Sharon-heart	F
Rose of Sharon-red bird	F
Rose of Sharon-woodbridge	F
<i>Ilex aquifolium</i>	
Balkans holly	F
Gold coast holly	F
<i>Ilex aquipernyi</i>	
San Jose holly	C, F
<i>Ilex attenuata</i>	
Savannah holly	C, F
<i>Ilex comuta</i>	
Burford holly	C, F
Dwarf Burford holly	F
Needle point holly	C, F
<i>Ilex crenata</i>	
Compacta-dwarf Japanese holly	C, F
Convexa holly	C, F
Dwarf Chinese holly	C, F
Green luster holly	C, F
Helleri-Heller's Japanese holly	C, F
Hetzii's Japanese holly	C, F
Stokesii Japanese holly	C, F
<i>Ilex glabra</i>	
Compacta-compact inkberry holly	C, F
Nordica-inkberry holly	C, F
<i>Ilex meserveae</i>	
Blue girl holly	F
Blue boy holly	F
Ebony magic holly	F
<i>Ilex vomitoria</i>	
Nana-dwarf yaupon holly	C, F
Pendula-weeping yaupon holly	C, F
yaupon holly	C, F
<i>Illicium annisatum</i>	
Mystery gardenia	C, F
<i>Itea ilicifolia</i>	
Henry Garnet holly leaf sweetspire	C, F
<i>Ixora collinea</i>	
Ixora	C, F
<i>Juniperus chinensis</i>	
Torulosa-hollywood juniper	C, F
Media-old gold juniper	C, F
Pfitzerana-pfitzer juniper	C, F
Pfitzerana glauca-blue juniper	C, F
Sea green juniper	F
<i>Juniperus conferta</i>	
Emerald sea shore juniper	C, F
Shore juniper	C, F

<i>Juniperus horizontalis</i>	
Plumosa-Andorra juniper	C, F
Bar Harbor juniper	C, F
Blue rug juniper	C, F
Creeping juniper	C, F
Dwarf Andorra juniper	C, F
Huntington blue juniper	C, F
Wiltonii-blue carpet juniper	C, F
<i>Juniperus procumbens</i>	
Nana-dwarf Japaneses garden juniper	C, F
<i>Juniperus prostrata</i>	
Prostrata juniper	C, F
<i>Juniperus sabina</i>	
Broadmoor juniper	F
Foemina-Hicks juniper	C, F
Savin juniper	C, F
Tamariscifolia-Tam juniper	F
<i>Juniperus scopulorum</i>	
Emerald green juniper	F
<i>Juniperus squamata</i>	
Blue juniper	F
Blue star juniper	F
Parsonii juniper	C, F
<i>Kalmia latifolia</i>	
Mountain laurel	C, F
<i>Lagerstroemia indica</i>	
Crape myrtle	C, F
<i>Lantana spp.</i>	
Lantana	C, F
<i>Leucothoe axillaris</i>	
Coast leucothoe	C, F
<i>Leucothoe fontainesiana</i>	
Drooping leucothoe	C, F
<i>Ligustrum lucidum</i>	
Glossy privet	C, F
<i>Ligustrum ovalifolium</i>	
California privet	F
<i>Ligustrum japonicum</i>	
Japanese privet	C, F
Wax ligustrum	C, F
Yellow tip ligustrum	C, F
<i>Ligustrum texanum</i>	
Howardi privet	F
Wax leaf privet	C, F
<i>Ligustrum vicaryi</i>	
Vicary golden privet	F
<i>Ligustrum vulgare</i>	
Lodense privet	C, F
<i>Livistona chinensis</i>	
Chinese fountain palm	F
<i>Lonicera periclymenum</i>	
Flowering woodbine	F
Serotina woodbine	F
<i>Lonicera sempervirens</i>	
Trumpet honeysuckle	F
<i>Mahonia bealei</i>	
Leather leaf mahonia	C, F

<i>Mahonia repens</i>	
Creeping mahonia	C, F
<i>Myrica cerifera</i>	
Wax myrtle	C, F
<i>Nandina domestica</i>	
Compacta-dwarf heavenly bamboo	C, F
Harbour dwarf-heavenly bamboo	C, F
Heavenly bamboo	C, F
Nana compacta-heavenly bamboo	C, F
Nana purpurea-heavenly bamboo	C, F
Woods dwarf-heavenly bamboo	C, F
<i>Nerium oleander</i>	
Hardy red oleander	F
Oleander	C, F
Ruby lace oleander	F
<i>Osmanthus fortunei</i>	
Fortunes osmanthus	C, F
<i>Phoenix roelofenii</i>	
Pigmy date palm	C, F
<i>Photinia fraseri</i>	
Fraser's photinia	C, F
<i>Pieris japonica</i>	
Lily-of-the-valley	C, F
Mountain fire lily-of-the-valley	C, F
Snowdrift lily-of-the-valley	C, F
Temple bells lily-of-the-valley	C, F
Valley rose lily-of-the-valley	C, F
Valley valentine lily-of-the-valley	C, F
<i>Pieris japonica x forestii</i>	
Forest flame lily-of-the-valley	C, F
<i>Pinus mugo</i>	
Mugo-mugho pine	C, F
<i>Pittosporum tobira</i>	
Wheeler's dwarf pittosporum	C, F
Green pittosporum	C, F
<i>Plumbago ariculata</i>	
Blue cape plumbago	F
<i>Plumbago capensis</i>	
Plumbago	C, F
<i>Podocarpus macrophyllus</i>	
Yewpine	C, F
<i>Polystichum polyblepharum</i>	
Tassel fern	C, F
<i>Potentilla fragiformis</i>	
Cinquefoil	F
<i>Potentilla fruticosa</i>	
Cinquefoil	C, F
Gold drop potentilla	F
Goldfinger potentilla	C, F
Red ace potentilla	C, F
Sunset potentilla	C, F
Tangerine potentilla	C, F
<i>Potentilla verna</i>	
Spring cinquefoil	C, F
<i>Prunus glandulosa</i>	
Dwarf pink flowering almond	C, F

<i>Pyracantha fortuneana</i>	
Lalendel Monrovia pyracantha	F
Monon pyracantha	F
Red elf hybrid pyracantha	F
Rutgers hybrid pyracantha	C, F
Santa Cruz pyracantha	F
Victory pyracantha	F
<i>Raphiolepis indica</i>	
Charisma-Monruce raphiolepis	F
Enchantress-Moness raphiolepis	F
India hawthorn	C, F
Springtime-Monme raphiolepis	F
<i>Raphiolepis ovata</i>	
Roundleaf raphiolepis	F
<b><i>Rhododendron</i> spp. (including azalea)</b>	
<b>Azaleas</b>	
<i>Rhododendron calendulacem</i>	
Flame azalea	F
Cannon's double azalea	C, F
Golden flare azalea	C, F
Klondike azalea	C, F
<i>Rhododendron indica</i>	
Formosa azalea	C, F
Waucabusa azalea	C, F
<i>Rhododendron kaempferi</i>	
Blue danube azalea	C, F
<i>Rhododendron kerume</i>	
Coral bells azalea	C, F
Hino pink azalea	C, F
Mildred azalea	C, F
Snow azalea	C, F
<i>Rhododendron obtusum</i>	
Hino-crimson azalea	C, F
<i>Rhododendron satsumi</i>	
Gumpo pink azalea	C, F
Higasa azalea	F
Reijn azalea	C, F
<i>Rhododendron</i> spp. hybrids	
Carror azalea	C, F
Fashion azalea	C, F
Gerard Christina azalea	F
Girard Roberta azalea	F
Helmut vogel azalea	F
Hersey red azalea	F
Hot shot azalea	C, F
Hume azalea	F
Inga azalea	F
Irene Koster azalea	C, F
President Clay azalea	C, F
Tradition azalea	C, F

**Rhododendrons**

<i>Rhododendron campylocarpum</i>	
Butterfly rhododendron	F
<i>Rhododendron carolinianum x dauricum</i>	
PJM rhododendron	F
<i>Rhododendron catawbiense</i>	
Catawba rhododendron	C, F
Catawba album rhododendron	C, F
Lord Roberts rhododendron	C, F
Rocket rhododendron	C, F
<i>Rhododendron caucasicum x ponticum</i>	
Cunningham White rhododendron	F
<i>Rhododendron forrestii x griersonianum</i>	
Elizabeth rhododendron	C, F
<i>Rhododendron forrestii repens</i>	
Gomer Waterer rhododendron	C, F
<i>Rhododendron griffithianum</i>	
Jean Marie rhododendron	F
<i>Rhododendron impeditum</i>	
Rhododendron	C, F
<i>Rhododendron maximum</i>	
Rhodie max (rosebay)	F
<i>Rhododendron mucronulatum</i>	
Rhododendron	F
<i>Rhododendron ponticum</i>	
Chionoides rhododendron	F
Daphnoides rhododendron	C, F
<i>Rhododendron racemosum</i>	
Dwarf scarlet wonder rhododendron	C, F
Tribly rhododendron	C, F
Unique rhododendron	C, F
Vulcan rhododendron	C, F
<i>Rhododendron sassthigiatim x carolinianum</i>	
Ramapo rhododendron	F
<i>Rhododendron hybrid spp.</i>	
America rhododendron	C, F
English Roseum rhododendron	F
Nova Zembla rhododendron	C, F
Scintillation rhododendron	C, F
<i>Rhus lancea</i>	
Africa sumac	C, F
<i>Rosa rugosa</i>	
Ramanas rose	F
<i>Rosmarinus officinalis</i>	
Rosemary	F
<i>Skimmia japonica</i>	
Japanese skimmia	C, F
<i>Skimmia reevesiana</i>	
Reeve's skimmia	C, F
<i>Spiraea bumalda</i>	
Anthony Waterer spiraea	C, F
<i>Spiraea japonica</i>	
Dolchia spiraea	C, F
Japanese alpine spiraea	C, F
Shirobana spiraea	C, F
<i>Spiraea vanhouttii</i>	
Bridal wreath	C, F

<i>Syringa rothomagensis</i>	
Chinese lilac	C, F
<i>Syringa vulgaris</i>	
Common lilac	F
<i>Taxus cuspidata</i>	
Japanese yew	F
<i>Tecomaria capensis</i>	
Cape honeysuckle	C, F
<i>Temstroemia gymnanthera</i>	C, F
<i>Thuja occidentalis</i>	
Emerald arborvitae	F
Globosa-globe arborvitae	C, F
Little giant-dwarf arborvitae	F
Nigra-dark American arborvitae	F
Pyramidalis-pyramid arborvitae	F
Rheingold arborvitae	F
Techny arborvitae	F
<i>Thuja orientalis</i>	
ALPUS nana-dwarf golden arborvitae	F
Minima glauca-dwarf arborvitae	F
<i>Veitchia merrilli</i>	
Christmas palm	F
<i>Viburnum bodnantense</i>	
Pink dawn viburnum	C, F
<i>Viburnum davidii</i>	
David viburnum	F
<i>Viburnum japonicum</i>	
Viburnum	F
<i>Viburnum lantana</i>	
Wayfaring tree	F
<i>Viburnum opulus sterile</i>	
Common snowball viburnum	F
<i>Viburnum plicatum tomentosum</i>	
Doublefile viburnum	F
<i>Viburnum x pragense</i>	
Viburnum	F
<i>Viburnum setigerum</i>	
Tea viburnum	F
<i>Viburnum tinus compactum</i>	
Spring bouquet viburnum	F
<i>Viburnum trilobum</i>	
Cranberry bush	C, F
<i>Viburnum trilobum compactum</i>	
Dwarf cranberry bush	C, F
<i>Weigela florida</i>	
Bristol ruby weigela	F
Java red weigela	F
Minuet weigela	F
<i>Xylosma congestum</i>	
Xyloema	F
<i>Yucca filamentosa</i>	
Yucca	C, F

**Ground Covers**

<u>Scientific Name/Common Name</u>	<u>Recommended Treatment Method:</u> C = Container Grown F = Field Grown
<i>Ammophila breviligulata</i> Beachgrass	C, F
<i>Arctotheca calendula</i> Cape weed	F
<i>Carex</i> spp. Variegated carex	C, F
<i>Carpobrotus edulis</i> Largeleaf iceplant	F
<i>Cortaderia selloana</i> Pampas grass	C, F
<i>Delosperma alba</i> White iceplant	F
<i>Descampsia caespitosa</i> Descampsia	C, F
<i>Drosanthemum floribundum</i> Trailing rosea iceplant	F
<i>Drosanthemum hispidum</i> Iceplant	C, F
<i>Festuca ovina glauca</i> Blue fescue	C, F
<i>Gazania</i> spp. Gazania	C, F
<i>Hakonechloa macroaureola</i> Golden hakonechloa	C, F
<i>Hedera canariensis</i> Algerian ivy	F
<i>Hedera helix</i> English ivy	C, F
<i>Hemerocallis</i> spp. Daylily	C, F
<i>Hosta lancifolia</i> Narrow leafed plantain lily	C
<i>Hosta lancifolia</i> Albo-marginata hosta	C, F
<i>Hypericum</i> spp. St. Johnswort	F
<i>Jasminum nitidum</i> Angelwing jasmine	C, F
<i>Lampranthus spectabilis</i> Trailing iceplant	F
<i>Liriope gigantea</i> White lily turf	F
<i>Liriope muscari</i> Big blue lily turf	C, F
Lilac beauty lily turf	C, F
Majestic lily turf	C, F
Monroe white lily turf	C, F
Silvery sunproof lily turf	C, F
Variegated lily turf	C, F
<i>Liriope spicata</i> Green/creeping lily turf	C, F
<i>Lonicera japonica</i> Japanese honeysuckle	F

<i>Miscanthus sinensis</i>	
Eulalia grass	C, F
<i>Ophiopogon japonicus</i>	
Dwarf Mondo grass	C, F
Mondo grass	C, F
<i>Osteospermum fruticosum</i>	
Trailing African daisy	F
<i>Pachysandra terminalis</i>	
Japanese spurge	C, F
<i>Pennisetum alopecuroides</i>	
Fountain grass	C, F
<i>Phalaris arundinacea picta</i>	
Ribbon grass	C, F
<i>Trachelospermum asiaticum</i>	
Asaian jasmine	C, F
<i>Vinca minor</i>	
Dwarf periwinkle	F

### Ornamental Bulbs

Gallery 75 Dry Flowable may be applied for control of susceptible annual weeds in ornamental bulbs, e.g., bulbous iris, daffodil (narcissus), hyacinth and tulip. Apply Gallery 75 Dry Flowable to the soil surface 2 to 4 weeks after planting, but prior to the emergence of annual weeds. Gallery 75 Dry Flowable may also be applied following bulb emergence but prior to flowering and after flowering. For fall planted bulbs, apply Gallery 75 Dry Flowable again in late winter or early spring to weed-free soil surfaces.

#### Special Use Precautions:

Do not apply to tulip plants that have emerged to a height greater than 3/4 inch.  
Do not apply to gladiolus prior to emergence or if corms are less than one inch in diameter.  
Do not apply to bulbs while they are flowering.

### Non-bearing† Fruit and Nut Trees and Non-bearing† Vineyards

<u>Common Name</u>	<u>Recommended Treatment Method:</u>
	<u>C = Container Grown</u>
	<u>F = Field Grown</u>
almond	F
apple	F
apricot	F
avocado	F
blackberry	F
blueberry	F
boysenberry	F
cherry, sour	F
cherry, sweet	F
currant	F
dewberry	F
elderberry	F
fig	F
filbert	F
gooseberry	F
grape, American	F
grape, European	F
grapefruit	F
kiwi	F
lemon	F
loganberry	F

macadamia nut	F
nectarine	F
olive	F
orange	F
peach	F
pear	F
pecan	F
pistachio	F
plum	F
prune	F
pomegranate	F
raspberry	F
walnut, black	F
walnut, English	F

†Non-bearing fruit and nut trees and non-bearing vineyards are defined as plants which will not bear fruit for at least one year after treatment.

### Christmas Tree Plantations

#### Gallery 75 Dry Flowable Alone

Apply Gallery 75 Dry Flowable as a directed spray to the soil surface or as an overtop spray to established plantings of field grown Christmas tree species listed in this label. Do not apply to seedbeds or seedling transplant beds. Apply only to established plantings. Established plants are defined as those that have been transplanted into their final growing location for a sufficient period of time to allow the soil to be firmly settled around the roots from packing and rainfall or irrigation. Follow all instructions provided in the "General Information" section of this label.

#### Tank-Mix Combinations

Tank-mix combinations of Gallery 75 Dry Flowable plus other labeled herbicides may be used as directed or overtop sprays in established Christmas tree plantings. When applied according to use directions, these tank mixes will provide control of susceptible weed species listed on the respective product labels. Refer to tank mix product labels for specific use directions, precautions and limitations before use.

**Gallery 75 Dry Flowable/Roundup:** Apply tank mix combinations of Gallery 75 Dry Flowable plus Roundup only as directed sprays in Christmas tree plantings. When applied according to use directions, Gallery 75 Dry Flowable plus Roundup herbicide will provide postemergence control of susceptible weed species listed on the Roundup label and residual preemergence control of susceptible weed species listed on the Gallery 75 Dry Flowable label. Refer to the Roundup label for specific use directions, precautions and limitations before use.

#### Precautions:

Do not apply sprays containing Roundup over the top of Christmas tree plantings.

Extreme care must be exercised to prevent contact of sprays containing Roundup with foliage or stems of Christmas trees or other desirable plants or severe damage or death may result.

### Non-cropland Areas

Gallery 75 Dry Flowable is recommended as a preemergence herbicide for control of certain broadleaf weeds on industrial sites, utility substations, highway guard rails, sign posts and delineators.

Apply Gallery 75 Dry Flowable in late summer to early fall, in early spring, or anytime prior to germination of target weeds. Areas to be treated should be free of established weeds or existing weeds should be controlled with postemergence herbicides.

Refer to the "General Information" section prior to using this product on non-cropland.

Gallery 75 Dry Flowable is compatible and can be tank mixed with other herbicides registered for use on non-cropland. Applied as directed, tank mixes containing Gallery 75 Dry Flowable will provide control of susceptible weed species listed on the respective labels. All directions, precautions and limitations on the respective product labels apply to the tank mix use.

---

### **Warranty Disclaimer**

---

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. DOWELANCO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

### **Inherent Risks of Use**

---

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application or other factors, all of which are beyond the control of DowElanco or the seller. All such risks shall be assumed by Buyer.

---

### **Limitation of Remedies**

---

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at DowElanco's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of product used.

DowElanco shall not be liable for losses or damages resulting from handling or use of this product unless DowElanco is promptly notified of such loss or damage in writing. In no case shall DowElanco be liable for consequential or incidental damages or losses.

The terms of the "Warranty Disclaimer" above and this "Limitation of Remedies" cannot be varied by any written or verbal statements or agreements. No employee or sales agent of DowElanco or the seller is authorized to vary or exceed the terms of the "Warranty Disclaimer" or this "Limitation of Remedies" in any manner.

Copyright © 1992 DowElanco

Revised 3/21/92

#### **Amendments**

- 1) Revised weed controlled list (excluding grass weeds) to match Snapshot 5 TG label.
- 2) Revised plant species which may be treated list to match Snapshot 2.5 TG label.
- 3) Tall fescue removed from "Established Cool Season Turf Grasses" list.
- 4) Buffalograss (*Buchloe dactyloides*) added to "Established Warm Season Turf Grasses" list.
- 5) Added use on ornamental bulbs
- 6) Added use on Christmas tree plantations.