

1M23

62719-145

1 of 11

**DowElanco**

**Gallery\***  
75 Dry Flowable

**Specialty Herbicide**


A selective preemergence herbicide for control of certain broadleaf weeds in:

- Established turf
- Landscape ornamentals
- Container grown ornamentals
- Nursery stock
- Ground covers
- Ornamental bulbs
- Non-bearing fruit and nut trees and non-bearing vineyards
- Christmas tree plantations
- Non-cropland

**Active Ingredient:**

isoxaben: N-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers .....75%  
 Inert Ingredients .....25%  
 Total .....100%  
 Contains 0.75 pounds active ingredient per pound.

EPA Reg. No. 62713-145  
EPA Est. 464-MI-1  
Net Weight 1 lb



**Precautionary Statements**

**Hazards to Humans and Domestic Animals**

**Keep Out of Reach of Children**

**CAUTION / PRECAUCION:**  
Precaucion al usuario: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

**Causes Eye Irritation • Harmful If Inhaled**

Avoid ingestion, breathing dust or spray mist, and contact with skin, eyes, or clothing. Use eye protection when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**First Aid**

**If in eyes:** Flush eyes with plenty of water. Call a physician if irritation develops.

**If swallowed:** Call a physician or Poison Control Center.

**If on skin:** Wash with plenty of soap and water. Get medical attention if irritation develops.

**If inhaled:** Remove individual to fresh air. If not breathing, provide cardiopulmonary resuscitation assistance and get medical attention.

**Environmental Hazards**

Do not contaminate water when disposing of equipment washwaters. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift may result in reduced germination or emergence of non-target plants adjacent to treated area.

**Notice:** Read the entire label. Use only according to label directions. Before buying or using this product, read Warranty Disclaimer and Limitation of Remedies sections elsewhere on this label.

In case of emergency endangering health or the environment involving this product, call collect 517-636-4400.

**Agricultural Chemical:** Do not ship or store with food, feeds, drugs or clothing.

**BEST AVAILABLE COPY**

**Gallery\* 75 Dry Flowable**

2 of 11

### Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Chemigation: Do not apply Gallery 75 Dry Flowable through any type of irrigation system.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container. Do not store in direct sunlight. Do not store at temperatures above 120°F. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Dispose of in a sanitary landfill, or incineration, or by burning if allowed by state and local authorities. If burned, stay out of smoke.

### General Information

Gallery 75 Dry Flowable herbicide is a preemergence herbicide for control of certain broadleaf weeds in established turf, landscape ornamentals, container grown ornamentals, nursery stock, ground covers, ornamental bulbs, non-bearing fruit and nut trees and non-bearing vineyards, Christmas tree plantations and non-cropland areas. Apply Gallery 75 Dry Flowable in late summer to early fall, in early spring, or anytime prior to germination of target weeds, or immediately after cultivation.

Gallery 75 Dry Flowable does not control established weeds. Existing weeds should be controlled by cultivation or with postemergence herbicides. Weed residues, prunings, and trash should be removed or thoroughly mixed into the soil prior to treatment. Soil in non-turf areas should be in good condition and free of clods at the time of application. Gallery 75 Dry Flowable is stable on the soil surface for up to 21 days, but must be activated to be effective. A single rainfall or sprinkler irrigation of 0.5 inches or more, or flood irrigation after application is necessary to activate Gallery 75 Dry Flowable. If Gallery 75 Dry Flowable is not activated by rainfall or irrigation within 21 days of application, erratic weed control may result. In non-turf areas, if weeds emerge due to lack of rainfall or irrigation, shallow cultivate to a depth of 1 to 2 inches to destroy existing weeds or remove by hand.

In Arizona: The state of Arizona has not approved Gallery 75 Dry Flowable for use on plants grown for commercial production such as: turf sod farms, ornamentals and nursery stock grown for resale, and non-bearing fruit and nut trees and non-bearing vineyards.

### Mixing and Application Directions

#### Gallery 75 Dry Flowable - Alone

Check to be sure spray equipment is clean and not contaminated. Using clean water, fill the tank to 1/2 of the final volume required and start agitation. Slowly add the correct quantity of Gallery 75 Dry Flowable to the spray tank, continue agitation and complete filling the tank. Maintain agitation during filling and through application. Agitation should be sufficient to create a rippling or rolling action on the liquid surface. Sparger pipe agitation generally provides the best agitation. In-line screens and strainers should be no finer than 16 mesh.

If spraying and agitation is stopped, Gallery 75 Dry Flowable may settle to the bottom of the spray tank. If settling occurs, material must be re-suspended before continuing spray application. Clean the spray tank, lines and screens thoroughly after use.


#### Gallery 75 Dry Flowable in Tank Mixes

Add Gallery 75 Dry Flowable to the spray tank as described above and fill the spray tank to 3/4 of the final volume required. Add other formulations in this order: (1) other dry flowables, (2) wettable powders, (3) aqueous suspensions, (4) flowables, (5) liquids, (6) solutions and emulsifiable or liquid concentrates. Allow two to three minutes between the addition of each product, and agitate continuously until the product(s) are completely dispersed in water. Maintain agitation during filling and through application. If a build-up of materials is observed on the walls of the spray tank, wash the tank with soapy water between fillings, rinse and then continue the spraying operation. Follow label directions for each material added to the tank.

#### Application

Apply Gallery 75 Dry Flowable at 0.66 to 1.33 lb/acre with a properly calibrated low pressure herbicide sprayer that provides uniform distribution. Nozzle screens should be no finer than 50 mesh (50 mesh is finer than 16 mesh). In-line screens and strainers should be no finer than 16 mesh. Apply Gallery 75 Dry Flowable in 10 to 200 gallons of water carrier per acre. As the spray volume decreases, the importance of accurate calibration and uniform application increases. Take precautions to avoid spray drift when applying Gallery 75 Dry Flowable. Drift may result in reduced germination or emergence of non-target plants adjacent to the treated area. Maintain agitation from mixing through application. Avoid boom overlaps that will increase rates above those recommended. Calibrate application equipment prior to use, according to manufacturer's directions. Check calibration frequently to be sure equipment is working properly and distributing spray uniformly.

Notes: The container cap for Gallery 75 Dry Flowable may be used to measure the amount of product required to treat small areas. When filled to the level of the inner lip, the cap contains approximately 1/2 ounce which is the amount of Gallery 75 Dry Flowable needed to treat an area of 1,000 square feet.



When filled to this level, container cap contains 1/2 oz (14 grams) of Gallery 75 Dry Flowable.

BEST AVAILABLE COPY

Handwritten notes:   
New rate 1.33 lb/A   
1.07 lb/A   
old increase in rate is OK because of manufacturer use

3 of 11

## Weeds Controlled or Suppressed by Gallery 75 Dry Flowable

Weeds controlled when applied at 0.66 lb/acre (0.25 oz or 7 grams/1000 sq ft)

### Common Name

aster, slender  
bursage, annual  
calery, wild  
chickweed, common  
clover, white  
cudweed, purple  
fiddleneck, coast  
flaxse, redstem  
fleabane, blackleaved  
fleabane, dwarf  
groundcherry, lanceleaf  
henbit  
horseweed  
knotweed, prostrate  
lambquarters, common  
mallow, little  
mustard, Indian  
mustard, wild  
nightshade, black  
pepperweed, Virginia  
pigweed  
pineappleweed  
plantain, slender  
purslane, common  
radish, wild  
ragweed, common  
rocket, London  
rockpurslane, desert  
shepherdspurse  
sibara  
smartweed, Pennsylvania

sowthistle, annual  
speedwell, purslane  
spurge, spotted  
telegraphplant  
thistle, Russian  
velvetleaf

In addition to the weeds listed at the 0.66 lb/acre rate, the following weeds will be controlled at 1.0 lb/acre (0.38 oz or 10.5 grams/1000 sq ft)

### Common Name

aster, heath  
bittercress  
bittercress, hairy  
brassbuttons, southern  
carrot, wild  
chickweed, common  
chickweed, Russian  
dandelion  
eclipta  
gallinoga, hairy  
geranium, Carolina  
groundsel, common  
ladythumb  
lettuce, prickly  
mallow, dwarf  
marretail

### Scientific Name

*Aster exilis*  
*Ambrosia scanthicarpa*  
*Apium leptophyllum*  
*Stellaria media*  
*Triticum repens*  
*Gnaphalium purpureum*  
*Ameiopsis intermedia*  
*Erodium cicutarium*  
*Coryza bonariensis*  
*Coryza ramosissima*  
*Physalis lanceifolia*  
*Lamium amplexicaule*  
*Coryza canadensis*  
*Polygonum aviculare*  
*Chenopodium album*  
*Malva parviflora*  
*Brassica juncea*  
*Sinapis arvensis*  
*Solanum nigrum*  
*Lepidium virginicum*  
*Amaranthus* spp.  
*Matricaria matricaria toides*  
*Plantago elongata*  
*Portulaca oleracea*  
*Raphanus raphanistrum*  
*Ambrosia artemisiifolia*  
*Sisymbrium irio*  
*Calandrinia ciliata*  
*Capsella bursa-pastoris*  
*Sibara virginica*  
*Polygonum pennsylvanicum*  
*Sonchus oleraceus*  
*Veronica peregrina*  
*Euphorbia maculata*  
*Heterotheca grandiflora*  
*Salsola iberica*  
*Abutilon theophrasti*

### Scientific Name

*Aster ericoides*  
*Cardamine oligosperma*  
*Cardamine hirsuta*  
*Cotula australis*  
*Daucus carota*  
*Phyllanthus urinaria*  
*Cerastium vulgatum*  
*Taraxacum officinale*  
*Eclipta prostrata*  
*Gallinoga ciliata*  
*Geranium carolinianum*  
*Senecio vulgaris*  
*Polygonum persicaria*  
*Lactuca serriola*  
*Malva rotundifolia*  
*Hippuris vulgata*

mayweed  
morningglory, ivyleaf  
mustard, black  
pennywort  
phyllanthus, long-stalk  
plantain, bracted  
plantain, broadleaf  
plantain, buckhorn  
pokeweed, common  
rockpurslane, redmaids

sida, prickly  
sorrel, red  
speedwell, thymeleaf  
spurge, hyssop  
sweetclover, yellow  
tansymustard, green

woodsorrel, yellow

*Anthemis cotula*  
*Ipomoea hederacea*  
*Brassica nigra*  
*Hydrocotyle* spp.  
*Phyllanthus tenellus*  
*Plantago aristata*  
*Plantago major*  
*Plantago lanceolata*  
*Phytolacca americana*  
*Calandrinia ciliata* var. *menziesii*  
*Sida spinosa*  
*Rumex acetosella*  
*Veronica serpyllifolia*  
*Euphorbia hyssopifolia*  
*Melilotus officinalis*  
*Descurainia pinnata* spp. *brachycarpa*  
*Oxalis stricta*

In addition to the weeds listed at the 0.66 lb/acre and 1.0 lb/acre rates, the following weeds will be controlled at 1.33 lb/acre (0.5 oz or 14 grams/1000 sq ft)

### Common Name

burclover, California  
detuna  
doglennel  
eveningprimrose  
fescue, rattail  
flaxse, whitestem  
goosefoot, nettleleaf  
jimsonweed  
knotweed, silversheath  
  
kochia  
medic, black  
mullein, turkey  
nettle, burning  
oxongus, prickly  
pimpernel, scarlet  
sowthistle, spiny  
spurge, petty  
spurge, prostrate  
sunflower  
swinecress  
thistle, musk  
willoweed, pericle  
woodsorrel, creeping

### Scientific Name

*Medicago polymorpha*  
*Desura* spp.  
*Eupatorium capillitium*  
*Oenothera* spp.  
*Vulpia myuros*  
*Erodium moschatum*  
*Chenopodium murale*  
*Desura stramonium*  
*Polygonum argyrocoleon*  
*Kochia scoparia*  
*Medicago lupulina*  
*Eriogonum cicutarium*  
*Urtica urens*  
*Picris echioides*  
*Anagallis arvensis*  
*Sonchus asper*  
*Euphorbia pepus*  
*Euphorbia humifusa*  
*Helianthus* spp.  
*Coronopus didymus*  
*Carduus nutans*  
*Epilobium periclymenum*  
*Oxalis corniculata*

In addition to the weeds controlled, the following weeds will be partially controlled or suppressed at 1.33 lb/acre (0.5 oz or 14 grams/1000 sq ft)

### Common Name

bindweed, field  
carpetweed  
dock, curly  
mallow, Venice  
milkweed, honeyvine  
morningglory, tall  
pusley, Florida

### Scientific Name

*Cynodorus arvensis*  
*Xanthoxa verticillata*  
*Rumex crispus*  
*Ficaria verna*  
*Ampelodesmos abidus*  
*Ipomoea purpurea*  
*Rhizoglyphis scabra*

**BEST AVAILABLE COPY**

## Approved Uses

### Established Turfgrasses

Gallery 75 Dry Flowable is recommended as a preemergence treatment for control of certain broadleaf weeds in established cool season and warm season turfgrasses.

Apply Gallery 75 Dry Flowable in late summer to early fall, in early spring, or anytime prior to germination of target weeds.

Refer to "General Information" section of this label prior to using this product on turf.

#### Special Use Precautions

Apply Gallery 75 Dry Flowable only on established turf. Apply to newly seeded turf (including overseeded turf) only after seedlings are established (three leaf stage and tillering) and well rooted. Do not overseed established turf sooner than 60 days following an application of Gallery 75 Dry Flowable.

Do not use Gallery 75 Dry Flowable on golf course putting greens or tees.

Do not use Gallery 75 Dry Flowable on dichondra.

Do not use Gallery 75 Dry Flowable on grass grown for seed.

#### Treatment of Turf Species Not Listed on the label for Gallery 75 Dry Flowable

Users who wish to use Gallery 75 Dry Flowable on turf species not recommended on this label may determine the suitability for such uses by treating a small area at a recommended rate. Prior to treatment of larger areas, the treated area should be observed for any sign of herbicidal injury during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious to the target turf species. The user assumes the responsibility for any turf damage or other liability resulting from use of Gallery 75 Dry Flowable on turf species not recommended on this label.

#### Tank Mixing

Gallery 75 Dry Flowable may be tank mixed with Surflan® A.S. and applied to warm season turfgrass only as a preemergence treatment to provide broad spectrum annual grass and broadleaf weed control. Gallery 75 Dry Flowable may also be applied as a separate treatment to supplement Balan 2.5G or Team 2G applications in cool and warm season turf. Gallery 75 Dry Flowable may be tank mixed with postemergence broadleaf herbicides registered for use on established turf to control existing broadleaf weeds and provide residual preemergence broadleaf weed control. Applied as directed, Gallery 75 Dry Flowable tank mixes will provide control of susceptible weed species listed on the respective labels. When using Gallery 75 Dry Flowable in tank mix combinations with other products, read and follow all applicable use directions, precautions, and limitations on the respective product labels.

Gallery 75 Dry Flowable is recommended for use on the following turf species:

#### Established Cool Season Turf Grasses

##### Common Name

bentgrass, creeping  
bentgrass, colonial  
bluegrass, Kentucky  
fescue, chewing  
fescue, creeping red  
fescue, sheep  
fescue, tall  
ryegrass, perennial

##### Scientific Name

*Agrostis scolonifera*  
*Agrostis tenuis*  
*Poa pratensis*  
*Festuca rubra* var. *commutata*  
*Festuca rubra*  
*Festuca ovina*  
*Festuca arundinacea*  
*Lolium perenne*

#### Established Warm Season Turf Grasses

##### Common Name

bahiagrass  
Bermudagrass  
buffalograss  
centipedegrass  
St. Augustinegrass  
zoysiagrass  
zoysiagrass

##### Scientific Name

*Paspalum notatum*  
var. *Saurae parodi*  
*Cynodon dactylon*  
*Buchloe dactyloides*  
*Eremochloa ophiuroides*  
*Stenotaphrum secundatum*  
*Zoysia japonica*  
*Zoysia tenuifolia*

#### Landscape Ornamentals, Container Grown Ornamentals, Nursery Stock, Ground Covers, Ornamental Bulbs, Non-bearing† Fruit and Nut Trees and Non-bearing† Vineyards, and Christmas Tree Plantations

† Non-bearing fruit and nut trees and non-bearing vineyards are defined as plants which will not bear fruit for at least one year after treatment.

Gallery 75 Dry Flowable is recommended as a preemergence treatment for control of certain broadleaf weeds in landscape ornamentals, container grown ornamentals, nursery stock, ground covers, ornamental bulbs, non-bearing fruit and nut trees and non-bearing vineyards, and Christmas tree plantations.

Apply Gallery 75 Dry Flowable anytime prior to germination of target weeds, or immediately after cultivation.

Note: Refer to the "General Information" section of this label for use precautions and information on mixing and application, application rates, and weeds controlled prior to using this product.

#### Special Use Precautions

To avoid possible injury, do not apply Gallery 75 Dry Flowable to: Nursery seedbeds or forest or Christmas tree seeding transplant beds.

Unrooted liners or cuttings that have been planted in pots for the first time.

Pots less than four inches wide.

Bedding plants or areas where bedding plants will be planted or transplanted within one year following application.

Ground covers until they are established and well rooted.

Do not apply Gallery 75 Dry Flowable to newly transplanted container plants, nursery stock, ground covers, ornamental bulbs, non-bearing fruit and nut trees and non-bearing vineyards or Christmas trees until soil or potting media has been settled by packing and irrigation or rainfall and no cracks are present or plant injury may occur.

When planting into a site treated with Gallery 75 Dry Flowable in the past 6 months, use untreated soil as fill around roots when replacing plants or injury may occur.

Repeat applications at 1.0 pound per acre or more of Gallery 75 Dry Flowable should not be made sooner than 60 days after a previous application of Gallery 75 Dry Flowable. Do not apply more than a total of 4.0 pounds per acre of Gallery 75 Dry Flowable within a 12 month period.

**BEST AVAILABLE COPY**

**Treatment of Plant Species Not Listed on the label  
or Gallery 75 Dry Flowable**

Users who wish to use Gallery 75 Dry Flowable on plant species not recommended on this label may determine the suitability for such uses by making a trial application of Gallery 75 Dry Flowable at a recommended rate to a small number of plants. Prior to using Gallery 75 Dry Flowable on a larger number of plants, the treated plants should be observed for any sign of herbicidal injury during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious to the target plant species. The user assumes the responsibility for any plant damage or other liability resulting from use of Gallery 75 Dry Flowable on plant species not recommended on this label.

**Note: Do not apply Gallery 75 Dry Flowable to the following plant species or injury may occur:**

- Ajuga spp. (bugle)
- Brassica spp. (mustard)
- Euonymus alatus 'compacta' (dwarf burning bush)
- Euphorbia spp. (spurge)
- Hydrangea spp. (hydrangea)
- Iberis spp. (candytuft)
- Juniperus horizontalis 'Prince of Wales' (Prince of Wales Juniper)
- Maleolucua quinquenervis (cayuput tree)
- Rhododendron carolinianum (Carolina rhododendron)
- Rhododendron catawbiense 'Roseum elegans' (Roseum elegans rhododendron)
- Sedum spp. (stonecrop)
- Yucca recurvifolia (green yucca)

**Tank Mixing**

Gallery 75 Dry Flowable may be tank mixed with Roundup herbicide or other registered postemergence herbicides to control existing unwanted vegetation and provide residual preemergence broadleaf weed control in ornamental areas. Gallery 75 Dry Flowable may also be tank mixed with Surflan A.S. herbicide and applied preemergence to provide broad spectrum control of annual grasses and broadleaf weeds in ornamental areas. Applied as directed, Gallery 75 Dry Flowable tank mixes will provide control of susceptible weed species listed on the respective labels. When using Gallery 75 Dry Flowable in tank mix combination with other products, read and follow all applicable use directions, precautions, and limitations on the respective product labels.

**Note: Do not apply sprays containing Roundup over the top of ornamental plants. Extreme care must be exercised to prevent contact of sprays containing Roundup with foliage or stems of turfgrasses, trees, shrubs, or other desirable vegetation since severe damage or death may result. If spraying Roundup in areas adjacent to desirable plants, use a shield to prevent spray from contacting foliage or stems of desirable plants.**

**Gallery 75 Dry Flowable may be used on the following established plant species:**

(Note: Limitations on recommended treatment methods)

**Trees**

Scientific Name/ Common Name	Recommended Treatment Method: C = Container Grown E = Field Grown
<i>Abies concolor</i> White fir	F
<i>Abutilon hybridum</i> Albus-flowering maple	F
Luteus-flowering maple	F
Roseus-flowering maple	F
Tangerine-flowering maple	F
Vesuvius red-flowering maple	F

<i>Acer ginnale</i> Flame maple	F
<i>Acer rubrum</i> Red maple	F
Red sunset maple	F
<i>Acer saccharinum</i> Silver maple	C, F
<i>Arecastrum romanoffianum</i> Queen palm	C, F
<i>Betula nigra</i> River birch	C, F
<i>Betula papyrifera</i> Paper birch	F
<i>Brachycton papuense</i> Bottle tree	C, F
<i>Bucida buxeras</i> Black olive	F
<i>Ceratonia siliqua</i> Carob	F
<i>Cercis canadensis</i> Redbud	C, F
<i>Chamaecyparis obtusa</i> Filicoides-tenspray cypress	F
Gracile-slender Hinoki cypress	F
<i>Chamaecyparis pisifera</i> Sawara-false cypress	F
Squamosa-moss cypress	F
<i>Chamaedorea cataractarum</i> Palm	F
<i>Chamaedorea costaricensis</i> Palm	F
<i>Chamaedorea elegans</i> Parlor palm	F
<i>Cornus florida</i> Cloud nine dogwood	C, F
Flowering dogwood	C, F
<i>Crategeus viridis</i> Green hawthorn	F
<i>Cupaniopsis anacardioides</i> Carrot wood	F
<i>Cupressus glabra</i> Arizona cypress	F
<i>Elaeagnus angustifolia</i> Russian olive	C, F
<i>Eucalyptus camaldulensis</i> Red gum eucalyptus	F
<i>Eucalyptus cinerea</i> Mealy eucalyptus	F
Silver dollar eucalyptus	F
<i>Eucalyptus microtheca</i> Coolibah tree	C, F
<i>Eucalyptus sideroxylon</i> Red ironbark eucalyptus	F
<i>Ficus benjamina</i> Ficus	C, F
Mini ficus	C, F
<i>Fraxinus ulex</i> Shamash	C, F
<i>Ginkgo biloba</i> Maidenhair tree	F
<i>Gleditsia triacanthos</i> Honey locust	F
Shademaster honey locust	F
<i>Heteromeles arbustiflora</i> Toyon	F
<i>Nitidum floridanum</i> Florida anise-tree	C, F
<i>Juniperus virginiana</i> Eastern red cedar	C, F
<i>Liquidambar styraciflua</i> American sweet gum	F

**BEST AVAILABLE COPY**

<i>Magnolia grandiflora</i>	
Southern magnolia	C, F
<i>Morus alba</i>	
White mulberry	F
<i>Musa aluminata</i>	
Banana	C, F
<i>Oxydendrum arboreum</i>	
So. wood	C, F
<i>Picea abies</i>	
Norway spruce	F
Pendula-weeping Norway spruce	F
Repens-spreading Norway spruce	F
<i>Picea glauca conica</i>	
Dwarf Alberta spruce	F
<i>Picea pungens</i>	
Dwarf globe blue spruce	C, F
Glauca-Colorado blue spruce	F
Hoopii-Hoop's blue spruce	F
Koster-Koster blue spruce	F
<i>Pinus aristata</i>	
Bristlecone pine	F
<i>Pinus canariensis</i>	
Canary Island pine	F
<i>Pinus contorta</i>	
Shore pine, beach pine	F
<i>Pinus edulis</i>	
Eldarica pine	C, F
<i>Pinus leucodermis</i>	
Bosnian pine	C, F
<i>Pinus mugo</i>	
Pumilio-shrubby swiss mountain pine	C, F
<i>Pinus nigra</i>	
Austrian black pine	C, F
<i>Pinus radiata</i>	
Monterey pine	F
<i>Pinus strobus</i>	
Eastern white pine	C, F
<i>Pinus sylvestris</i>	
Columnar Scotch pine	C, F
Scotch pine	C, F
<i>Pinus thunbergiana</i>	
Japanese black pine	F
<i>Platanus occidentalis</i>	
American sycamore	F
<i>Platanus racemosa</i>	
California sycamore	F
<i>Podocarpus</i> spp.	
Podocarpus	F
<i>Populus deltoides</i>	
Cottonwood	F
<i>Prosopis chilensis</i>	
Chilean mesquite	C, F
<i>Prunus yedoensis</i>	
Yoshino flowering cherry	F
<i>Quercus ilicifolia</i>	
Burr oak	F
<i>Quercus palustris</i>	
Pin oak	F
<i>Quercus phellos</i>	
Willow oak	C, F
<i>Quercus rubra</i>	
Red oak	C, F
<i>Quercus virginiana</i>	
Live oak	C, F
<i>Salix babylonica</i>	
Babylon weeping willow	F
Corkscrew willow	F
<i>Sequoiadendron giganteum</i>	
Giant sequoia	F
<i>Swietenia mahoganii</i>	
Mahogany	F

<i>Tabebuia caribbea</i>	
Yellow tab	F
<i>Taxus canadensis</i>	
Eastern hemlock	C, F
<i>Ulmus parvifolia</i>	
Chinese elm	F
<i>Washingtonia robusta</i>	
Mexican fan palm	F

### Ornamental Shrubs

Scientific Name/ Common Name	Recommended Treatment Method: C = Container Grown F = Field Grown
<i>Abelia grandiflora</i>	
Edward Goucher abelia	C, F
Glossy abelia	C, F
<i>Acacia abyssinica</i>	
Abyssinica acacia	C, F
<i>Acacia redolens</i>	
Prostrate acacia	F
<i>Acacia stenophylla</i>	
Shoestring acacia	C, F
<i>Acalypha wilkesiana</i>	
Copper leaf	C, F
<i>Acer ginnala</i>	
Amur maple	C, F
<i>Acer palmatum</i>	
Coral bark Japanese maple	C, F
Dwarf Japanese maple	C, F
<i>Agave americana</i>	
Century plant	F
<i>Astilbe chinensis</i>	
False spiraea	C, F
<i>Athyrium nipponicum</i>	
Japanese painted fern	C, F
<i>Baccharis pilularis</i>	
Coyotebush	F
<i>Berberis gladywensis</i>	
William Penn barberry	C, F
<i>Berberis mentorensis</i>	
Mentor barberry	C, F
<i>Berberis thunbergii</i>	
Atropurea-redleaf Japanese barberry	F
Aurea-golden Japanese barberry	F
Crimson pygmy barberry	C, F
Rose glow barberry	C, F
<i>Louisa-villae</i> spp.	
Barbara Karst	F
California gold	F
Pink pixie	C, F
Scarlet O'Hara	F
Temple fire	C, F
Texas dawn	F
<i>Buxus microphylla japonica</i>	
Japanese boxwood	C, F
<i>Buxus microphylla Koreana</i>	
Korean boxwood	F
<i>Callistemon citrinus</i>	
Lemon bottlebrush	F
<i>Callistemon viminalis</i>	
Weeping bottlebrush	C, F
<i>Calluna vulgaris</i>	
Spring torch scotch heather	C, F
<i>Camellia japonica</i>	
Camellia	C, F
<i>Cassia artemisioides</i>	
Feathery cassia	C, F

BEST AVAILABLE COPY

<i>Caenothur</i> spp.			
Wild lilac	F		
<i>Cephalotaxus drupacea</i>			
Plum yew	C, F		
<i>Ceraeum tomentosum</i>			
Snor-in-summer	C, F		
<i>Chamaecyparis obtusa</i>			
Kostari cypress	F		
Nana-dwarf Hinoki cypress	F		
Torulosa cypress	F		
<i>Chamaecyparis pisifera</i>			
Fillera-thread cypress	F		
<i>Chrysalidocarpus nuscens</i>			
Areca palm	F		
<i>Clethra alnifolia</i>			
Summe: sweet	C, F		
<i>Claytonia japonica</i>			
Japanese claytonia	C, F		
<i>Cornus alba</i>			
Sibirica-Siberian dogwood	F		
<i>Cornus stolonifera</i>			
Baileyi-red-osier dogwood	F		
Flaviramea-yellowtwig dogwood	F		
<i>Cotinus coggygria</i>			
Royal purple smoke tree	C, F		
<i>Cotinus dammeri</i>			
Coral beauty smoke tree	C, F		
Eichholz smoke tree	C, F		
<i>Cotoneaster adpressus</i>			
Praecox-early cotoneaster	F		
<i>Cotoneaster apiculatus</i>			
Cranberry cotoneaster	C, F		
<i>Cotoneaster congestus</i>			
Pyrenees cotoneaster	F		
<i>Cotoneaster dammeri</i>			
Bearberry cotoneaster	C, F		
<i>Cotoneaster himalayana</i>			
Himalayan cotoneaster	F		
<i>Cotoneaster horizontalis</i>			
Rock cotoneaster	C, F		
<i>Cycas revoluta</i>			
Sago palm	C, F		
<i>Cytisus praecox</i>			
Hollandia-warminster broom	F		
<i>Cytisus scoparius</i>			
Lane-Scotch broom	F		
<i>Daphne odora</i>			
Fragrant daphne	C, F		
<i>Deutzia crenata</i>			
Nakiana-dwarf deutzia	F		
<i>Deutzia gracilis</i>			
Slender gracilis	F		
<i>Dodonaea viscosa</i>			
Hopsad bush	F		
<i>Eleagnus pungens</i>			
Fruitland silver berry	C, F		
<i>Erica cinerea</i>			
Purple bell heather	C, F		
<i>Erica x darleyana</i>			
Mediterranean pink heather	C, F		
<i>Erica vagans</i>			
Cornish heather	C, F		
<i>Euonymus fortunei</i>			
Canadale gold euonymus	F		
Emerald gold euonymus	F		
Sunspot euonymus	C, F		
<i>Euonymus japonica</i>			
Silver king euonymus	F		
Variegated evergreen euonymus	C, F		
<i>Euonymus histschovica</i>			
Spreading euonymus	F		
<i>Forsythia intermedia</i>			
Border forsythia	C, F		
<i>Gardenia jasminoides</i>			
August beauty gardenia	C, F		
Gardenia	C, F		
Radiosa gardenia	C, F		
<i>Gaultheria shallon</i>			
Santal/lemon leaf	C, F		
<i>Geleum sempervirens</i>			
Carolina jessamine	C, F		
<i>Genista pilosa</i>			
Woodwaxen	F		
<i>Hibiscus rosa-sinensis</i>			
Rose Estay-hibiscus	C, F		
<i>Hibiscus syriacus</i>			
Rose of Sharon-heart	F		
Rose of Sharon-red bird	F		
Rose of Sharon-woodbridge	F		
<i>Ilex aquifolium</i>			
Balkans holly	F		
Gold coast holly	F		
<i>Ilex aquiparvifolia</i>			
San Jose holly	C, F		
<i>Ilex attenuata</i>			
Savannah holly	C, F		
<i>Ilex cornuta</i>			
Burford holly	C, F		
Dwarf Burford holly	F		
Needle point holly	C, F		
<i>Ilex crenata</i>			
Compacta-dwarf Japanese holly	C, F		
Convexa holly	C, F		
Dwarf Chinese holly	C, F		
Green luster holly	C, F		
Helleri-Heller's Japanese holly	C, F		
Hetzii's Japanese holly	C, F		
Stokesii Japanese holly	C, F		
<i>Ilex glabra</i>			
Compacta-compact inkberry holly	C, F		
Nordica-inkberry holly	C, F		
<i>Ilex meserveae</i>			
Blue girl holly	F		
Blue boy holly	F		
Ebony magic holly	F		
<i>Ilex vomitoria</i>			
Nana-dwarf yaupon holly	C, F		
Pendula-weeping yaupon holly	C, F		
yaupon holly	C, F		
<i>Illicium anisatum</i>			
Mystery gardenia	C, F		
<i>Itea ilicifolia</i>			
Henry Garnet holly leaf sweetspire	C, F		
<i>Ixora collina</i>			
Ixora	C, F		
<i>Juniperus chinensis</i>			
Torulosa-hollywood juniper	C, F		
Media-old gold juniper	C, F		
Pfitzerana-pfitzer juniper	C, F		
Pfitzerana glauca-blue juniper	C, F		
Sea green juniper	F		
<i>Juniperus conferta</i>			
Emerald sea shore juniper	C, F		
Shore juniper	C, F		
<i>Juniperus horizontalis</i>			
Plumosa-Andorra juniper	C, F		
Bar Harbor juniper	C, F		
Blue rug juniper	C, F		
Creeping juniper	C, F		
Dwarf Andorra juniper	C, F		
Huntington blue juniper	C, F		
Witons-blue carpet juniper	C, F		

**BEST AVAILABLE COPY**

<i>Juniperus procumbens</i>		<i>Pieris japonica</i>	
Nana-dwarf Japanese garden juniper	C, F	Lily-of-the-valley	C, F
<i>Juniperus prostrata</i>		Mountain fire lily-of-the-valley	C, F
Prostrata juniper	C, F	Snowdrift lily-of-the-valley	C, F
<i>Juniperus sabina</i>		Temple bells lily-of-the-valley	C, F
Broadmoor juniper	F	Valley rose lily-of-the-valley	C, F
Foemina-Hicks juniper	C, F	Valley valentine lily-of-the-valley	C, F
Savin juniper	C, F	<i>Pieris japonica x forestii</i>	
Tamariscifolia-Tam juniper	F	Forest flame lily-of-the-valley	C, F
<i>Juniperus scopulorum</i>		<i>Pinus mugo</i>	
Emerald green juniper	F	Mugo-mugho pine	C, F
<i>Juniperus squamata</i>		<i>Pitiosporum tobira</i>	
Blue juniper	F	Wheeler's dwarf pitiosporum	C, F
Blue star juniper	F	Green pitiosporum	C, F
Parsoni juniper	C, F	<i>Plumbago auriculata</i>	
<i>Kalmia latifolia</i>		Blue cape plumbago	F
Mountain laurel	C, F	<i>Plumbago capensis</i>	
<i>Lagerstroemia indica</i>		Plumbago	C, F
Crape myrtle	C, F	<i>Podocarpus macrophyllus</i>	
<i>Lantana spp.</i>		Yewpine	C, F
Lantana	C, F	<i>Polystichum polyblepharum</i>	
<i>Leucothoe axillaris</i>		Tassel fern	C, F
Coast leucothoe	C, F	<i>Potentilla fragiformis</i>	
<i>Leucothoe fontanesiana</i>		Cinquefoil	F
Drooping leucothoe	C, F	<i>Potentilla fruticosa</i>	
<i>Ligustrum lucidum</i>		Cinquefoil	C, F
Glossy privet	C, F	Gold drop potentilla	F
<i>Ligustrum ovalifolium</i>		Goldfinger potentilla	C, F
California privet	F	Red ace potentilla	C, F
<i>Ligustrum japonicum</i>		Sunset potentilla	C, F
Japanese privet	C, F	Tangerine potentilla	C, F
Wax ligustrum	C, F	<i>Potentilla verna</i>	
Yellow tip ligustrum	C, F	Spring cinquefoil	C, F
<i>Ligustrum taxanum</i>		<i>Prunus glandulosa</i>	
Howard privet	F	Dwarf pink flowering almond	C, F
Wax leaf privet	C, F	<i>Pyracantha fortuneana</i>	
<i>Ligustrum vicaryi</i>		Lalendal Monrovia pyracantha	F
Vicary golden privet	F	Monon pyracantha	F
<i>Ligustrum vulgare</i>		Red ell hybrid pyracantha	F
Lodense privet	C, F	Rutgers hybrid pyracantha	C, F
<i>Livistone chinensis</i>		Santa Cruz pyracantha	F
Chinese fountain palm	F	Victory pyracantha	F
<i>Lonicera periclymenum</i>		<i>Raphiolepis indica</i>	
Flowering woodbine	F	Charisma-Monruce raphiolepis	F
Serotina woodbine	F	Enchantress-Moness raphiolepis	F
<i>Lonicera sempervirens</i>		Infra hawthorn	C, F
Trumpet honeysuckle	F	Springtime-Monme raphiolepis	F
<i>Mahonia bealei</i>		<i>Raphiolepis ovata</i>	
Leather leaf mahonia	C, F	Roundleaf raphiolepis	F
<i>Mahonia repens</i>		<i>Rhododendron ssp. (including azalea)</i>	
Creeping mahonia	C, F	<b>Azalea</b>	
<i>Myrica cerifera</i>		<i>Rhododendron calendulaceum</i>	
Wax myrtle	C, F	Flame azalea	F
<i>Nandina domestica</i>		Cannon's double azalea	C, F
Compacta-dwarf heavenly bamboo	C, F	Golden flare azalea	C, F
Harbour dwarf-heavenly bamboo	C, F	Klondike azalea	C, F
Heavenly bamboo	C, F	<i>Rhododendron indica</i>	
Nana compacta-heavenly bamboo	C, F	Armoes azalea	C, F
Nana purpurea-heavenly bamboo	C, F	Waucabusa azalea	C, F
Woods dwarf-heavenly bamboo	C, F	<i>Rhododendron kaempferi</i>	
<i>Nerium oleander</i>		Blue danube azalea	C, F
Hardy red oleander	F	<i>Rhododendron keruma</i>	
Oleander	C, F	Coral bells azalea	C, F
Ruby lace oleander	F	Hino pink azalea	C, F
<i>Osmantbus fortunei</i>		Mildred azalea	C, F
Fortunes osmanthus	C, F	Snow azalea	C, F
<i>Phoenix rostrata</i>		<i>Rhododendron obtusum</i>	
Pigmy date palm	C, F	Hino-crimson azalea	C, F
<i>Photinia fraseri</i>		<i>Rhododendron satsumi</i>	
Fraser's photinia	C, F	Gumpo pink azalea	C, F
		Higasa azalea	F
		Reijn azalea	C, F



<i>Rhododendron</i> spp. hybrids		<i>Syringa rohaniensis</i>	
Carron azalea	C, F	Chinese lilac	C, F
Fashion azalea	C, F	<i>Syringa vulgaris</i>	
Gerard Christina azalea	F	Common lilac	F
Girard Roberts azalea	F	<i>Taxus cuspidata</i>	
Helmut Vogel azalea	F	Japanese yew	F
Hersey red azalea	F	<i>Tecoma capensis</i>	
Hot shot azalea	C, F	Cape honeysuckle	C, F
Hume azalea	F	<i>Ternstroemia gymnanthera</i>	C, F
Inga azalea	F	<i>Thuja occidentalis</i>	
Irene Koster azalea	C, F	Emerald arborvitae	F
President Clay azalea	C, F	Globosa-globe arborvitae	C, F
Tradition azalea	C, F	Little giant-dwarf arborvitae	F
<b>Rhododendrons</b>		Nigra-dark American arborvitae	F
<i>Rhododendron campylocarpum</i>		Pyramidalis-pyramid arborvitae	F
Butterfly rhododendron	F	Pinegold arborvitae	F
<i>Rhododendron carolinianum</i> x <i>deurium</i>		Tachy arborvitae	F
PJM rhododendron	F	<i>Thuja orientalis</i>	
<i>Rhododendron catawbiense</i>		Aurea nano-dwarf golden arborvitae	F
Catawba rhododendron	C, F	Minima glauca-dwarf arborvitae	F
Catawba album rhododendron	C, F	<i>Veitchia marilli</i>	
Lord Roberts rhododendron	C, F	Christmas palm	F
Rocket rhododendron	C, F	<i>Viburnum bochantense</i>	
<i>Rhododendron caucasicum</i> x <i>ponticum</i>		Pink dawn viburnum	C, F
Cunningham White rhododendron	F	<i>Viburnum davidii</i>	
<i>Rhododendron forrestii</i> x <i>griersonianum</i>		David viburnum	F
Elizabeth rhododendron	C, F	<i>Viburnum japonicum</i>	
<i>Rhododendron forrestii repens</i>		Viburnum	F
Gomer Waterer rhododendron	C, F	<i>Viburnum lentana</i>	
<i>Rhododendron griffithianum</i>		Wayfaring tree	F
Jean Marie rhododendron	F	<i>Viburnum opulus sterile</i>	
<i>Rhododendron impeditum</i>		Common snowball viburnum	F
Rhododendron	C, F	<i>Viburnum plicatum tomentosum</i>	
<i>Rhododendron maximum</i>		Doublefile viburnum	F
Rhodie max (rosebay)	F	<i>Viburnum x pragensis</i>	
<i>Rhododendron mucronulatum</i>		Viburnum	F
Rhododendron	F	<i>Viburnum setigerum</i>	
<i>Rhododendron ponticum</i>		Tea viburnum	F
Chionodoxa rhododendron	F	<i>Viburnum tinus compactum</i>	
Daphnoides rhododendron	C, F	Spring bouquet viburnum	F
<i>Rhododendron racemosum</i>		<i>Viburnum trilobum</i>	
Dwarf scarlet wonder rhododendron	C, F	Cranberry bush	C, F
Tribly rhododendron	C, F	<i>Viburnum trilobum compactum</i>	
Unique rhododendron	C, F	Dwarf cranberry bush	C, F
Vulcan rhododendron	C, F	<i>Weigela florida</i>	
<i>Rhododendron sassthipiatim</i> x <i>carolinianum</i>		Bristol ruby weigela	F
Ramapo rhododendron	F	Java red weigela	F
<i>Rhododendron hybrid</i> spp.		Minuet weigela	F
America rhododendron	C, F	<i>Xylocoma congestum</i>	
English Roseum rhododendron	F	Xylocoma	F
Nova Zambia rhododendron	C, F	<i>Yucca filamentosa</i>	
Scintillation rhododendron	C, F	Yucca	C, F
<i>Rhus lancea</i>			
Africa sumac	C, F		
<i>Rosa rugosa</i>			
Remanes rose	F		
<i>Rosmarinus officinalis</i>			
Rosemary	F		
<i>Stimmiis japonica</i>			
Japanese stimmiis	C, F		
<i>Stimmiis revesiana</i>			
Reeve's stimmiis	C, F		
<i>Spiraea bumalda</i>			
Anthony Waterer spiraea	C, F		
<i>Spiraea japonica</i>			
Dolichia spiraea	C, F		
Japanese alpine spiraea	C, F		
Shirobana spiraea	C, F		
<i>Spiraea vanhouttei</i>			
Bridal wreath	C, F		

BEST AVAILABLE COPY

102811

**Ground Covers**

Scientific Name/ Common Name	Recommended Treatment Method: C = Container Grown F = Field Grown
<i>Ammophila breviflorus</i>	C, F
Beadgrass	
<i>Arctostaphylos californica</i>	F
Cape weed	
Carex spp.	C, F
Variegated carex	
<i>Carpenteria edulis</i>	F
Largeleaf iceplant	
<i>Conoclinium subterminale</i>	C, F
Pampas grass	
<i>Dactyloctenium aegyptium</i>	F
White iceplant	
<i>Decaspermum caespitosum</i>	C, F
Decaspermia	
<i>Dioscorea oppositifolia</i>	F
Trailing rose iceplant	
<i>Drosera rotundifolia</i>	C, F
Iceplant	
<i>Festuca ovina glauca</i>	C, F
Blue fescue	
Gazania spp.	
Gazania	C, F
<i>Halenia laetevirens</i>	C, F
Golden halimolobos	
<i>Hedera cuneata</i>	F
Algerian ivy	
<i>Hedera helix</i>	C, F
English ivy	
<i>Hemipocallis</i> spp.	C, F
Daylily	
<i>Hosta lancifolia</i>	C
Narrow leafed plantain lily	
<i>Hosta lancifolia</i>	C, F
Abso-marginata hosta	
<i>Hypericum</i> spp.	F
St. Johnswort	
<i>Jasminum nudum</i>	C, F
Angelwing jasmine	
<i>Lampranthus spectabilis</i>	F
Trailing iceplant	
<i>Liriope glauca</i>	F
White lily turf	
<i>Liriope muscari</i>	C, F
Big blue lily turf	
Black beauty lily turf	C, F
Majestic lily turf	C, F
Morroe white lily turf	C, F
Silvery sunproof lily turf	C, F
Variegated liriope lily turf	C, F
<i>Liriope spicata</i>	C, F
Greentwining lily turf	
<i>Lonicera japonica</i>	F
Japanese honeysuckle	
<i>Macrorhizus sinensis</i>	C, F
Edulis grass	
<i>Ophiopogon japonicus</i>	C, F
Dwarf Mondo grass	C, F
Mondo grass	F
<i>Osteospermum frutescens</i>	C, F
Trailing African daisy	
<i>Pachyzanthes terminalis</i>	C, F
Japanese spurge	
<i>Perisetum alopecuroides</i>	C, F
Fountain grass	

Recommended  
Treatment Method:  
C = Container Grown  
F = Field Grown

<i>Phalaris amabilis</i>	C, F
Ribbon grass	
<i>Trachypogon sativum</i>	C, F
Asian juncus	
<i>Vriesea minor</i>	F
Dwarf periwinkle	

**Ornamental Bulbs**

Galery 75 Dry Flowable may be applied for control of susceptible annual weeds in ornamental bulbs, e.g., bulbs in the soil surface, iris and tulip. Apply Galery 75 Dry Flowable to the soil surface 2 to 4 weeks after planting, but prior to the emergence of annual weeds. Galery 75 Dry Flowable may also be applied following bulb emergence but prior to flowering and after flowering. For late planted bulbs, apply Galery 75 Dry Flowable again in late winter or early spring to weed-free soil surfaces.

**Special Use Precautions:**

Do not apply to bulb plants that have emerged to a height greater than 3/4 inch.  
Do not apply to gladiolus prior to emergence or if corms are less than one inch in diameter.  
Do not apply to bulbs while they are flowering.

**Non-bearing† Fruit and Nut Trees and Non-bearing† Vineyards**

Scientific Name/ Common Name	Recommended Treatment Method: C = Container Grown F = Field Grown
almond	F
apple	F
apricot	F
avocado	F
blackberry	F
blueberry	F
boysenberry	F
cherry, sour	F
cherry, sweet	F
currant	F
dateberry	F
elderberry	F
fig	F
hibert	F
gooseberry	F
grape, American	F
grape, European	F
grapefruit	F
kiwi	F
lemon	F
loganberry	F
macadamia nut	F
nectarine	F
olive	F
orange	F
peach	F
pear	F
pecan	F
plum	F
prune	F
ponygranate	F
raspberry	F
walnut, black	F
walnut, English	F

† Non-bearing fruit and nut trees and non-bearing vineyards are defined as plants which will not bear fruit for at least one year after treatment.

## Christmas Tree Plantations

### Gallery 75 Dry Flowable Alone

Apply Gallery 75 Dry Flowable as a directed spray to the soil surface or as an overtop spray to established plantings of field grown Christmas tree species listed in this label. Do not apply to seedbeds or seeding transplant beds. Apply only to established plantings. Established plants are defined as those that have been transplanted into their final growing location for a sufficient period of time to allow the soil to be firmly settled around the roots from packing and rainfall or irrigation. Follow all instructions provided in the "General Information" section of this label.

### Tank Mix Combinations

Tank mix combinations of Gallery 75 Dry Flowable plus other labeled herbicides may be used as directed or overtop sprays in established Christmas tree plantings. When applied according to use directions, these tank mixes will provide control of susceptible weed species listed on the respective product labels. Refer to tank mix product labels for specific use directions, precautions and limitations before use.

**Gallery 75 Dry Flowable plus Roundup:** Apply tank mix combinations of Gallery 75 Dry Flowable plus Roundup only as directed sprays in Christmas tree plantings. When applied according to use directions, Gallery 75 Dry Flowable plus Roundup herbicide will provide postemergence control of susceptible weed species listed on the label for Roundup and residual preemergence control of susceptible weed species listed on the label for Gallery 75 Dry Flowable. Refer to the label for Roundup for specific use directions, precautions and limitations before use.

### Precautions:

Do not apply sprays containing Roundup over the top of Christmas tree plantings.

Extreme care must be exercised to prevent contact of sprays containing Roundup with foliage or stems of Christmas trees or other desirable plants or severe damage or death may result.

## Non-cropland Areas

Gallery 75 Dry Flowable is recommended as a preemergence herbicide for control of certain broadleaf weeds in noncropland areas, including industrial sites, utility substations, highway guard rails, sign posts and delineators.

Apply Gallery 75 Dry Flowable anytime prior to germination of target weeds. Areas to be treated should be free of established weeds or existing weeds should be controlled with postemergence herbicides.

Refer to the "General Information" section prior to using this product on non-cropland.

Gallery 75 Dry Flowable is compatible and can be tank mixed with other herbicides registered for use on non-cropland. Applied as directed, tank mixes containing Gallery 75 Dry Flowable will provide control of susceptible weed species listed on the respective labels. All directions, precautions and limitations on the respective product labels apply to the tank mix use.

## Warranty Disclaimer

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. DowElanco makes no other express or implied warranty of merchantability or fitness for a particular purpose or any other express or implied warranty.

## Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application or other factors, all of which are beyond the control of DowElanco or the seller. All such risks shall be assumed by Buyer.

## Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at DowElanco's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of product used.

DowElanco shall not be liable for losses or damages resulting from handling or use of this product unless DowElanco is promptly notified of such loss or damage in writing. In no case shall DowElanco be liable for consequential or incidental damages or losses.

The terms of the "Warranty Disclaimer" above and this "Limitation of Remedies" cannot be varied by any written or verbal statements or agreements. No employee or sales agent of DowElanco or the seller is authorized to vary or exceed the terms of the "Warranty Disclaimer" or this "Limitation of Remedies" in any manner.

\*Trademark of DowElanco  
DowElanco • Indianapolis, IN 46268, U.S.A.

LABEL CODE 113-75-002    DATE CODE 992  
EPA APPROVAL 08/18/92

### Revisions include:

- 1) Revised weeds controlled list (excluding grass weeds) to match Snapshot 2.5 TG label.
- 2) Revised plant species which may be treated to match Snapshot 2.5 TG label.
- 3) Tall fescue removed from "Established Cool Season Turf Grasses" list.
- 4) Buffelgrass (*Buchloe dactyloides*) added to "Established Warm Season Turf Grasses" list.
- 5) Added use on ornamental bulbs.
- 6) Added use on Christmas tree plantations.
- 7) Added instructions for use of container cap as dose measuring device.

**BEST AVAILABLE COPY**