

(Container Label)

Turf Fertilizer - Contains Balan* 1.25%

A granular fertilizer containing Balan for the preemergence control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual weed grasses in established turfgrass.

Active Ingredients:

benefin: <i>N</i> -butyl- <i>N</i> -ethyl- <i>a,a,a</i> -trifluoro-2,6-dinitro- <i>p</i> -toluidine	1.25%
Inert Ingredients	98.75%
Total	100.00%

Contains 0.46 Pounds of Active Ingredient Per 50 Pounds Bag

Guaranteed Fertilizer Analysis	
Total Nitrogen (N)	18.0%
Available Phosphoric Acid (P ₂ O ₅)	5.0%
Soluble Potash (K ₂ O)	9.0%

Keep Out of Reach of Children

WARNING

AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See back or container/side panel for additional precautionary statements.

Notice: Read the entire label. Use only according to label directions. Before buying this product, read Limited Warranty and Disclaimer Section elsewhere on this label.

In case of emergency endangering health or the environment involving this product, call collect 517-636-4400.

Do Not Ship or Store with Food, Feeds, Drugs, or Clothing.

EPA Reg. No. 62719-148

EPA Est. No. _____

*Trademark of DowElanco
DowElanco • Indianapolis, Indiana 46268 U.S.A.

Herbicide
Net Weight - 50 lb

BEST AVAILABLE COPY

62719-148



Precautionary Statements

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

WARNING

Harmful If Swallowed, Inhaled Or Absorbed Through The Skin

Avoid breathing dust and contact with skin or clothing. The granular formulation may cause skin sensitization reactions in certain individuals. Use protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

First Aid

If swallowed: Call physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or, if available, by administering 1 to 2 tablespoons of syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation develops.

If in eyes: Flush with plenty of water. Call a physician if irritation develops.

If inhaled: If breathing difficulty occurs, move individual to fresh air and get medical attention.

Environmental Hazards

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read all directions carefully before applying.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General Information

This product is a granular fertilizer containing Balan, a preemergence herbicide, which is applied to established turfgrasses for the control of the winter annual grass: annual bluegrass (*Poa annua*); and summer annual grasses: smooth and hairy crabgrass, goosegrass (silver crabgrass or crowfootgrass), barnyardgrass (watergrass) and green and yellow foxtail. Turf fertilizer containing Balan controls these weeds by killing their seeds as they germinate. Apply this product before these annual weeds germinate since it does not control established weeds. This product will not harm most nearby established ornamentals.

BEST AVAILABLE COPY

DowElanco's Turf Fertilizer containing Balan can be used on the following established cool-and warm-season turfgrasses:

Established Cool-Season Turfgrasses

- Bentgrass fairways
- Kentucky bluegrass
- Perennial ryegrass
- Fescues

Established Warm-Season Turfgrasses

- Bahiagrass
- Bermudagrass
- Centipedegrass
- St. Augustinegrass
- Tall fescue
- Zoysiagrass

General Use Precautions

Do not apply this product to golf course putting greens, dichondra, or newly sprigged areas of bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass. Any cultural practices that disturb the soil, such as aeration or verticutting, should be done prior to application.

Application Directions

Rates and Frequency of Application: This product should be applied one to two weeks prior to the anticipated germination of the target annual grass weeds. If application time does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control may be erratic or poor. Optimum weed control performance will be obtained if treated areas are irrigated soon after application.

Broadcast Applications Rates:

	<u>Cool-Season Turfgrasses</u>	<u>Warm-Season Turfgrasses</u>
Summer Annual Grass Weeds	160 lb/acre (3.7 lb/1000 sq ft)	240 lb/acre (5.5 lb/1000 sq ft)
Winter Annual Grass Weeds	160 lb/acre	240 lb/acre

In areas with extended growing seasons or where desired, a second application at the recommended rate can be made 8 to 10 weeks after following the initial application on cool-season turfgrasses and 10 to 12 weeks following the initial application on warm season turfgrasses.

Reseeding

Chemicals that effectively control annual weed species may also effect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after an application rate of 160 pounds per acre and 12 to 16 weeks after an application rate of 240 pounds per acre.

Application Equipment

Applications should be made using a properly calibrated granular fertilizer spreader which will uniformly distribute the product over the target area. Avoid skips and excessive overlaps as poor weed control or turfgrass injury may occur.

[Editor's note: Subregistrant may insert appropriate Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies statements.]

©Copyright 1994 by DowElanco

BEST AVAILABLE COPY