

62719-150

10-2-2001

1/5

Turf Fertilizer - Contains Team* 1.15%

A Granular Fertilizer Containing Team for the Preemergence Control of Crabgrass, Foxtail, Goosegrass, *Poa annua* and Other Annual Weed Grasses in Established Turfgrasses.

Active Ingredients:

benefin: <i>N</i> -butyl- <i>N</i> -ethyl-trifluoro-2,6-dinitro- <i>p</i> -toluidine.....	0.76%
trifluralin: -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine	0.39%
Inert Ingredients.....	98.85%
Total.....	100.00%

Guaranteed Fertilizer Analysis†	
Total Nitrogen (N)	XX.X%
Available Phosphoric Acid (P ₂ O ₅)	XX.X%
Soluble Potash (K ₂ O)	XX.X%

[†Editor's note: See subregistrant label for specific fertilizer analysis of the fertilizer blend used.]
Contains 0.575 pounds active ingredient per 50 pound bag.

Keep Out of Reach of Children

WARNING

See back or container/side panel for additional precautionary statements.

Notice: Read the entire label. Use only according to label directions. Before buying using this product, read Limited Warranty and Disclaimer Section elsewhere on this label, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.



Do Not Ship or Store with Food, Feeds, Drugs, or Clothing.

EPA Reg. No. 62719-150

EPA Est. No. _____

*Trademark of Dow AgroSciences LLC
Dow AgroSciences • Indianapolis, Indiana 46268 U.S.A.

Herbicide
Net Weight - 50 Pounds



Precautionary Statements

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

WARNING

Causes substantial but temporary eye injury • Harmful if absorbed through skin •

Do not get in eyes or on clothing. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

Personal Protective Equipment:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks
- Protective eyewear

User Safety Requirements: Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

- Wear goggles or face shield.
- Users should wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should removed PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: ~~Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention. Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.~~

If swallowed: ~~Call physician or Poison Control Center. Do not induce vomiting. Drink promptly a large quantity of milk, egg white, gelatin solution, or if these are not available, large quantities of water. Avoid alcohol. Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.~~

If on skin or clothing: ~~Wash with plenty of soap and water. Get medical attention. Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call poison control center or doctor for treatment advice.~~

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner which will directly expose

canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

General Information

This turf fertilizer product contains Team*, a preemergence herbicide for application to established turfgrasses for the control of the winter annual grass: annual bluegrass (*Poa annua*); and summer annual grasses: smooth and hairy crabgrass, goosegrass (silver crabgrass or crowfootgrass), barnyardgrass (watergrass) and green and yellow foxtail. This product also provides partial control of prostrate spurge (*Euphorbia humistrata*), spotted spurge (*Euphorbia maculata*), creeping spurge (*Euphorbia serpens*), creeping woodsorrel (*Oxalis corniculata*) and yellow woodsorrel (*Oxalis stricta*).

General Use Precautions and Restrictions

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Application Restrictions: Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Entry Restrictions: Do not enter or allow others to enter treated areas until dusts have settled. If incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

Reseeding: Do not reseed treated areas for minimum of 8 weeks after a single application of 174 pounds per acre or 12 to 16 weeks after a single application of 260 pounds per acre. For split applications, do not reseed treated areas for 8 to 10 weeks after the last 130 pounds per acre rate or 10 to 12 weeks after the last 174 pounds per acre rate.

Do not apply this product to golf course putting greens, dichondra, or newly sprigged areas of bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass. Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application.

This Product is Recommended for Use on the Following Established Warm and Cool Season Perennial Turfgrass Species:

Established Cool Season Turfgrasses

- Bentgrass fairways
- Kentucky bluegrass
- Perennial ryegrass
- Fescues

Established Warm Season Turfgrasses

- Bahiagrass
- Bermudagrass
- Centipedegrass
- St. Augustinegrass
- Tall fescue
- Zoysiagrass

4/5

Application

This product should be applied one to two weeks prior to the anticipated germination of the target annual weeds. Apply using a properly calibrated granular fertilizer spreader which will uniformly distribute the product over the target area. Avoid skips and excessive overlaps as poor weed control or turfgrass injury may occur. Optimum weed control performance will be obtained if treated areas are irrigated soon after application. This product will not harm most nearby ornamentals.

In areas with extended growing seasons or where desired, a second application at the recommended rate can be made 8 to 10 weeks following the initial application on cool season turfgrasses and 10 to 12 weeks following the initial application on warm season turfgrasses.

Broadcast Application Rates:

	Cool Season Turfgrasses	Warm Season Turfgrasses
Summer Annual Grass Weeds (Spring Application)	130 to 260 lb/acre 3 to 6 lb/1000 sq ft	260 lb/acre 6 lb/1000 sq ft
Winter Annual Grass Weeds (Fall application)	130 to 174 lb/acre 3 to 4 lb/1000 sq ft	260 lb/acre 6 lb/1000 sq ft

Split Applications for Extended Control and Lawn Care Operator Programs:

As an alternative to single application programs, this product may also be applied in split applications. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time.

Apply this product at 130 pounds plus 130 pounds per acre for Cool Season Turfgrasses and in the northern half of the Warm Season Turfgrass area of the U.S. or 174 pounds plus 174 pounds per acre in the southern half of the Warm Season Turfgrasses area of the U.S. The second application should be made 8 to 10 weeks following the 130 pounds per acre rate and 10 to 12 weeks following the 174 pounds per acre rate.

Spreader Settings

[Editor's note: Subregistrant label may include a table of recommended spreader equipment and spreader settings for the fertilizer blend used.]

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

5/5

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of ~~Dow AgroSciences~~ or the Seller. All such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

*Trademark of Dow AgroSciences
EPA-Accepted: 03/31/99