

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

MAY 1 2008

Ms. Kimberly S. Gilbert Product Registration Dow AgroScience LLC 9330 Zionsville Road Indianapolis, IN 46268

SUBJECT: Application for Pesticide Notification (PRN 98-10)

Request General Label Change & Optional Label Claims

EPA Reg. No. 62719-150

Application Dated April 9, 2008

Dear Registrant:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 04/09/08 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

Linda Arrington

Notifications & Minor Formulations Team Leader Registration Division (7505P)

Office of Pesticide Programs

308/2E April 9, 2008



Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U. S. Environmental Protection Agency
One Potomac Yard
2777 S. Crystal Drive
Arlington, VA 22202

TURF FERTILIZER – CONTAINS TEAM 1.15% (A.I. BENEFIN, TRIFLURALIN) EPA REGISTRATION NUMBER: 62719-150 NOTIFICATION OF MINOR LABEL CHANGE PER PR NOTICE 98-10

Enclosed please find labeling for the notification action of Turf Fertilizer – Contains Team[®] 1.15% herbicide. The following changes have been made by notification:

- 1. Updated trademark symbol throughout.
- 2. Added to list of optional label claims.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Contents of Submission

- Transmittal document (this letter)
- Application for Pesticide, EPA Form 8570-1
- Label entitled Turf Fertilizer-Contains Team 1.15% (P4A/Turf Fert Contains Team 1.15%/Notif/04-09-08)
 (19 Pages plus Registration Notes) (5 Copies)

If you require further information, please contact Cindy Loy, Regulatory Specialist at (317) 337-4655 or Paula McKinnies, Registration Assistant for this product, at (317) 337-4679.

Sincerely,

Kimberly S. Gilbert Regulatory Leader (317) 337-4685

(317) 337-4649 (FAX)

Enclosures

Trademark of Dow AgroSciences LLC

Please read instructions on reverse before completing form.	Form Approved	3/3 . OMB No. 2070-0060, Approval expires 2-28-9
SEPA Environmental Protect Washington, DC	tion Agency	Registration Amendment Other
Applica	tion for Pesticide - Section	I
1. Company/Product Number Dow AgroSciences LLC/62719-150	2. EPA Product Menager Joanne I. Miller	3. Proposed Classification ✓ None Restricted
 Company/Product (Name) Dow AgroSciences LLC/Turf Fert-Contains Team 1 	.15% PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268 Check if this is a new address		In accordance with FIFRA Section 3(c)(3) ilar or identical in composition and labeling
	Section - II	
Amendment - Explain below. Resubmission in response to Agency letter dated	8-10 and EPA regulations at 40 CFR 152.46 I understand that it is a violation of 18 U.S. nt with the terms of PR Notice 98-10 and 40	ed ation. slow. s, and no other changes have been made to the C. Sec. 1001 to willfully make any false statement to
	Section - III	
1. Material This Product Will Be Packaged In: Child-Resistant Packaging Yes No * Certification must be submitted 3. Location of Net Contents Information 1. Material This Product Will Be Packaging Unit Packaging No. per Unit Packaging wgt. A. Size(s)		2. Type of Container Metal Plastic Glass Paper Other (Specify) cation of Label Directions
6. Manner in Which Label is Affixed to Product	hograph Other	
Ste	per glued enciled	
	Section - IV	
Contact Point <i>(Complete items directly below for identific)</i> Name Kimberly S. Gilbert	Title Regulatory Manager	Telephone No. (Include Area Code) (317) 337-4685 (fax: 317-337-4649)

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowlingly false or misleading statement may be punishable by fine or imprisonment cross-

3. Title

5. Date

Regulatory Manager

April 9, 2008

both under applicable law.

Kimberly S. Gilbert (ksgibert@dow.com)

2. Signature

4. Typed Name

6. Date Application

(Stamped)

Received

Turf Fertilizer - Contains Team® 1.15%

EPA Reg. No. 62719-150

Registration Notes:

Source label text based on EPA accepted text dated January 15, 2008 and notification coded "P4A / Turf Fert Contains Team 1.15% / Notif / 01-30-08."

Changes by notification:

- 1. Updated trademark symbol throughout.
- 2. Added to list of optional label claims.

[®]Trademark of Dow AgroSciences LLC

(Base label for rigid containers 5 gal or less):

Turf Fertilizer - Contains Team® 1.15%

Herbicide

A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual weed grasses in established turfgrass

and/or

Controls [tough] weeds [such as annual bluegrass] [smooth, hairy crabgrass] [goosegrass] [silver crabgrass] [crowfoot] [barnyardgrass] [watergrass] [green foxtail] [yellow foxtail] [prostrate spurge] [spotted spurge] [creeping spurge] [creeping woodsorrel] [yellow woodsorrel] and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/prevention than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass and other weeds from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types and/or including bentgrass

and/or

For use on cool season and warm season turfgrass

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

and/o

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as]

[crabgrass] [goosegrass] [barnyardgrass]

and/or

University tested [and researched]

and/or

Labeled for use around a [wide] variety of trees and shrubs

and/or

Never see crabgrass again

Active Ingredients:

را) 2 ge 2

trifluralin: α, α, α -trifluoro-2,6-dinitro- <i>N</i> , <i>N</i> -dipropyl- <i>p</i> -toluidine	0.39%
Other Ingredients	
Total	

Contains 0.575 lb active ingredient per 50 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N)	XX.X%
Available Phosphoric Acid (P ₂ O ₅)	XX.X%
Soluble Potash (K ₂ O)	

Keep Out of Reach of Children

WARNING

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Personal Protective Equipment (PPE)

All loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material
- · Goggles or faceshield

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-150

Trademark of Dow AgroSciences LLC.
Produced for
Dow AgroSciences
9330 Zionsville Road
Indianapolis, Indiana 46268

Net	We	ight		
IACL	AAC	IMIIL		

EPA Est. No.

(Base label for nonrigid containers any size):

Turf Fertilizer - Contains Team® 1.15%

Herbicide

A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual weed grasses in established turfgrass

and/or

Controls [tough] weeds [such as annual bluegrass] [smooth, hairy crabgrass] [goosegrass] [silver crabgrass] [crowfoot] [barnyardgrass] [watergrass] [green foxtail] [yellow foxtail] [prostrate spurge] [spotted spurge] [creeping spurge] [creeping woodsorrel] [yellow woodsorrel] and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/prevention than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass and other weeds from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types and/or including bentgrass

and/or

For use on cool season and warm season turfgrass

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]

and/or

University tested [and researched]

and/or

Labeled for use around a [wide] variety of trees and shrubs

and/or

Never see crabgrass again

Active Ingredients:

trifluralin: α, α, α -trifluoro-2,6-dinitro- <i>N</i> , <i>N</i> -dipropyl- <i>p</i> -toluidine	0.39%
Other Ingredients	98.85%
Total	

Contains 0.575 lb active ingredient per 50 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N)	XX.X%
Available Phosphoric Acid (P ₂ O ₅)	
Soluble Potash (K ₂ O)	

Keep Out of Reach of Children

WARNING

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Personal Protective Equipment (PPE)

All loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- · Shoes plus socks
- Chemical-resistant gloves made of any waterproof material
- Goggles or faceshield

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water: Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- · Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Ecracye contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

ge 6

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-150

EPA Est. No.

Trademark of Dow AgroSciences LLC
Produced for
Dow AgroSciences
9330 Zionsville Road
Indianapolis, Indiana 46268

Net Weight	
------------	--

Page 7 /93

(Base label for refillable rigid containers greater than 5 gal):

Turf Fertilizer - Contains Team® 1.15%

Herbicide

A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual weed grasses in established turfgrass

and/or

Controls [tough] weeds [such as annual bluegrass] [smooth, hairy crabgrass] [goosegrass] [silver crabgrass] [crowfoot] [barnyardgrass] [watergrass] [green foxtail] [yellow foxtail] [prostrate spurge] [spotted spurge] [creeping spurge] [creeping woodsorrel] [yellow woodsorrel]

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/prevention than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass and other weeds from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types and/or including bentgrass

and/or

For use on cool season and warm season turfgrass

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

and/o

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]

and/or

University tested [and researched]

and/oi

Labeled for use around a [wide] variety of trees and shrubs

and/or

Never see crabgrass again

Active Ingredients:

trifluralin: α , α , α -trifluoro-2,6-dinitro- N , N -dipropyl- p -toluidine	0.39%
Other Ingredients	
Total	

Contains 0.575 lb active ingredient per 50 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N)	XX.X%
Available Phosphoric Acid (P ₂ O ₅)	XX.X%
Soluble Potash (K ₂ O)	XX.X%

Keep Out of Reach of Children

WARNING

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Personal Protective Equipment (PPE)

All loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material
- · Goggles or faceshield

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Rengve contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.



Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Reuse: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

[®] Trademark o	f Dow	AgroSciences	LLC

EPA Reg. No. 62719-150

Produced for Dow AgroSciences 9330 Zionsville Road Indianapolis, Indiana 46268

Net	Weight	

EPA Est. No. __

(Base label for nonrefillable rigid containers larger than 5 gal):

Turf Fertilizer - Contains Team® 1.15%

Herbicide

A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual weed grasses in established turfgrass

and/or

Controls [tough] weeds [such as annual bluegrass] [smooth, hairy crabgrass] [goosegrass] [silver crabgrass] [crowfoot] [barnyardgrass] [watergrass] [green foxtail] [yellow foxtail] [prostrate spurge] [spotted spurge] [creeping spurge] [creeping woodsorrel] [yellow woodsorrel] and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/o

Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/prevention than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass and other weeds from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types and/or including bentgrass

and/or

For use on cool season and warm season turfgrass

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

and/o

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]

and/or

University tested [and researched]


and/oi

Labeled for use around a [wide] variety of trees and shrubs

and/or

Never see crabgrass again

Active Ingredients:



trifluralin: α,α,α -trifluoro-2,6-dinitro- <i>N</i> , <i>N</i> -dipropyl- <i>p</i> -toluidine	0.39%
Other Ingredients	
Total	

Contains 0.575 lb active ingredient per 50 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N)	XX.X%
Available Phosphoric Acid (P ₂ O ₅)	XX.X%
Soluble Potash (K ₂ O)	XX.X%

Keep Out of Reach of Children

WARNING

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Personal Protective Equipment (PPE)

All loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- · Shoes plus socks
- Chemical-resistant gloves made of any waterproof material
- Goggles or faceshield

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remeve contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Page 12

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Reuse: nrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

v

EPA Reg. No. 62719-150

EPA Est. No.

[®]Trademark of Dow AgroSciences LLC Produced for Dow AgroSciences 9330 Zionsville Road Indianapolis, Indiana 46268

Net Weight _____

(Label booklet cover):

Turf Fertilizer - Contains Team® 1.15%

Herbicide

A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual weed grasses in established turfgrass

and/or

Controls [tough] weeds [such as annual bluegrass] [smooth, hairy crabgrass] [goosegrass] [silver crabgrass] [crowfoot] [barnyardgrass] [watergrass] [green foxtail] [yellow foxtail] [prostrate spurge] [spotted spurge] [creeping spurge] [creeping woodsorrel] [yellow woodsorrel]

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [barnyardgrass] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/prevention than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass and other weeds from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types and/or including bentgrass

and/or

For use on cool season and warm season turfgrass

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]

and/o

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as]

[crabgrass] [goosegrass] [barnyardgrass]

and/or

University tested [and researched]

and/oi

Labeled for use around a [wide] variety of trees and shrubs

and/or

Never see crabgrass again

Active Ingredients:

Page 14

trifluralin: α,α,α -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine	
Other Ingredients	98.85% 100.00%
Contains 0.575 lb active ingredient per 50 lb bag.	
[Editor's note: Supplemental distributor will add fertilizer analysis and defertilizer carrier used.]	erivation statement for
Guaranteed Fertilizer Analysis Total Nitrogen (N) Available Phosphoric Acid (P ₂ O ₅) Soluble Potash (K ₂ O)	XX.X%
Keep Out of Reach of Children WARNING	
Refer to inside of label booklet for additional precautionary information in Use.	ncluding Directions for
Notice: Read the entire label. Use only according to label directions. Before Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies a Use. If terms are unacceptable, return at once unopened.	
In case of emergency endangering health or the environment involving this pro	oduct, call 1-800-992-5994.
Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.	
EPA Reg. No. 62719-150	EPA Est. No.
®Trademark of Dow AgroSciences LLC Produced for Dow AgroSciences 9330 Zionsville Road	
Indianapolis, Indiana 46268	Net Weight

(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals

WARNING

Causes Substantial But Temporary Eye Injury • Harmful If Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Personal Protective Equipment (PPE)

All loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material
- Goggles or faceshield

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Application Restrictions: Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Only protected handlers may be in the area during application.

Entry Restrictions: Do not enter or allow others to enter treated areas until dusts have settled. If soil incorporation is required after application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If incorporation is accomplished by watering-in, except for those people involved in the watering-in, do not enter or allow others to enter the treated area until the watering-in is complete the surface is dry after the watering-in.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Storage and Disposal

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage: Store in original container. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Nonrefillable rigid containers 5 gallons or less:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Nonrefillable nonrigid containers:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Refillable rigid containers larger than 5 gal:

Container Reuse: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable rigid containers larger than 5 gal:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the

rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

General Information

This product is a granular fertilizer containing Team®, a preemergence herbicide for application to established turfgrass for control of annual bluegrass (*Poa annua*), smooth and hairy crabgrass, goosegrass (silver crabgrass or crowfoot), barnyardgrass (watergrass) and green and yellow foxtail. This product also provides partial control of prostrate spurge (*Euphorbia humistrata*), spotted spurge (*Euphorbia maculata*), creeping spurge (*Euphorbia serpens*), creeping woodsorrel (*Oxalis corniculata*) and yellow woodsorrel (*Oxalis stricta*). This product will not harm most nearby ornamental trees and shrubs.

General Use Precautions and Restrictions

- Do not apply this product to golf course putting greens.
- Do not apply to dichondra.
- Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until turf is well established.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.
- Do not make more than 2 applications of this product per year.
- Maximum Single Application Rate: Cool season turf 200 lb of Turf Fertilizer Contains Team 1.15% per acre; warm season turf 260 lb of Turf Fertilizer Contains Team 1.15% per acre.
- Maxmum Yearly Rate: Cool season turf 400 lb of Turf Fertilizer Contains Team 1.15% per acre; warm season turf 520 lb of Turf Fertilizer Contains Team 1.15% per acre.
- Minimum Re-Treatment Interval: Cool season turf 56 to 70 days: warm season turf 70 to 84 days.
- This product must be watered in as soon as possible after application.

Reseeding: Do not reseed treated areas for at least 8 weeks after a single application of 174 lb of product per acre, or 12 to 16 weeks after a single application of 260 lb of product per acre unless injury can be tolerated. For split applications, do not reseed treated areas for 8 to 10 weeks after the last 130 lb of product per acre rate is applied, or 10 to 12 weeks after the last 174 lb of product per acre rate is applied unless injury can be tolerated.

This product is recommended for use on the following established turfgrass species:

Established Cool Season Turfgrass Established Warm Season Turfgrass

bentgrass fairways fescues Kentucky bluegrass perennial ryegrass bahiagrass
Bermudagrass
centipededgrass
St. Augustinegrass
tall fescue
zoysiagrass

Application Directions

Apply the product using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. A



splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

Rates, Frequency and Timing of Application

Broadcast Application Rates

	Cool Season Turfgrass	Warm Season Turfgrass
summer annual grass weeds	130 to 200 lb/acre	260 lb/acre
(spring application)	(3 to 5 lb/1000 sq ft)	(6 lb/1000 sq ft)
winter annual grass weeds	130 to 174 lb/acre	260 lb/acre
(fall application)	(3 to 4 lb/1000 sq ft)	(6 lb/1000 sq ft)

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application. Apply this product 1 to 2 weeks prior to the germination of annual grass weeds since it does not control established weeds. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor.

In areas where germination of the annual weed grass species continues for extended periods of time, make a second application at the recommended rate 8 to 10 weeks after the first application on cool season turfgrass and 10 to 12 weeks after the first application on warm season turfgrass.

This product may also be applied in split applications. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time. For a split application, apply this product at 130 lb plus 130 lb per acre for cool season turfgrass and in the northern half of the warm season turfgrass area, or 174 lb plus 174 lb per acre in the southern half of the warm season turfgrass area. Make the second application 8 to 10 weeks following the 130 lb per acre rate and 10 to 12 weeks following the 174 lb per acre rate.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT PERMITTED BY LAW, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of the Seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

- 1 Refund of purchase price paid by buyer or user for product bought, or
- 2 Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®_{TM}Trademark of Dow AgroSciences LLC EPA accepted 01/15/08