




Turf Fertilizer - Contains Team* 0.92%
EPA Reg. No. 62719-151

**Directions for Use for Established Ornamental
Plants in Landscape Settings**

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for this product.

Directions for Use

Established Ornamental Plants in Landscape Settings

Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

Application

Turf Fertilizer - Contains Team* 0.92% herbicide is recommended for preemergence control of annual grasses and certain broadleaf weeds in a wide variety of established ornamental trees, shrubs, and flowers. It should be applied one to two weeks prior to the anticipated germination of the target annual weeds. Use applicators designed to apply herbicide or insecticide granules. Calibrate the applicator according to the manufacturer's directions prior to use and check frequently during application to be sure the equipment is working properly and distributing the granules uniformly. Do not apply more than the recommended amount. This product will control annual weeds only where it is applied to the soil surface. An even distribution of the granules with no skips or misses is important for satisfactory weed control.

Application timing: Apply one (1) to two (2) weeks prior to the germination of weed seeds or to clean cultivated weed free areas. Do not apply this product to plantings when the foliage is wet.

Soil Preparation: This product can be surface applied around established plants or on ornamental beds. Apply to a smooth soil surface (no large clods) to assure uniform distribution. Clean-cultivate and hoe to remove existing weeds before applying this product.

Activation: Optimum weed control performance will be obtained when followed by overhead irrigation or rainfall, within a few hours after application.

Broadcast Application Rates:

Season of Application	Pounds per Acre	Pounds per 1000 sq ft
Spring Application (Summer annuals)	217 to 326 lb/acre	5 to 7.5 lb/1000 sq ft
Fall Application (Winter annuals)	326 lb/acre	7.5 lb/1000 sq ft

In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 - 12 weeks following the initial application.

BEST COPY AVAILABLE

This product may be used for annual weed control in landscape settings on the following established ornamental plant species:

Ornamental Trees:

<u>Common Name</u>	<u>Scientific Name</u>	<u>Common Name</u>	<u>Scientific Name</u>
almond	<i>Prunus dulcis</i>	maple, sugar	<i>Acer saccharum</i>
apple, crabapple	<i>Malus</i> spp.	oak, pin	<i>Quercus palustris</i>
apricot	<i>Prunus armeniaca</i>	oak, scarlet	<i>Quercus coccinea</i>
ash, white	<i>Fraxinus americana</i>	peach	<i>Prunus persica</i>
baldcypress	<i>Taxodium distichum</i>	pine, Austrian	<i>Pinus nigra</i>
birch, European white	<i>Betula pendula</i>	pine, eastern white	<i>Pinus strobus</i>
blackgum	<i>Nyssa sylvatica</i>	pine, Japanese black	<i>Pinus thunbergiana</i>
cherry	<i>Prunus</i> spp.	pine, loblolly	<i>Pinus taeda</i>
chestnut, Chinese	<i>Castanea mollissima</i>	pine, red	<i>Pinus resinosa</i>
cottonwood	<i>Populus deltoides</i>	pine, scotch	<i>Pinus sylvestris</i>
dogwood, flowering	<i>Cornus florida</i>	planetree, London	<i>Platanus acerifolia</i>
dogwood, kousa	<i>Cornus kousa</i>	plum	<i>Prunus</i> spp.
douglasfir	<i>Pseudotsuga menziesii</i>	redbud, eastern	<i>Cercis canadensis</i>
fir, balsam	<i>Abies balsamea</i>	spruce, Colorado	<i>Picea pungens</i>
hemlock, Canada	<i>Tsuga canadensis</i>	spruce, Norway	<i>Picea abies</i>
honeylocust	<i>Gleditsia triacanthos</i>	spruce, white	<i>Picea glauca</i>
larch, Japanese	<i>Larix kaempferi</i>	sweetgum	<i>Liquidambar styraciflua</i>
locust, black	<i>Robinia pseudoacacia</i>	sycamore	<i>Platanus occidentalis</i>
maple, Norway	<i>Acer platanoides</i>	tuliptree	<i>Liriodendron tulipifera</i>
maple, red	<i>Acer rubrum</i>	walnut, black	<i>Juglans nigra</i>
maple, silver	<i>Acer saccharinum</i>		

Ornamental Shrubs:

<u>Common Name</u>	<u>Scientific Name</u>	<u>Common Name</u>	<u>Scientific Name</u>
andromeda, Japanese	<i>Pieris japonica</i>	forsythia	<i>Forsythia</i> spp.
arbutus, American	<i>Thuja occidentalis</i>	guava, pineapple	<i>Feijoa sellowiana</i>
azalea	<i>Rhododendron</i> spp.	holly	<i>Ilex</i> spp.
barberry, Japanese	<i>Berberis thunbergii</i>	honeysuckle	<i>Lonicera</i> spp.
barberry, mento	<i>Berberis mentorensis</i>	indiahawthorn	<i>Raphiolepis indica</i>
boxwood, common	<i>Buxus sempervirens</i>	juniper	<i>Juniperus</i> spp.
boxwood, littleleaf	<i>Buxus microphylla</i>	laurel, mountain	<i>Kalmia latifolia</i>
camellia, Japanese	<i>Camellia japonica</i>	lilac, common	<i>Syringa vulgaris</i>
camellia, sasanqua	<i>Camellia sasanqua</i>	mockorange	<i>Philadelphus</i> spp.
cherry laurel, American	<i>Prunus caroliniana</i>	pittosporum, Japanese	<i>Pittosporum tobira</i>
cinquefoil	<i>Potentilla</i> spp.	privet	<i>Ligustrum</i> spp.
cleome, Japanese	<i>Cleyera japonica</i>	redcedar, eastern	<i>Juniperus virginiana</i>
cotoneaster, cranberry	<i>Cotoneaster apiculata</i>	rhododendron	<i>Rhododendron</i> spp.
cotoneaster, zabel	<i>Cotoneaster zabelii</i>	spiraea, vanhoutte	<i>Spiraea vanhouttei</i>
deutzia	<i>Deutzia</i> spp.	viburnum	<i>Viburnum</i> spp.
elaeagnus, silverberry	<i>Elaeagnus pungens</i>	weigela	<i>Weigela</i> spp.
euonymus, spreading	<i>Euonymus kiautschovica</i>	willow	<i>Salix</i> spp.
euonymus, winged	<i>Euonymus alatus</i>	yew, anglojap	<i>Taxus media</i>
euonymus, wintercreeper	<i>Euonymus fortunei</i>	yew, Japanese	<i>Taxus cuspidata</i>
firethorn	<i>Pyracantha</i> spp.	yewpine	<i>Podocarpus macrophyllus</i>

Ornamental Groundcover Plantings:

<u>Common Name</u>	<u>Scientific Name</u>	<u>Common Name</u>	<u>Scientific Name</u>
aaronsbeard	<i>Hypericum calycinum</i>	lilyturf, bigblue	<i>Liriope muscari</i>
bellflower, adriatic	<i>Campanula elatines</i>	marigold	<i>Tagetes</i> spp.
bellflower, poscharsky	<i>Campanula poscharskyana</i>	myoporum	<i>Myoporum laetum</i>
ceanothus	<i>Ceanothus</i> spp.	plumbago, dwarf	<i>Ceratostigma umbaginoides</i>
coreopsis	<i>Coreopsis</i> spp.	rockrose	<i>Cistus</i> spp.
cotoneaster	<i>Cotoneaster</i> spp.	rosemary	<i>Rosmarinus officinalis</i>
coyote brush	<i>Baccharis pilularis</i>	rupturewort	<i>Herniaria glabra</i>
crown vetch	<i>Coronilla vana</i>	snow-in-summer	<i>Cerastium tomentosum</i>
daisy trailing African	<i>Osteospermum fruticosum</i>	speedwell	<i>Veronica</i> spp.
fern, asparagus	<i>Asparagus densiflorus</i>	st. johnswort	<i>Hypericum coris</i>
gazania	<i>Gazania</i> spp.	stonecrop (sedum)	<i>Sedum</i> spp.
germander	<i>Teucrium chamaedrys</i>	strawberry, beach	<i>Fragaria chiloensis</i>
ice plant, largeleaf	<i>Carpobrotus edulis</i>	thrift	<i>Ameria maritima</i>
ivy, Algerian	<i>Hedera canariensis</i>	verbena	<i>Verbena</i> spp.
ivy, English	<i>Hedera helix</i>	wirevine, creeping	<i>Muehlenbeckia axillaris</i>
lily-of-the-nile	<i>Agapanthus</i> spp.	yarrow, woolly	<i>Achillea tomentosa</i>

Roses and Other Established Flowers:

African daisy	dianthus	marigold	shasta daisy
aster (perennial)	dusty miller	marigold, cape	snapdragon
balsam	floss flower	morningglory	snow-on-the-mountain
blackeyed susan	forget-me-not	nasturtium	stock
calendula	four o'clock	nicotiana	sunflower
carnation	gaillardia	petunia	sweet alyssum
centaurea, velvet	gladiolus	phlox	sweet pea
chrysanthemum	golden glow	pincushion flower	sweet sultan
coreopsis	impatiens	poppy, California	sweet william
cornflower	ixora	portulaca	vinca
cosmos	lobelia	rose	yarrow
dahlia	lupine	salvia	zinnia

*Trademark of DowElanco

(Label Code) Approved ___/___/___
Initial printing.

Amendments: Adds ornamental uses not within the scope of the WPS (plants not grown for sale or research purposes) to uses for Turf Fertilizer - Contains Team 0.92%.

SEP 21 1994

62719-151

Turf Fertilizer - Contains Team* 0.92%

EPA Reg. No. 62719-151

Directions for Use for Established Ornamental Plants in Landscape Settings**ATTENTION**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for this product.

Directions for Use**Established Ornamental Plants in Landscape Settings**

Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

Application

Turf Fertilizer - Contains Team* 0.92% herbicide is recommended for preemergence control of annual grasses and certain broadleaf weeds in a wide variety of established ornamental trees, shrubs, and flowers. It should be applied one to two weeks prior to the anticipated germination of the target annual weeds. Use applicators designed to apply herbicide or insecticide granules. Calibrate the applicator according to the manufacturer's directions prior to use and check frequently during application to be sure the equipment is working properly and distributing the granules uniformly. Do not apply more than the recommended amount. This product will control annual weeds only where it is applied to the soil surface. An even distribution of the granules with no skips or misses is important for satisfactory weed control.

Application timing: Apply one (1) to two (2) weeks prior to the germination of weed seeds or to clean cultivated weed free areas. Do not apply this product to plantings when the foliage is wet.

Soil Preparation: This product can be surface applied around established plants or on ornamental beds. Apply to a smooth soil surface (no large clods) to assure uniform distribution. Clean-cultivate and hoe to remove existing weeds before applying this product.

Activation: Optimum weed control performance will be obtained when followed by overhead irrigation or rainfall, within a few hours after application.

Broadcast Application Rates:

Season of Application	Pounds per Acre	Pounds per 1000 sq ft
Spring Application (Summer annuals)	217 to 326 lb/acre	5 to 7.5 lb/1000 sq ft
Fall Application (Winter annuals)	326 lb/acre	7.5 lb/1000 sq ft

In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 - 12 weeks following the initial application.

BEST AVAILABLE COPY

This product may be used for annual weed control in landscape settings on the following established ornamental plant species:

Ornamental Trees:

<u>Common Name</u>	<u>Scientific Name</u>	<u>Common Name</u>	<u>Scientific Name</u>
almond	<i>Prunus dulcis</i>	maple, sugar	<i>Acer saccharum</i>
apple, crabapple	<i>Malus</i> spp.	oak, pin	<i>Quercus palustris</i>
apricot	<i>Prunus armeniaca</i>	oak, scarlet	<i>Quercus coccinea</i>
ash, white	<i>Fraxinus americana</i>	peach	<i>Prunus persica</i>
baldcypress	<i>Taxodium distichum</i>	pine, Austrian	<i>Pinus nigra</i>
birch, European white	<i>Betula pendula</i>	pine, eastern white	<i>Pinus strobus</i>
blackgum	<i>Nyssa sylvatica</i>	pine, Japanese black	<i>Pinus thunbergiana</i>
cherry	<i>Prunus</i> spp.	pine, loblolly	<i>Pinus taeda</i>
chestnut, Chinese	<i>Castanea mollissima</i>	pine, red	<i>Pinus resinosa</i>
cottonwood	<i>Populus deltoides</i>	pine, scotch	<i>Pinus sylvestris</i>
dogwood, flowering	<i>Cornus florida</i>	planetree, London	<i>Platanus acerifolia</i>
dogwood, kousa	<i>Cornus kousa</i>	plum	<i>Prunus</i> spp.
douglasfir	<i>Pseudotsuga menziesii</i>	redbud, eastern	<i>Cercis canadensis</i>
fir, balsam	<i>Abies balsamea</i>	spruce, Colorado	<i>Picea pungens</i>
hemlock, Canada	<i>Tsuga canadensis</i>	spruce, Norway	<i>Picea abies</i>
honeylocust	<i>Gleditsia triacanthos</i>	spruce, white	<i>Picea glauca</i>
larch, Japanese	<i>Larix kaempferi</i>	sweetgum	<i>Liquidambar styraciflua</i>
locust, black	<i>Robinia pseudoacacia</i>	sycamore	<i>Platanus occidentalis</i>
maple, Norway	<i>Acer platanoides</i>	tuliptree	<i>Liriodendron tulipifera</i>
maple, red	<i>Acer rubrum</i>	walnut, black	<i>Juglans nigra</i>
maple, silver	<i>Acer saccharinum</i>		

Ornamental Shrubs:

<u>Common Name</u>	<u>Scientific Name</u>	<u>Common Name</u>	<u>Scientific Name</u>
andromeda, Japanese	<i>Pieris japonica</i>	forsythia	<i>Forsythia</i> spp.
arborvitae, American	<i>Thuja occidentalis</i>	guava, pineapple	<i>Feijoa sellowiana</i>
azalea	<i>Rhododendron</i> spp.	holly	<i>Ilex</i> spp.
barberry, Japanese	<i>Berberis thunbergii</i>	honeysuckle	<i>Lonicera</i> spp.
barberry, mento	<i>Berberis mentorensis</i>	indiahawthorn	<i>Raphiolepis indica</i>
boxwood, common	<i>Buxus sempervirens</i>	juniper	<i>Juniperus</i> spp.
boxwood, littleleaf	<i>Buxus microphylla</i>	laurel, mountain	<i>Kalmia latifolia</i>
camellia, Japanese	<i>Camellia japonica</i>	lilac, common	<i>Syringa vulgaris</i>
camellia, sasanqua	<i>Camellia sasanqua</i>	mockorange	<i>Philadelphus</i> spp.
cherrylaurel, American	<i>Prunus caroliniana</i>	pittosporum, Japanese	<i>Pittosporum tobira</i>
cinquefoil	<i>Potentilla</i> spp.	privet	<i>Ligustrum</i> spp.
cleysa, Japanese	<i>Cleyera japonica</i>	redcedar, eastern	<i>Juniperus virginiana</i>
cotoneaster, cranberry	<i>Cotoneaster apiculata</i>	rhododendron	<i>Rhododendron</i> spp.
cotoneaster, zabel	<i>Cotoneaster zabelii</i>	spiraea, vanhoutte	<i>Spiraea vanhouttei</i>
deutzia	<i>Deutzia</i> spp.	viburnum	<i>Viburnum</i> spp.
elaeagnus, silverberry	<i>Elaeagnus pungens</i>	weigela	<i>Weigela</i> spp.
euonymus, spreading	<i>Euonymus kiautschovica</i>	willow	<i>Salix</i> spp.
euonymus, winged	<i>Euonymus alatus</i>	yew, anglojap	<i>Taxus media</i>
euonymus, wintercreeper	<i>Euonymus fortunei</i>	yew, Japanese	<i>Taxus cuspidata</i>
firethorn	<i>Pyracantha</i> spp.	yewpine	<i>Podocarpus macrophyllus</i>

Ornamental Groundcover Plantings:

Common Name

aaronsbeard
bellflower, adriatic
bellflower, poscharsky
ceanothus
coreopsis
cotoneaster
coyote brush
crown vetch
daisy trailing African
fern, asparagus
gazania
germander
ice plant, largeleaf
ivy, Algerian
ivy, English
lily-of-the-nile

Scientific Name

Hypericum calycinum
Campanula elatines
Campanula poscharskyana
Ceanothus spp.
Coreopsis spp.
Cotoneaster spp.
Baccharis pilularis
Coronilla vana
Osteospermum fruticosum
Asparagus densiflorus
Gazania spp.
Teucrium chamaedrys
Carpobrotus edulis
Hedera canariensis
Hedera helix
Agapanthus spp.

Common Name

lilyturf, bigblue
marigold
myoporum
plumbago, dwarf
rockrose
rosemary
rupturewort
snow-in-summer
speedwell
st. johnswort
stonecrop (sedum)
strawberry, beach
thrift
verbena
wirevine, creeping
yarrow, woolly

Scientific Name

Liriope muscari
Tagetes spp.
Myoporum laetum
Ceratostigma umbaginoides
Cistus spp.
Rosmarinus officinalis
Hemaria glabra
Cerastium tomentosum
Veronica spp.
Hypericum coris
Sedum spp.
Fragaria chiloensis
Armeria maritima
Verbena spp.
Muehlenbeckia axillaris
Achillea tomentosa

Roses and Other Established Flowers:

African daisy
aster (perennial)
balsam
blackeyed susan
calendula
carnation
centaurea, velvet
chrysanthemum
coreopsis
cornflower
cosmos
dahlia

dianthus
dusty miller
floss flower
forget-me-not
four o'clock
gaillardia
gladiolus
golden glow
impatiens
ixora
lobelia
lupine

marigold
marigold, cape
morningglory
nasturtium
nicotiana
petunia
phlox
pincushion flower
poppy, California
portulaca
rose
salvia

shasta daisy
snapdragon
snow-on-the-mountain
stock
sunflower
sweet alyssum
sweet pea
sweet william
vinca
yarrow
zinnia

BEST AVAILABLE COPY

*Trademark of DowElanco

(Label Code) Approved ___/___/___
Initial printing.

Amendments: Adds ornamental uses not within the scope of the WPS (plants not grown for sale or research purposes) to uses for Turf Fertilizer - Contains Team 0.92%