

11103

10719 152

10/9

DowElanco

**TURF FERTILIZER**  
contains  
**TEAM™ 1.25%**

A granular fertilizer containing TEAM™ for the preemergence control of crabgrass, foxtail, goosegrass, poa annua and other annual weed grasses in established turfgrasses.

Active Ingredients:

N-butyl-N-ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2, 6-dinitro-p-toluidine ..... 0.82%  
trifluralin:  $\alpha,\alpha,\alpha$ -trifluoro-2, 6-dinitro-N, N-dipropyl-p-toluidine ..... 0.43%

Inert Ingredients ..... 98.75%

Contains 0.625 pounds active ingredient per 50 pound lb. bag.

TEAM™ (benefin plus trifluralin, DowElanco)

**GUARANTEED ANALYSIS**

Total Nitrogen (N) ..... 24.00%

Available Phosphoric Acid ( $P_2O_5$ ) ..... 6.00%

Soluble Potash ( $K_2O$ ) ..... 12.00%

**PRECAUTIONARY STATEMENTS**

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

**WARNING**

Causes eye irritation. Do not get in eyes. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust and contact with skin or clothing. The granular formulation may cause skin sensitization reactions in certain individuals. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**STATEMENTS OF PRACTICAL TREATMENT**

If in eyes: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. See an ophthalmologist or physician immediately.

If swallowed: Call physician or Poison Control Center. Drink promptly a large quantity of milk, egg white, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol. Do not induce vomiting.

If on the skin: Wash with plenty of soap and water. Get medical attention if irritation develops.

**BEST AVAILABLE COPY**

2 of 4

If inhaled: If breathing difficulty occurs, move individual to fresh air and get medical attention. If not breathing, provide cardiopulmonary resuscitation assistance (mouth-to-mouth) and call a physician.

### **ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, or marshes). Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sites. Do not contaminate water when disposing of equipment washwaters.

### **STORAGE AND DISPOSAL**

**Storage:** Store in original container only. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### **DIRECTIONS FOR USE - Read All Directions Carefully Before Applying**

DowElanco's Turf Fertilizer containing Team 1.25% can be used on the following established cool and warm season turfgrasses:

#### **Established Cool Season Turfgrasses**

- Bentgrass Fairways
- Kentucky Bluegrass
- Perennial Ryegrass
- Fescues

#### **Established Warm Season Turfgrasses**

- Bahiagrass
- Bermudagrass
- Centipedegrass
- St. Augustinegrass
- Tall Fescue
- Zoysiagrass

DowElanco fertilizer containing Team 1.25% is a preemergence herbicide which is applied to established turfgrasses for the control of the winter annual grass: annual bluegrass (Poa annua); and summer annual grasses: smooth and hairy crabgrass, goosegrass (silver crabgrass or crow-footgrass), barnyardgrass (watergrass) and green and yellow foxtail. Turf fertilizer containing Team 1.25% also provides partial control of prostrate spurge (Euphorbia humistrata), spotted spurge (Euphorbia maculata), creeping spurge (Euphorbia maculata), creeping wood sorrel (Oxalis corniculata) and yellow wood sorrel (Oxalis stricta).

This product should be applied one to two weeks prior to the anticipated germination of the target annual weeds. Apply using a properly calibrated granular fertilizer spreader which will uniformly distribute the product over the target area. Avoid skips and excessive overlaps as poor control or turfgrass injury may occur. This product will not harm most nearby ornamentals.

**DO NOT MIX WITH OTHER PRODUCTS**

Rates and Frequency of Application:

	<u>Cool Season Turfgrasses</u>	<u>Warm Season Turfgrasses</u>
Summer Annual	160 - 240 lbs./acre	240 lbs./acre
Grass Weeds	<del>(3.7 - 5.5 lbs./1000 sq. ft.)</del> —(3.6 - 5.5 lbs./1000 sq. ft.)	(5.5 lbs./1000 sq. ft.)
Winter Annual	160 lbs./acre	240 lbs./acre
Grass Weeds		

In areas with extended growing seasons or where desired, a second application at the recommended rate can be made 8 to 10 weeks following the initial application on cool season turfgrasses and 10 to 12 weeks following the initial application on warm season turfgrasses.

**Split Applications for Extended Control and Lawn Care Operator Programs**

As an alternative to single application programs DowElanco Turf Fertilizer containing Team 1.25% may also be applied in split applications. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time.

Apply DowElanco Fertilizer containing Team 1.25% at 120 pounds plus 120 pounds per acre for Cool Season Turfgrass and in the northern half of the Warm Season Turfgrass area of the U. S. or 160 pounds plus 160 pounds per acre in the southern half of the Warm Season Turfgrasses area of the U.S. The second application should be made 8 to 10 weeks following the 120 pounds per acre rate and 10 to 12 weeks following the 160 pounds per acre rate.

Optimum weed control performance will be obtained if treated areas are irrigated soon after application.

**RESEEDING**

Do not reseed treated areas for a minimum of 8 weeks after a single application of 160 pounds per acre or 12 to 16 weeks after a single application of 240 pounds per acre. For split applications, do not reseed treated areas for 8 to 10 weeks after the last 120 pounds per acre rate or 10 to 12 weeks after the last 160 pounds per acre rate.

**GENERAL USE PRECAUTION**

Do not apply this product to golf course putting greens, dichondra, or newly sprigged areas of bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass. Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application.

**DISCLAIMER OF WARRANTIES**

DowElanco makes no warranties concerning this product or its use, which extend beyond the description on the label. All other warranties, expressed, or implied, including but not limited to, any implied warranties of merchantability and fitness for a particular purpose, are disclaimed. All statements concerning this product apply only when used as directed. This disclaimer of warranties does not apply where prohibited by operation of law.

**BEST AVAILABLE COPY**

4511

**LIMITATION OF DAMAGES**

The liability of DowElanco whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the return of the amount of the purchase price of this product and under no circumstances shall DowElanco be liable for special, indirect or consequential damages. This limitation of damages does not apply where prohibited by operation of law.

TEAM™ (benefin plus trifluralin, DowElanco)

EPA Est. No. XXXXX-XX-XX

EPA Reg. No. 62719-152

DowElanco

Indianapolis, IN 46268 U.S.A.

Team Turf Fertilizer/1.25

07/08/88

**Amendment:** Lower rate in rate range adjusted from 3.7 to 3.6 lb/1,000 sq ft.