

Reg # 62719-152

PM-25

194

(Product Label)

Logo - DowElanco

Turf Fertilizer - Contains Team* 1.25%

A Granular Fertilizer Containing Team for the Preemergence Control of Crabgrass, Foxtail, Goosegrass, *Poa Annua* and Other Annual Weed Grasses in Established Turfgrasses.

Active Ingredients:

| | |
|---|---------|
| <i>N</i> -butyl- <i>N</i> -ethyl- α,α,α -trifluoro-2,6-dinitro- <i>p</i> -toluidine | 0.82% |
| trifluralin (α,α,α -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine) | 0.43% |
| Inert Ingredients | 98.75% |
| Total..... | 100.00% |

Guaranteed Fertilizer Analysis†

| | |
|---|-------|
| Total Nitrogen (N)..... | 24.0% |
| Available Phosphoric Acid (P ₂ O ₅)..... | 6.0% |
| Soluble Potash (K ₂ O)..... | 12.0% |

[Editors note: See subregistrant label for specific fertilizer analysis of the fertilizer blend used.]
Contains 0.625 pounds active ingredient per 50 pound bag.

Keep Out of Reach of Children
WARNING

See back or container/side panel for additional precautionary statements.

Notice: Read the entire label. Use only according to label directions. Before buying this product, read Limited Warranty and Disclaimer Section elsewhere on this label.

In case of emergency endangering health or the environment involving this product, call collect 517-636-4400

Do Not Ship or Store with Food, Feeds, Drugs, or Clothing.

EPA Reg. No. 62719-152
EPA Est. No. _____

*Trademark of DowElanco

DowElanco • Indianapolis, Indiana 46268, U.S.A.

Herbicide
Net Weight - 50 Pounds

ACCEPTED
 Nov. 1, 1993
 NOV - 1 1993
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act
 as amended, for the pesticide
 registered under
 EPA Reg. No. 62719-152

294

Precautionary Statements

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

WARNING

Causes Eye Irritation • Do Not Get In Eyes • Harmful If Swallowed, Inhaled Or Absorbed Through The Skin

Avoid breathing dust and contact with skin or clothing. The granular formulation may cause skin sensitization reactions in certain individuals. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Statements of Practical Treatment

If in eyes: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. See an ophthalmologist or physician immediately.

If swallowed: Call physician or Poison Control Center. Drink promptly a large quantity of milk, egg white, gelatin solution, or if these are not available, large quantities of water. Avoid alcohol. Do not induce vomiting.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation develops.

If inhaled: If breathing difficulty occurs, move individual to fresh air and get medical attention. If not breathing, provide cardiopulmonary resuscitation assistance (mouth-to-mouth) and call a physician.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sties. Do not contaminate water when disposing of equipment washwaters..

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

General Information

This turf fertilizer product contains Team, a preemergence herbicide which is applied to established turfgrasses for the control of the winter annual grass: annual bluegrass (*Poa Annua*); and summer annual grasses: smooth and hairy crabgrass, goosegrass (silver crabgrass or crowfootgrass), barnyardgrass (watergrass) and green and yellow foxtail. This product also provides partial control of prostrate spurge (*Euphorbia humistrata*), spotted spurge (*Euphorbia maculata*), creeping spurge (*Euphorbia maculata*), creeping woodsorrel (*Oxalis corniculata*) and yellow woodsorrel (*Oxalis stricta*).

Important Use Precautions

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Reseeding: Do not reseed treated areas for minimum of 8 weeks after a single application of 160 pounds per acre or 12 to 16 weeks after a single application of 240 pounds per acre. For split applications, do not reseed treated areas for 8 to 10 weeks after the last 120 pounds per acre rate or 10 to 12 weeks after the last 160 pounds per acre rate.

Do not apply this product to golf course putting greens, dichondra, or newly sprigged areas of bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass. Any cultural practices that disturb the soil, such as aeration or verticutting, should be done prior to application.

This Product is Recommended for Use on the Following Established Warm and Cool Season Perennial Turfgrass Species:

Established Cool Season Turfgrasses

- Bentgrass fairways
- Kentucky bluegrass
- Perennial ryegrass
- Fescues

Established Warm Season Turfgrasses

- Bahiagrass
- Bermudagrass
- Centipedegrass
- Fescue, tall
- St. Augustinegrass
- Zoysiagrass

Application

This product should be applied one to two weeks prior to the anticipated germination of the target annual weeds. Apply using a properly calibrated granular fertilizer spreader which will uniformly distribute the product over the target area. Avoid skips and excessive overlaps as poor weed control or turfgrass injury may occur. Optimum weed control performance will be obtained if treated areas are irrigated soon after application. This product will not harm most nearby ornamentals.

In areas with extended growing seasons or where desired, a second application at the recommended rate can be made 8 to 10 weeks following the initial application on cool season turfgrasses and 10 to 12 weeks following the initial application on warm season turfgrasses.

Broadcast Application Rates:

| | Cool Season Turfgrasses | Warm Season Turfgrasses |
|---|---|----------------------------------|
| Summer Annual Grass Weeds (Spring Application) | 160 - 240 lb/acre 3.6 to 5.5 lb/1000 sq ft | 240 lb/acre 5.5 lb/1000 sq ft |
| Winter Annual Grass Weeds (Fall application) | 160 lb/acre | 240 lb/acre |

Split Applications for Extended Control and Lawn Care Operator Programs:

As an alternative to single application programs, DowElanco Turf Fertilizer containing Team may also be applied in split applications. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time.

Apply DowElanco Fertilizer containing Team 1.25% at 120 pounds plus 120 pounds per acre for Cool Season Turfgrasses and in the northern half of the Warm Season Turfgrass area of the U.S. or 160 pounds plus 160 pounds per acre in the southern half of the Warm Season Turfgrasses area of the U.S. The second application should be made 8 to 10 weeks following the 120 pounds per acre rate and 10 to 12 weeks following the 160 pounds per acre rate.

Spreader Settings

(Editors note: Subregistrant label may include a table of recommended spreader equipment and spreader settings for the fertilizer blend used.)

Warranty Disclaimer/Inherent Risks of Use/Limitation of Remedies

(Editors note: Subregistrant label may include appropriate product warranty and disclaimer statements.)

Copyright © 1993 DowElanco