


67719-152

01/11/2008

1/12



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
 Registration Division (7505P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:	Date of Issuance:
62719-152	1-11-08

Term of Issuance:

Name of Pesticide Product:
 Turf Fertilizer Contains Team
 1/25%

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):
 Dow AgroSciences LLC
 9330 Zionsville Rd
 Indianapolis, IN 46268

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA section 4(g)(2)(C) provided you agree in writing to:

Commercial Label

1. Change the Signal Word to "Warning".
2. To the First Aid section add "If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice." below the "If in Eyes" section.
3. Change the Hazards to Humans and Domestic Animals section to "Causes substantial but temporary eye injury. Harmful if absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals."
4. Change the PPE to "All loaders, applicators, and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks, chemical-resistant gloves (such as or made of any waterproof material), and goggles, face shield or safety glasses."
5. Change the entry restrictions currently on the label to "Do not enter or allow other people (or pets) to enter the treated area until dusts have settled. If watering-in is required after the application- except for those people involved in the watering-in do not enter or allow other

[Handwritten signature] 1-11-08


people (or pets) to enter the treated areas until the watering-in is complete and the surface is dry.”

- 6. The watering in text on the must read “This product must be watered in as soon as possible after application. Watering-in must be performed by the commercial applicator or the commercial applicator must provide the following instructions to the resident or owner in writing: This product must be watered in as soon as possible. Do not enter or allow others (including children or pets) to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.”
- 7. Remove “recommended” from the label.
- 8. To the Warranty section add “to the extent consistent with applicable law” in front of “seller makes no”, “seller shall not”, and “in no case”. Change “to the extent permitted by law” to “to the extent consistent with applicable law”.

Residential Label

- 9. Change the Signal Word to “Warning”.
- 10. To the First Aid section add “If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.” below the “If in Eyes” section.
- 11. Change the Hazards to Humans and Domestic Animals section to “Causes substantial but temporary eye injury. Harmful if absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.”
- 12. Remove “recommended” from the label.
- 13. To the Warranty section add “to the extent consistent with applicable law” in front of “seller makes no”, “seller shall not”, and “in no case”. Change “to the extent permitted by law” to “to the extent consistent with applicable law”.

Signature of Approving Official:



Erik Kraft, Acting Product Manager (23)
Herbicide Branch, Registration Division (7505P)

Date:

1-11-08

EPA Form 8570-6

You will submit one copy of your final printed labeling before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). If you have any questions please call Erik Kraft at 703-308-9358.

3/12

P4A /Turf Fert Contains Team 1.25% / MSTR Amend / 10-10-06

ACCEPTED
with COMMENTS
in EPA Letter Dated

1-11-08

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

[Sub Label A: Professional Uses]

(Label)

Turf Fertilizer - Contains Team™ 1.25%

62719-152

Lite

Herbicide

A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual weed grasses in established turfgrass

Active Ingredients:

benefin: <i>N</i> -butyl- <i>N</i> -ethyl- α,α,α -trifluoro-2,6-dinitro- <i>p</i> -toluidine	0.82%
trifluralin: α,α,α -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine	0.43%
Other Ingredients	98.75%
Total	100.00%

Contains 0.625 lb active ingredient per 50 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N)	XX.X%
Available Phosphoric Acid (P ₂ O ₅)	XX.X%
Soluble Potash (K ₂ O)	XX.X%

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Moderate Eye Irritation • Harmful If Swallowed

Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Personal Protective Equipment (PPE)

All loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

4/12

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Refer to back panel for Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-152

EPA Est. No. _____

™Trademark of Dow AgroSciences LLC

Dow AgroSciences LLC • Indianapolis, Indiana 46268 U.S.A.

Net Weight _____

5/12

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Application Restrictions: Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Entry Restrictions: Do not enter or allow others to enter treated areas until dusts have settled. If soil incorporation is required after application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

General Information

This product is a granular fertilizer containing Team™, a preemergence herbicide for application to established turfgrass for control of annual bluegrass (*Poa annua*), smooth and hairy crabgrass, goosegrass (silver crabgrass or crowfoot), barnyardgrass (watergrass) and green and yellow foxtail. This product also provides partial control of prostrate spurge (*Euphorbia humistrata*), spotted spurge (*Euphorbia maculata*), creeping spurge (*Euphorbia serpens*), creeping woodsorrel (*Oxalis corniculata*) and yellow woodsorrel (*Oxalis stricta*). This product will not harm most nearby ornamental trees and shrubs.

General Use Precautions and Restrictions

- Do not apply this product to golf course putting greens.
- Do not apply to dichondra.
- Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until turf is well established.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.
- Do not make more than 2 applications of this product per year.
- **Maximum Single Application Rate:** Cool season turf – 190 lb of Turf Fertilizer – Contains Team 1.25% per acre; warm season turf – 240 lb of Turf Fertilizer – Contains Team 1.25% per acre.
- **Maximum Yearly Rate:** Cool season turf – 380 lb of Turf Fertilizer – Contains Team 1.25% per acre; warm season turf – 480 lb of Turf Fertilizer – Contains Team 1.25% per acre.
- **Minimum Re-Treatment Interval:** Cool season turf – 56 to 70 days; warm season turf – 70 to 84 days.
- This product must be watered in as soon as possible after application.

Reseeding: Do not reseed treated areas for at least 8 weeks after a single application of 160 lb of product per acre, or 12 to 16 weeks after a single application of 240 lb of product per acre unless injury can be tolerated. For split applications, do not reseed treated areas for 8 to 10 weeks after the last 120 lb

6/12

of product per acre rate is applied, or 10 to 12 weeks after the last 160 lb of product per acre rate is applied unless injury can be tolerated.

This product is recommended for use on the following established turfgrass species:

Established Cool Season Turfgrass	Established Warm Season Turfgrass
bentgrass fairways	bahiagrass
fescues	Bermudagrass
Kentucky bluegrass	centipedegrass
perennial ryegrass	St. Augustinegrass
	tall fescue
	zoysiagrass

Application Directions

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

Rates, Frequency and Timing of Application

Broadcast Application Rates

	Cool Season Turfgrass	Warm Season Turfgrass
summer annual grass weeds (spring application)	120 to 190 lb/acre (2.75 to 4 lb/1000 sq ft)	240 lb/acre (5.5 lb/1000 sq ft)
winter annual grass weeds (fall application)	120 to 160 lb/acre (2.75 to 3.6 lb/1000 sq ft)	240 lb/acre (5.5 lb/1000 sq ft)

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application. Apply this product 1 to 2 weeks prior to the germination of annual grass weeds. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor.

In areas where germination of the annual weed grass species continues for extended periods of time, make a second application at the recommended rate 8 to 10 weeks after the first application on cool season turfgrass and 10 to 12 weeks after the first application on warm season turfgrass.

This product may also be applied in split applications. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time. For a split application, apply this product at 120 lb plus 120 lb per acre for cool season turfgrass and in the northern half of the warm season turfgrass area, or 160 lb plus 160 lb per acre in the southern half of the warm season turfgrass area. Make the second application 8 to 10 weeks following the 120 lb per acre rate and 10 to 12 weeks following the 160 lb per acre rate.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

7/12

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of the Seller. All such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

- 1 Refund of purchase price paid by buyer or user for product bought, or
- 2 Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

™Trademark of Dow AgroSciences LLC
EPA accepted __/__/__

8/12

[Sub Label B: Residential Uses]

(Label)

Super Team™ 1.25%

Herbicide

A fertilizer containing Team, a preemergence herbicide for control of crabgrass, foxtail, goosegrass, *Poa annua* and other annual weed grasses in established home lawns

[Editor's note: One or more of the following statements may be included on the label.]
and/or

Prevents crabgrass & other unsightly annual weeds
and/or

Encourages quick spring greenup
and/or

Greens your lawn with extended feeding nitrogen
and/or

Covers up to 15,000 sq ft (100' X 150')
and/or

Covers up to 5,000 sq ft (100' X 50')
and/or

Prevents crabgrass before it starts
and/or

Made in the USA

Active Ingredients:

benefin: N-butyl-N-ethyl- α,α,α -trifluoro-2,6-dinitro-p-toluidine	0.82%
trifluralin: α,α,α -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine	0.43%
Other Ingredients	98.75%
Total	100.00%

Contains 0.625 lb active ingredient per 50 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

Guaranteed Fertilizer Analysis	
Total Nitrogen (N)	XX.X%
Available Phosphoric Acid (P ₂ O ₅)	XX.X%
Soluble Potash (K ₂ O)	XX.X%

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Moderate Eye Irritation • Harmful If Swallowed

9/12

Avoid contact with eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater or rinsate.

Refer to back panel for Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-152

EPA Est. No. _____

™Trademark of Dow AgroSciences LLC

Dow AgroSciences LLC • Indianapolis, Indiana 46268 USA

Net Weight _____

10/12

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Application Restrictions: Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Entry Restrictions: Do not allow people or pets to enter the treated areas until dusts have settled. If watering-in is required after the application, do not enter or allow others to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: If empty: Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your solid waste agency or 1-800-992-5994 for disposal instructions.

Never place unused product down any indoor or outdoor drain.

How This Product Works

This product is a granular fertilizer containing Team™, a preemergence herbicide for application to established home lawns and other turfgrass areas for control of annual bluegrass (*Poa annua*), smooth and hairy crabgrass, goosegrass (silver crabgrass or crowfoot), barnyardgrass (watergrass) and green and yellow foxtail. This product also provides partial control of prostrate spurge, spotted spurge, creeping spurge, creeping woodsorrel and yellow woodsorrel. This product will not harm most nearby ornamental trees and shrubs.

General Use Precautions and Restrictions

- Do not apply this product to dichondra.
- Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until turf is well established.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.
- **Reseeding:** Do not reseed treated areas for at least 8 weeks after a single application of 3.6 lb of product per 1000 sq ft unless injury can be tolerated, or 12 to 16 weeks after a single application of 5.5 lb of product per 1000 sq ft unless injury can be tolerated.
- Do not make more than 2 applications of this product per year.
- **Maximum Single Application Rate:** 5.5 lb of Super Team 1.25% per 1000 sq ft.
- **Maximum Yearly Rate:** 11 lb of Super Team 1.25% per 1000 sq ft.
- **Minimum Re-Treatment Interval:** Cool season turf – 56 to 70 days; warm season turf – 70 to 84 days.

This product is recommended for use on the following established turfgrass species:

Established Cool Season Turfgrass

bentgrass fairways
fescues
Kentucky bluegrass
perennial ryegrass

Established Warm Season Turfgrass

bahiagrass
Bermudagrass
centipedegrass
St. Augustinegrass
tall fescue

11/12

zoysiagrass

How to Apply

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution. Optimum weed control performance will be obtained when an application is followed by sprinkler irrigation or rainfall within a few hours.

When to Apply

Apply this product 1 to 2 weeks prior to germination of target weeds since it does not control established weeds.

How Much to Apply

	Cool Season Turfgrass	Warm Season Turfgrass
summer annual grass weeds (spring application)	2.75 to 5.5 lb/1000 sq ft	5.5 lb/1000 sq ft
winter annual grass weeds (fall application)	2.75 to 3.6 lb/1000 sq ft	5.5 lb/1000 sq ft

Use the following rate table to determine the total amount of product to apply based upon total size of the area to be treated.

Size of Area to be Treated (sq ft)	Total Amount of Product Required/1000 sq ft		
	2.75 lb Rate	3.6 lb Rate	5.5 lb Rate
1000	2.75	3.6 lb	5.5 lb
2000	5.5	7.2 lb	11 lb
3000	8.3	10.8 lb	16.5 lb
4000	11	14.4 lb	22 lb
5000	13.8	18 lb	27.5 lb

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to

12/12

label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of the Seller. All such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

™Trademark of Dow AgroSciences LLC
EPA accepted ___/___/___