


Specimen Label



Pathfinder*

Ready-to-Use Herbicide

Forests,

Rangeland and Permanent Pastures,

For the Control of Woody Plants on Rights-of-Way, Industrial Sites, Non-Crop Areas, ▲
Non-Irrigation Ditch Banks, and Wildlife Openings, Including Grazed Areas on These Sites

Active Ingredient(s):

Triclopyr (3,5,6-trichloro-2-pyridinyloxyacetic acid), Butoxyethyl Ester 16.7%

Inert Ingredient(s): 83.3%

Acid Equivalent: Triclopyr - 12.01% - 1 lb/gal

Contains petroleum distillates

E.P.A. Registration No. 62719-176

E.P.A. Est. 464-MI-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUCION:

PRECAUCION AL USUARIO:

Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

**HARMFUL IF SWALLOWED,
INHALED OR ABSORBED
THROUGH SKIN • CAUSES EYE
IRRITATION**

Avoid Contact With Eyes, Skin, Or
Clothing • Avoid Breathing Mists or
Vapors • Avoid Contamination Of Food
• Wash Thoroughly After Handling
• Remove And Wash Contaminated
Clothing Before Reuse

STATEMENTS OF PRACTICAL TREATMENT

In case of skin contact, flush skin with plenty of water.
Get medical attention if irritation persists. If swallowed,
do not induce vomiting. Call a physician. If in eyes,
flush with plenty of water. Call a physician if irritation
persists.

Physical or Chemical Hazards

**COMBUSTIBLE • Do Not Use or Store
Near Heat or Open Flame. • Do Not Cut or
Weld Container.**

Environmental Hazards

This pesticide is toxic to fish. Keep out of lakes, ponds or
streams. Do not contaminate water by cleaning of
equipment or disposal of wastes. Do not contaminate
water when disposing of equipment washwaters.

NOTICE

Read and understand the entire label before using. Use
only according to label directions.

Before buying or using this product, read "WARRANTY
LIMITATIONS AND DISCLAIMER" elsewhere on this label.
If terms are not acceptable, return unopened package at
once to seller for full refund of purchase price paid.
Otherwise, use by the buyer or any other user constitutes
acceptance of the terms under WARRANTY LIMITATIONS
AND DISCLAIMER.

IN CASE OF AN EMERGENCY

endangering life or property involving this
product, call collect 517-636-4400

AGRICULTURAL CHEMICAL

**Do Not Ship or Store with Food, Feeds,
Drugs, or Clothing**

See Back Panel for Important Use Precautions.

*
Pathfinder

in forests, rangeland and permanent pastures.

GENERAL INFORMATION

PATHFINDER Herbicide is a ready-to-use product which is recommended for the control of unwanted woody plants through the use of basal bark application techniques and on non-crop areas including industrial manufacturing and storage sites, rights-of-way such as electrical power lines, communication lines, pipelines, road sides and railroads, fence rows, non-irrigation ditch banks and around farm buildings. Use on these sites may include application to grazed areas as well as establishment and maintenance of wildlife openings.

Salt Cedar (1)

Among the woody plant species controlled are:

Alder, Speckled	Alnus	Hercules Club	Persimmon,
Australian Pine	Alder, Red	Hickory, Mockernut	Common
Basswood	Ash, Green	Hickory, Pignut	Pine, Jack
	Ash, White	Honeylocust	Pine, Loblolly
	Aspen(?)	Hornbeam	Pine,
	Beech,	(Blue Beech)	Ponderosa
	American	Locust, Black(?)	Pine, Red
	Birch, Black	Madrone, Pacific	Pine, White
	Birch, Gray	Manzanita,	Poison Ivy
	Birch, Paper	Greenleaf	Poison Oak
	Blackberry	Maple, Mountain	Poplar, Balsam
	Black Locust	Maple, Red	Redcedar,
	Blackgum	Maple, Silver	Eastern
	Boxelder	Maple, Sugar	Sassafras(?)
Brazilian	Cherry Black(?)	Maple, Vine	Sumac, Smooth(?)
Pepper	Cherry, Choke	Mesquite (1/3)	Sumac,
	Cherry, Pin	Mountain-Laurel	Staghorn(?)
	Cottonwood	Oak, Black(?)	Sweetgum
	Dogwood,	Oak, Blackjack(?)	Sycamore
	Flowering	Oak, Chestnut	Tanoak
	Dogwood,	Oak, Post(?)	Walnut
	Red-Osier	Oak, Red	Waxmyrtle
Elm,	Elm, American	Oak, Scarlet	Willow
Winged (1)	Gamberry	Oak, Water	Yaupon
	Guava	Oak, White	Yellow Poplar
	Hackberry	Olive, Autumn	
	Hazel	Olive, Russian	

(1) Some resprouting may occur. Maple, Bigleaf (1)
Maple, Stripped

DIRECTIONS FOR USE (Ready-To-Use, No Mixing Required)

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not use for manufacturing or formulating.

Do not apply this product through any type of irrigation system.

LOW VOLUME BASAL BARK TREATMENT: To control susceptible woody plants with stems less than 6 inches in basal diameter, apply PATHFINDER Herbicide with a backpack or knapsack sprayer using low pressure and a solid cone or flat fan nozzle. Spray the basal parts of brush and tree trunks in a manner which thoroughly wets the lower 12 to 15 inches of stems, including the root collar area, but not to the point of runoff. Herbicide concentration should vary with size and susceptibility of species treated. Apply at any time, including the winter months, except when snow or water prevent spraying to the ground line.

TREATMENT OF CUT STUMPS: To control resprouting, apply undiluted PATHFINDER Herbicide to wet the area adjacent to the cambium and bark around the entire circumference and the sides of freshly cut stumps. Sides of stumps should be thoroughly wetted down to the root collar area, but not to the point of runoff. Treatments may be applied throughout the year, except when snow or water prevent spraying to the ground line. Control may be reduced with treatment during periods of moisture stress as in late summer. Stumps should be cut so that they are approximately level to facilitate uniform PATHFINDER Herbicide coverage.

NOTE: These treatment methods may be used to treat susceptible woody species on range and permanent pasture land provided that no more than 1.5 gallons of PATHFINDER Herbicide are applied per acre. Large plants or species requiring higher rates PATHFINDER may not be completely controlled.

USE PRECAUTIONS

Apply this product only as specified on this label.

Do not apply PATHFINDER Herbicide directly to, or otherwise permit it to come into direct contact with grapes, tobacco, vegetable crops, flowers or other desirable broadleaf plants and do not permit spray mists containing it to drift onto them.

AVOID INJURIOUS SPRAY DRIFT: Applications should be made only when there is little or no hazard from spray drift. Very small quantities of spray, which may not be visible, may seriously injure susceptible plants. Do not spray when wind is blowing toward susceptible crops or ornamental plants near enough to be injured.

With ground equipment, spray drift can be reduced by using spray pressures no greater than are required to obtain adequate coverage; by using large droplet producing nozzle tips; and by spraying when wind velocity is low. Do not apply with nozzles that produce a fine droplet spray. Do not use a mistblower.

Do not apply on snow or frozen ground.

Untreated trees occasionally can be affected by movement of the herbicide through root grafting with the treated trees. Since this herbicide moves within the treated plant, do not use Pathfinder Herbicide on parts of a multiple stem plant if injury to the untreated portions (cut or standing stems) cannot be tolerated.

Do not apply on ditches used to transport irrigation water. Do not apply where runoff or irrigation water may flow onto agricultural land as injury to crops may result.

Do not graze lactating dairy animals or harvest hay from treated areas for one year following treatment.

Do not graze areas treated with more than 1.5 gallons PATHFINDER per acre for one year following treatment.

Withdraw livestock from forage treated with 1.5 gallons or less of PATHFINDER Herbicide at least 3 days before slaughter during the year of treatment.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store above 28°F or agitate before use.

DISPOSAL:

Prohibitions - Open dumping is prohibited.

Pesticide Disposal - Pesticide, spray mixture, or residue that cannot be used according to label instructions must be disposed of according to applicable federal, state, or local procedures.

Container Disposal - Triple rinse (or equivalent) with 10 drops or more of a liquid hand soap and water or an oil based product such as kerosene or diesel fuel and spray residue on undesirable vegetation in target area. Offer containers for recycling or reconditioning where allowed, or puncture and dispose of in a sanitary landfill, or by incineration if approved by state and local procedures.

General - Consult federal, state, or local disposal authorities for approved alternative procedures.

Be sure that use of this product conforms to all applicable regulations.

INSERT B

Container Disposal for Refillable Containers: Close all openings which have been opened during use and replace all caps. Return the empty container to a collection site designated by Dow Chemical. If the container has been damaged and cannot be returned according to the recommended procedures, contact the Dow Chemical Customer Service Center at 1-800-238-1470 to obtain proper handling instructions.

WARRANTY LIMITATIONS AND DISCLAIMER

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions therein under normal conditions of use. THIS IS THE ONLY WARRANTY MADE ON THIS PRODUCT. NO OTHER EXPRESS AND NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE IS MADE OUTSIDE OF THIS LABEL.

Therefore, neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), under abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes, etc.) or under conditions not reasonably foreseeable to or beyond the control of seller.

When buyer or user suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), buyer or user must promptly notify in writing DowElanco of any claims to be eligible to receive either remedy given below. The EXCLUSIVE REMEDY OF THE BUYER OR USER and the LIMIT OF LIABILITY of DowElanco or any other seller will be one of the following, at the election of DowElanco:

(1) Refund of purchase price paid by buyer or user for product bought, or

(2) Replacement of amount of product used

The seller will not be liable for consequential or incidental damages or losses

The terms of this Warranty Limitations And Disclaimer cannot be varied by any written or verbal statements or agreements. Any employee or sales agent of the seller is not authorized to vary or exceed the terms of this Warranty Limitations And Disclaimer in any manner.

12167-L1

390

STREAMLINE BASAL BARK TREATMENT (Southern

States): To control or suppress susceptible woody plants for conifer release, ~~only~~ apply DowElanco BASAL Herbicide with a backpack or knapsack sprayer using equipment which provides a directed straight-stream spray. Apply sufficient spray to one side of stems less than 3 inches in basal diameter to form a treated zone that is 6 inches in height. When the optimum amount of spray mixture is applied, the treated zone should widen to encircle the stem within approximately 30 minutes. Treat both sides of stems which are 3 to 4 inches in basal diameter. Direct the spray at bark that is approximately 12 to 24 inches above ground. Pines (loblolly, slash, shortleaf, and Virginia) up to 2 inches in diameter breast height (dbh) can be controlled by directing the spray at the point approximately 4 feet above ground. Best results are achieved when applications are made to young vigorously growing stems which have not developed the thicker bark characteristic of slower growing, under-story trees in older stands. This technique is not recommended for scrub and live oak species, including blackjack, turkey, post, live, bluejack and laurel oaks. Apply from approximately 6 weeks prior to hardwood leaf expansion in the spring until approximately 2 months after leaf expansion is completed. Do not apply when snow or water prevent spraying at the desired height above ground level. or in rangeland and pasture

INSERT 2

DowElanco

Indianapolis, Indiana 46268 U.S.A.

*Trademark of DowElanco