

PM23
62719-178
12-18

DowElanco IDXXXX

GALLERY TURF FERTILIZER

A Balanced Turf Fertilizer Containing GALLERY, a Selective Preemergence Herbicide for Control of Certain Broadleaf Weeds and Suppression of Annual Grasses in Established Turfgrasses.

Active Ingredients:

isoxaben*: N-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]- 2,6-dimethoxybenzamide and isomers.....	0.38%
Inert Ingredients.....	49.62%

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen.....	18.0%
Available Phosphoric Acid (P ₂ O ₅).....	5.0%
Soluble Potash (K ₂ O).....	9.0%

NET WEIGHT - 25 POUNDS

Contains 0.09 pounds active ingredient per 25 pound bag.

GALLERY(R) - (isoxaben, DowElanco)

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back or container/side panel for additional precautionary statements.

E.P.A. Reg. No. 62719-XXX


E.P.A. Est. No. XXXXXXX

FN9063

DowElanco

4040 Vincennes Circle, Suite 601

Indianapolis, Indiana 46268, U.S.A.



pg 2 of 6

PRECAUTIONARY STATEMENTS

Keep Out of Reach of Children

Hazards to Humans and Domestic Animals

CAUTION

Causes eye irritation. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust and contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

First Aid

If in eyes: Flush eyes with plenty of water. Get medical attention if irritation develops.


If swallowed: Do not induce vomiting. Call a physician or Poison Control Center. If available, administer activated charcoal (6-8 heaping teaspoonfuls) with a large quantity of water. Do not induce vomiting or give anything by mouth to an unconscious person. Immediately transport to a medical care facility and see a physician.

If inhaled: Remove individual to fresh air. Get medical attention if breathing difficulty occurs. If not breathing, give artificial respiration, preferably cardiopulmonary resuscitation and get medical attention immediately.

If on Skin: Wash with plenty of soap and water. Get medical attention if irritation develops.

Environmental Hazards

Do not apply directly to water or wetlands. Drift or inadvertent application to areas adjacent to treated turfgrasses may result in reduced germination or emergence of nontarget plants. Do not contaminate water when disposing of equipment washwaters.



DIRECTIONS FOR USE

Read all directions carefully before applying this product.

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

This turf fertilizer product contains GALLERY, a preemergence herbicide for control of certain broadleaf weeds and suppression of annual grasses in established turfgrasses. Apply in late summer to early fall, or in early spring, prior to emergence of target weeds.

GENERAL USE PRECAUTIONS


GALLERY Turf Fertilizer does not control established weeds. Existing weeds should be controlled with postemergence herbicides. To insure optimal fertilizer and herbicidal response, water treated areas as soon as possible after application. A single rainfall or sprinkler irrigation of 0.5 inch or more is required to activate this product.

Do not apply GALLERY Turf Fertilizer to non-turf areas such as flower beds or gardens. Do not allow "drift" from rotary-type applicators to reach such areas. GALLERY prevents seed emergence and may injure bedding plants.

Do not apply GALLERY Turf Fertilizer to dichondra.

Do not apply GALLERY Turf Fertilizer to golf course putting greens.


Users who wish to use GALLERY Turf Fertilizer on turfgrass species not recommended on this label may determine the suitability for such uses by treating a small area at a recommended rate. Prior to treating a larger area, the treatment should be observed for any sign of herbicidal injury for 30 to 60 days to determine if the treatment is safe to the target turf species. THE USER ASSUMES RESPONSIBILITY FOR ANY CROP DAMAGE OR OTHER LIABILITY RESULTING FROM USE OF GALLERY ON TURF SPECIES NOT RECOMMENDED ON THIS LABEL.



GALLERY TURF FERTILIZER IS RECOMMENDED FOR USE ON THE FOLLOWING TURFGRASS SPECIES.

Common Name	Scientific Name
ESTABLISHED COOL SEASON GRASSES	
Bentgrass, colonial	<i>Agrostis tenuis</i>
Bentgrass, creeping	<i>Agrostis stolonifera</i>
Bluegrass, Kentucky	<i>Poa pratensis</i>
Fescue, chewings	<i>Festuca rubra</i> var. <i>commutata</i>
Fescue, creeping red	<i>Festuca rubra</i>
Fescue, sheep	<i>Festuca ovina</i>
Fescue, tall	<i>Festuca arundinaceae</i>
Ryegrass, perennial	<i>Lolium perenne</i>

ESTABLISHED WARM SEASON GRASSES	
Bahia grass	<i>Paspalum notatum</i> var. <i>Saurae parodi</i>
Bermudagrass	<i>Cynodon dactylon</i>
Centipedegrass	<i>Eremochloa ophiuroides</i>
Fescue, tall	<i>Festuca arundinaceae</i>
St. Augustinegrass	<i>Stenotaphrum secundatum</i>
Zoysiagrass	<i>Zoysia japonica</i>
Zoysiagrass	<i>Zoysia tenuifolia</i>



APPLICATION

EQUIPMENT AND APPLICATION TECHNIQUES

Apply GALLERY Turf Fertilizer using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to manufacturer's directions. Check frequently to be sure equipment is operating properly and distributing uniformly. Avoid use of spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or crop injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. A splash board mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.


BROADCAST APPLICATION RATES

Apply GALLERY Turf Fertilizer at a broadcast rate of 200 to 250 pounds per acre (4.6 to 5.7 pounds per 1000 square feet). A minimum rate of 200 pounds per acre is recommended for optimum results.

WEEDS CONTROLLED

Weeds controlled by GALLERY Turf Fertilizer applied prior to weed germination at 200 to 250 pounds per acre.

Common Name	Scientific Name
Burweed, lawn	<i>Soliva pterosperma</i>
Buttercup	<i>Ranunculus spp.</i>
Celery, wild	<i>Apium leptophyllum</i>
Chickweed, common	<i>Stellaria media</i>
Chickweed, mouseear	<i>Cerastium vulgatum</i>
Foalsparsley	<i>Aethusa cynapium</i>
Geranium, Carolina	<i>Geranium carolinianum</i>
Henbit	<i>Lamium amplexicaule</i>
Knotweed, prostrate	<i>Polygonum aviculare</i>
Mallow, little	<i>Malva parviflora</i>
Parsley - piert	<i>Alchemilla arvensis</i>
Pennywort, lawn	<i>Hydrocotyle sithorpioides</i>
Pineappleweed	<i>Matricaria matricarioides</i>
Speedwell, thymeleaf	<i>Veronica serpyllifolia</i>
Spurge, prostrate	<i>Euphorbia humistrata</i>
Spurge, spotted	<i>Euphorbia maculata</i>
Spurry, corn	<i>Spergula arvensis</i>
Woodsorrel, creeping	<i>Oxalis corniculata</i>
Woodsorrel, yellow	<i>Oxalis stricta</i>



Weeds partially controlled by a single application; or, controlled by repeat applications of GALLERY Turf Fertilizer applied prior to weed germination at 200 to 250 pounds per acre.

Common Name	Scientific Name
Clover spp. (1)	<i>Trifolium spp.</i>
Clover, white (1)	<i>Trifolium repens</i>
Dandelion (1)	<i>Taraxacum officinale</i>
Medic, black	<i>Medicago lupulina</i>
Plantain, broadleaf	<i>Plantago major</i>
Plantain, buckhorn	<i>Plantago lanceolata</i>
Plantago, slender	<i>Plantago elongata</i>

1. From seed only

PS-1096

Weeds suppressed or partially controlled by GALLERY Turf Fertilizer applied prior to weed germination at 200 to 250 pounds per acre.

Common Name	Scientific Name
Bluegrass, annual	Poa annua
Crabgrass	Digitaria spp.
Ryegrass, annual (Italian)	Lolium multiflorum

DISCLAIMER OF WARRANTIES

DowElanco makes no warranties concerning this product or its use, which extend beyond the description of the label. All other warranties, expressed or implied, including but not limited to, and implied warranties of merchantability and fitness for a particular purpose, are disclaimed. All statements concerning this product apply only when used as directed. This disclaimer of warranties does not apply where prohibited by operation of law.

LIMITATION OF DAMAGES

The liability of DowElanco, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the amount of the purchase price of this product and under no circumstances shall DowElanco be liable for special, indirect or consequential damages. This limitation of damages does not apply where prohibited by operation of law.

Copyright(c) - DowElanco - 1989

DowElanco
40 Vincennes Circle, Suite 601
Indianapolis, Indiana 46268, U.S.A.

Revised 12/14/89

