



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

May 7, 2019

Jennifer Hughes  
Regulatory Manager  
Dow AgroSciences, LLC  
9330 Zionsville Rd., 308/2E  
Indianapolis, IN 46268-1054

Subject: Notification per PRN 98-10 – Update MOA and Resistance Management Sections per  
PR Notices 2017-1 & 2017-2  
Product Name: Gallery Turf Fertilizer  
EPA Registration Number: 62719-178  
Application Date: November 29, 2018  
Decision Number: 550649

Dear Ms. Hughes:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped “Notification” and will be placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you have any questions, please contact me at [ondish.mindy@epa.gov](mailto:ondish.mindy@epa.gov) or at (703)605-0723.

Sincerely,

A handwritten signature in cursive script that reads "Mindy Ondish".

Mindy Ondish  
Product Manager 23  
Herbicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

**[Sub Label A: Professional Uses]**

<b>ISOXABEN</b>	<b>GROUP</b>	<b>21</b>	<b>HERBICIDE</b>
-----------------	--------------	-----------	------------------

# Gallery<sup>®</sup> Turf Fertilizer

## HERBICIDE

**A turf fertilizer containing Gallery, a preemergence herbicide for control of certain broadleaf weeds and suppression of annual grasses in landscape ornamentals, container grown ornamentals, field grown ornamentals, and home lawns.**

**Intended for use by commercial pest control operators**

**[XX Lbs] Covers up to XX,000 sq ft (XXX' x XX')**  
and/or

**This bag covers XX,000 sq ft**

Not for sale, distribution or use in New York State

<b>Group</b>	<b>21</b>	<b>HERBICIDE</b>
--------------	-----------	------------------

### NOTIFICATION

62719-178

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

05/07/2019

**Active Ingredient:**

isoxaben: N-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers.....	0.38%
Other Ingredients .....	99.62%
Total .....	100.00%

**Guaranteed Fertilizer Analysis: XX-XX-XX †**

Total Nitrogen (N).....	XX%
XX% Ammoniacal Nitrogen	
XX% Nitrate Nitrogen	
XX% Water-Insoluble Nitrogen*	
XX% Urea Nitrogen**	
XX% Slowly available water soluble Nitrogen.	
XX% Other Water Soluble Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )*** .....	XX%
Soluble Potash (K <sub>2</sub> O).....	XX%
Calcium (Ca) .....	XX%
Magnesium (Mg) .....	XX%
XX% Water Soluble Magnesium (Mg)	
Sulfur****	
XX% Combined Sulfur (S)	
Boron (B) .....	XX%
Chlorine (Cl) [Max, Not more than] .....	XX%
Cobalt (Co).....	XX%
Copper (Cu).....	XX%
Iron (Fe).....	XX%
XX% Chelated Iron (Fe)	

XX% Water Soluble Iron (Fe) Manganese (Mn)XX%  
 Water Soluble Manganese (Mn) ..... XX%  
 Molybdenum (Mo) ..... XX%  
 Nickel (Ni)..... XX%  
 Sodium (Na)..... XX%  
 Zinc (Zn)..... XX%  
 XX% Chelated Zinc (Zn)

\*[XX]% Water insoluble nitrogen from biosolids.

\*\*[XX]% coated slow release Nitrogen (N)

\*\*[XX]% Slowly available [urea] nitrogen from [polycoated] [polymer/sulfur] coated urea

\*\*[XX]% Slowly available water soluble nitrogen

\*\*[XX]% Slowly available nitrogen from methylene ureas

\*\*[XX]% Slowly available nitrogen from ureaform

\*\*[XX]% Slowly available nitrogen from BioSolids

\*\*[XX]% Stabilized urea nitrogen or [urea] nitrogen stabilized with dicyandiamide

\*\*\*[XX]% Free Sulfur (S)

Derived From: [XX]<sup>†</sup>

Primary plant foods derived from: [XX]<sup>†</sup>

Secondary plant foods derived from [XX]<sup>†</sup>

Iron derived from: [XX]<sup>†</sup>

Density – XX lb./cu.ft.

**[Editor's Note: the following information may be added in the boxes above, this footnote must be deleted from the label after completion of the above Fertilizer Analysis data]**

<sup>†</sup> Urea, methylene urea, sulfur coated urea, polymer coated urea, poly[mer] coated sulfur coated urea, urea formaldehyde, Ureaform, ammonium sulfate, corn gluten, bio-solids (Milorganite, Nutri-Pel, composted poultry manure), activated sewage sludge, diammonium phosphate, monoammonium phosphate, magnesium ammonium phosphate hexa-hydrate, triple super phosphate, ammoniated phosphates, sulfate of potash, muriate of potash, sulfate of potash-magnesia, potassium nitrate, copper oxide, copper sulfate, iron/ferric oxide, iron/ferrous sucrate, iron/ferrous oxy-sulfate, iron/ferrous sulfate, manganese oxide, molybdcic oxide, manganese sulfate, manganese sucrate, zinc oxide, zinc sulfate, sodium borate, boron, sulfur, sodium molybdate, stabilized nitrogen, potassium magnesium sulfate, ferrous sulfate monohydrate, magnesium sulfate or limestone.

## Keep Out Of Reach Of Children

# CAUTION

If this product is being sold in a state requiring the publication of information regarding its metal content, information regarding the contents and levels of metals in this product is available at the following website: <http://www.aapfco.org/metals.html>

Information regarding the contents and levels of metals in this product is available on the Internet at:

- California Regulatory Site: <https://apps1.cdfa.ca.gov/FertilizerProducts/>
- Oregon Regulatory Site: [http://oda.state.or.us/dbs/heavy\\_metal/search.lasso](http://oda.state.or.us/dbs/heavy_metal/search.lasso)
- Washington Regulatory Site: <http://agr.wa.gov/> or <http://agr.wa.gov/PestFert/Fertilizers/ProductDatabase.aspx>
- <http://www.aapfco.org/metals.html>
- <http://www.regulatory-info-lpi.com>

It is a violation of law in Maryland and New Jersey to apply this product to turf at a rate that results in greater than 0.7 lb per 1,000 sq.ft. of readily available nitrogen or more than 0.9 lb per 1,000 sq.ft. of total nitrogen with a least 20% of which shall consist of slow release nitrogen per application. Additional restrictions apply to total nitrogen applied to turf per season. Other state and local restrictions may apply. [Maryland or New Jersey Applications]

No more than 0.7 lb per 1,000 sq.ft. of readily available nitrogen during any given 30 day period. No more than 0.9 lb (cool season grasses) or 1.0 lb (warm season grasses) per 1,000 sq.ft. of total nitrogen with a least 20% of which shall consist of slow release nitrogen during any given 30 day period. [Virginia Applications]

Do not apply near water storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back onto your lawn and garden. [Maryland Applications]

This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under §8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; 2) establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) reestablishing or repairing a turf area. [Maryland Applications which contain Phosphorus] Not for use on established lawns or grass in MD. [If phos >5% and is not classified as a "starter" fertilizer]

In Virginia and New Jersey this lawn/turf fertilizer contains phosphorous and is only for nonagricultural use on (i) turf during its first growing season, (ii) on turf areas being repaired or renovated, and (iii) on turf where a soil test performed within the last 3 years indicates a phosphorous deficiency. This fertilizer is not for the routine maintenance of turf. [Virginia and New Jersey Applications which contain Phosphorus]

Apply only to actively growing turf. Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands in the driveway, sidewalk, or street, back onto your lawn. Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use. [FL Language for Specialty Products]

The maximum rate of application of this product must not exceed the lesser of the applicable rate determined from the [Table [XX]: Use Rates or Broadcast Application Rates], and the applicable rate determined in the State Nutrient Restrictions table.

#### STATE NUTRIENT RESTRICTIONS

State	Readily Available Nitrogen* lbs/1,000 sq ft	Total Nitrogen lbs/1,000 sq ft
MD, NH, VA	0.7	0.9
NJ	0.7	1.0
XXX	XXX	XXX

\* Nitrogen present beyond the amount show as Readily Available must be in the form of Slow-Release nitrogen

---

## Precautionary Statements

---

### Hazards to Humans and Domestic Animals

## CAUTION

**Causes Eye Irritation • Harmful If Swallowed, Inhaled Or Absorbed Through The Skin**

**Avoid breathing dust and contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.**

### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### First Aid

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or inadvertent application to areas adjacent to treated turfgrasses may result in reduced germination or emergence of nontarget plants. Do not contaminate water when disposing of equipment washwaters.

**Ground Water Advisory:** This pesticide has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

**Surface Water Advisory:** This pesticide may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soil and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of isoxaben from runoff water and sediment.

### **Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

#### **(Storage and Disposal for rigid containers 5 gal or less)**

##### **Storage and Disposal**

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

#### **(Storage and Disposal for nonrigid containers any size)**

##### **Storage and Disposal**

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

#### **(Storage and Disposal for refillable rigid containers greater than 5 gal)**

##### **Storage and Disposal**

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container.

Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**(Storage and Disposal for nonrefillable rigid containers larger than 5 gal)**

**Storage and Disposal**

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**Refer to label booklet for Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994 [or 1-800-XXX-XXXX].

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-178

EPA Est. No. \_\_\_\_\_

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

**Produced by/for**  
**Manufactured by/for:**  
**Distributed by/for:**  
**Guaranteed by:**

**Dow AgroSciences LLC**  
**9330 Zionsville Road**  
**Indianapolis, IN 46268**

**NET WEIGHT** \_\_\_\_\_

(cover/shipping container):

<b>ISOXABEN</b>	<b>GROUP</b>	<b>21</b>	<b>HERBICIDE</b>
-----------------	--------------	-----------	------------------

# Gallery® Turf Fertilizer

## HERBICIDE

**A turf fertilizer containing Gallery, a preemergence herbicide for control of certain broadleaf weeds and suppression of annual grasses in landscape ornamentals, container grown ornamentals, field grown ornamentals, and home lawns.**

**Intended for use by commercial pest control operators**

**[XX lbs] Covers up to XX,000 sq ft (XXX' x XX')**  
and/or

**This bag covers XX,000 sq ft**

Not for sale, distribution or use in New York State

<b>Group</b>	<b>21</b>	<b>HERBICIDE</b>
--------------	-----------	------------------

**Active Ingredient:**

isoxaben: N-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]- 2,6-dimethoxybenzamide	0.38%
and isomers.....	0.38%
Other Ingredients .....	99.62%
Total .....	100.00%

**Guaranteed Fertilizer Analysis: XX-XX-XX †**

Total Nitrogen (N).....	XX%
XX% Ammoniacal Nitrogen	
XX% Nitrate Nitrogen	
XX% Water-Insoluble Nitrogen*	
XX% Urea Nitrogen**	
XX% Slowly available water soluble Nitrogen.	
XX% Other Water Soluble Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )*** .....	XX%
Soluble Potash (K <sub>2</sub> O).....	XX%
Calcium (Ca) .....	XX%
Magnesium (Mg) .....	XX%
XX% Water Soluble Magnesium (Mg)	
Sulfur****	
XX% Combined Sulfur (S)	
Boron (B) .....	XX%
Chlorine (Cl) [Max, Not more than] .....	XX%
Cobalt (Co).....	XX%
Copper (Cu).....	XX%
Iron (Fe).....	XX%
XX% Chelated Iron (Fe)	
XX% Water Soluble Iron (Fe) Manganese (Mn)XX%	



Water Soluble Manganese (Mn) ..... XX%  
 Molybdenum (Mo) ..... XX%  
 Nickel (Ni)..... XX%  
 Sodium (Na)..... XX%  
 Zinc (Zn)..... XX%  
 XX% Chelated Zinc (Zn)

\*[XX]% Water insoluble nitrogen from biosolids.

\*\*[XX]% coated slow release Nitrogen (N)

\*\*[XX]% Slowly available [urea] nitrogen from [polycoated] [polymer/sulfur] coated urea

\*\*[XX]% Slowly available water soluble nitrogen

\*\*[XX]% Slowly available nitrogen from methylene ureas

\*\*[XX]% Slowly available nitrogen from ureaform

\*\*[XX]% Slowly available nitrogen from BioSolids

\*\*[XX]% Stabilized urea nitrogen or [urea] nitrogen stabilized with dicyandiamide

\*\*\*\*[XX]% Free Sulfur (S)

Derived From: [XX]<sup>†</sup>

Primary plant foods derived from: [XX]<sup>†</sup>

Secondary plant foods derived from [XX]<sup>†</sup>

Iron derived from: [XX]<sup>†</sup>

Density – XX lb./cu.ft.

**[Editor's Note: the following information may be added in the boxes above, this footnote must be deleted from the label after completion of the above Fertilizer Analysis data]**

<sup>†</sup> Urea, methylene urea, sulfur coated urea, polymer coated urea, poly[mer] coated sulfur coated urea, urea formaldehyde, Ureaform, ammonium sulfate, corn gluten, bio-solids (Milorganite, Nutri-Pel, composted poultry manure), activated sewage sludge, diammonium phosphate, monoammonium phosphate, magnesium ammonium phosphate hexa-hydrate, triple super phosphate, ammoniated phosphates, sulfate of potash, muriate of potash, sulfate of potash-magnesia, potassium nitrate, copper oxide, copper sulfate, iron/ferric oxide, iron/ferrous sucate, iron/ferrous oxy-sulfate, iron/ferrous sulfate, manganese oxide, molybdcic oxide, manganese sulfate, manganese sucate, zinc oxide, zinc sulfate, sodium borate, boron, sulfur, sodium molybdate, stabilized nitrogen, potassium magnesium sulfate, ferrous sulfate monohydrate, magnesium sulfate or limestone.

## Keep Out Of Reach Of Children

# CAUTION

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

**Refer to inside of label booklet for additional precautionary information including Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994 [or 1-800-XXX-XXXX].

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-178

EPA Est. No. \_\_\_\_\_

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

**Produced by/for:**  
**Manufactured by/for:**  
**Distributed by/for:**  
**Guaranteed by:**

**Dow AgroSciences LLC**  
**9330 Zionsville Road**  
**Indianapolis, IN 46268**

**NET WEIGHT \_\_\_\_\_**

(Page 1 through end):

---

## Precautionary Statements

---

### Hazards to Humans and Domestic Animals

**Causes Eye Irritation • Harmful If Swallowed, Inhaled Or Absorbed Through The Skin**

**Avoid breathing dust and contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.**

### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### First Aid

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or inadvertent application to areas adjacent to treated turfgrasses may result in reduced germination or emergence of nontarget plants. Do not contaminate water when disposing of equipment washwaters.

**Ground Water Advisory:** This pesticide has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

**Surface Water Advisory:** This pesticide may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soil and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after

application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of isoxaben from runoff water and sediment.

---

## Directions for Use

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all directions carefully before applying this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements of this label about personal protective equipment (PPE) and restricted entry interval. The requirements in the box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow workers to enter during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material.
- Shoes plus socks

### Non-Agricultural Use Requirements

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**Entry Restrictions for Non-WPS Uses:** When this product is applied to turf and ornamental plantings in landscape settings, do not enter or allow others to enter the treated area until dusts have settled. If soil incorporation is required after the application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

### Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

#### Nonrefillable rigid containers 5 gallons or less:

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate

into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**Nonrefillable nonrigid containers:**

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**Refillable rigid containers larger than 5 gal:**

**Container Handling:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**Nonrefillable rigid containers larger than 5 gal:**

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

## Product Information

This turf fertilizer product contains Gallery®, a preemergence herbicide for control of certain broadleaf weeds and suppression of annual grasses in established turfgrasses, landscape ornamentals, container grown ornamentals, and field grown ornamentals. Apply in late summer to early fall, in early spring, or anytime prior to germination of target weeds.

## Use Precautions

Gallery Turf Fertilizer controls weeds germinating from seed. Gallery Turf Fertilizer does not control established weeds. Existing weeds should be controlled with postemergence herbicides. To ensure optimal fertilizer and herbicidal response, water treated areas as soon as possible after application. A single rainfall or sprinkler irrigation of 0.5 inch or more is required to activate this product.

### Use Restrictions

Do not apply more than 833 lb of Gallery Turf Fertilizer per acre.

**Chemigation:** Do not apply Gallery Turf Fertilizer through any type of irrigation system.

Not for sale, distribution or use in New York State.

Do not apply by air.

Do not apply Gallery Turf Fertilizer to turfgrass grown for seed.

### Treatment of Turfgrass or Ornamental Species not Listed on the Label for Gallery Turf Fertilizer

Users who wish to use Gallery Turf Fertilizer on a plant species not listed on this label must determine the suitability for such use by treating a small area or a small number of plants at a specified rate. Prior to treating a larger area, observe the treated area/plants for any sign of herbicidal injury during 30 to 60 days of typical growing conditions.

### Application Instructions

Apply Gallery Turf Fertilizer using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or plant injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. When using a drop-type spreader, a splashboard mounted below the release point will provide more even granule distribution.

**[Editor's note: With available data Supplemental Distributor may add the following statement.]**  
**These settings were calibrated and field tested. However, age, condition of spreader speed of operations and evenness of terrain may require slightly different settings for desired coverage.**

Check equipment frequently to verify calibration and proper functioning.

**[Editor's note: Supplemental Distributor may add a table listing recommended application equipment and calibration settings.]**

Brand and Model of Spreader	Application Rates	Calibration Settings at Specified Rate					
		lb/1000 sq ft	4.6	XX	XX	XX	XX
	lb/acre	200	XX	XX	XX	XX	250
	Lb nitrogen / 1000 sq ft	XX	XX	XX	XX	XX	XX
XXXXXXX		XX	XX	XX	XX	XX	XX

[AND/OR]

Calibration Settings			
Rate	lbs. Product per 1,000 sq ft	lbs. Nitrogen per 1,000 sq ft	lbs A.I per acre

<b>Low</b>	<b>4.6</b>	<b>XX</b>	<b>XX</b>
<b>Medium</b>	<b>XX</b>	<b>XX</b>	<b>XX</b>
<b>High</b>	<b>5.7</b>	<b>XX</b>	<b>XX</b>

[AND/OR]

Brand and Model of Spreader	Calibration Settings (lb/1000 sq ft)*		
	Low	Medium	High
XXXXXXX	XX	XX	XX
XXXXXXX	XX	XX	XX
XXXXXXX	XX	XX	XX

[\*These settings are approximate for the application rate shown per 1,000 square feet. You may need to adjust the spreader setting depending on walking speed, spreader condition and product]

### Broadcast Application Rates

Apply Gallery Turf Fertilizer at a broadcast rate of 200 to 250 lb per acre (4.6 to 5.7 lb per 1000 sq ft). A minimum rate of 200 lb per acre (4.6 lb per 1000 sq ft) will provide optimum results. Use the higher rate in the rate range where heavy weed infestation is anticipated.

### Weeds Controlled or Suppressed

#### Weeds controlled when applied at 200 to 250 lb per acre (4.6 to 5.7 lb per 1000 sq ft)

##### Common Name

burweed, lawn  
buttercup  
celery, wild  
chickweed, common  
chickweed, mouseear  
foolsparsely  
geranium, Carolina  
henbit  
knotweed, prostrate  
mallow, little  
parsley - piert  
pennywort, lawn  
pineappleweed  
speedwell, thymeleaf  
spurge, prostrate  
spurge, spotted  
spurry, corn  
woodsorrel, creeping  
woodsorrel, yellow

##### Scientific Name

*Soliva pterosperma*  
*Ranunculus spp.*  
*Apium leptophyllum*  
*Stellaria media*  
*Cerastium vulgatum*  
*Aethusa cynapium*  
*Geranium carolinianum*  
*Lamium amplexicaule*  
*Polygonum aviculare*  
*Malva parviflora*  
*Alchemilla arvensis*  
*Hydrocotyle sibthorpioides*  
*Matricaria matricarioides*  
*Veronica serpyllifolia*  
*Euphorbia humistrata*  
*Euphorbia maculata*  
*Spergula arvensis*  
*Oxalis corniculata*  
*Oxalis stricta*

#### Weeds partially controlled by a single application or controlled by repeat applications when applied at 200 to 250 lb per acre (4.6 to 5.7 lb per 1000 sq ft)

##### Common Name

clover, spp.<sup>1</sup>  
clover, white<sup>1</sup>  
dandelion<sup>1</sup>  
medic, black  
plantain, broadleaf  
plantain, buckhorn  
plantago, slender

##### Scientific Name

*Trifolium spp.*  
*Trifolium repens*  
*Taraxacum officinale*  
*Medicago lupulina*  
*Plantago major*  
*Plantago lanceolata*  
*Plantago elongata*

<sup>1</sup>From seed only

**Weeds suppressed or partially controlled when applied at 200 to 250 lb per acre (4.6 to 5.7 lb per 1000 sq ft)**

**Common Name**

bluegrass, annual  
crabgrass  
ryegrass, annual (Italian)

**Scientific Name**

*Poa annua*  
*Digitaria spp.*  
*Lolium multiflorum*

## **WEED RESISTANCE MANAGEMENT**

Gallery Turf Fertilizer, which contains the active ingredient isoxaben is a Group 21 herbicide based on the mode of action classification system of the Weed Science Society of America.

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is a best practice. A diversified weed management program may include the use of multiple herbicides with different modes of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistance.

The continued effectiveness of this product depends on the successful implementation of a weed resistance management program.

To aid in the prevention of developing weeds resistant to this product, users should:

- Scout fields before application to ensure herbicides and rates will be appropriate for the weed species and weed sizes present.
- Start with a clean field, using either a burndown herbicide application or tillage.
- If using post-emergence herbicides or tank mixes, control weeds early when they are relatively small.
- Apply full rates of Gallery Turf Fertilizer for the most difficult to control weed in the field at the specified time to minimize weed escapes.
- Scout fields after application to detect weed escapes or shifts in control of weed species.
- Control weed escapes before they reproduce by seed or proliferate vegetatively.
- Report any incidence of non-performance of this product against a particular weed to your local company representative, local retailer, or county extension agent.
- Contact your local company representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective modes of action for each target weed.
- If resistance is suspected, treat weed escapes with an herbicide having a mode of action other than Group 21 and/or use nonchemical methods to remove escapes, as practical, with the goal of preventing further seed production.
- Suspected herbicide-resistant weeds may be identified by these indicators:
  - Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
  - A spreading patch of non-controlled plants of a particular weed species; and
  - Surviving plants mixed with controlled individuals of the same species.

Additionally, users should follow as many of the following herbicide resistance management practices as is practical:

- Use a broad spectrum herbicide with other mode of action as a foundation in a weed control program, if appropriate.
- Utilize sequential applications of herbicides with alternative modes of action.



- Rotate the use of this product with non-Group 21 herbicides.
- Avoid making more than two sequential applications of Gallery Turf Fertilizer and any other Group 21 herbicides within a single growing season unless mixed with an herbicide with a different mode of action with an overlapping spectrum for the difficult-to-control weeds.
- Incorporate non-chemical weed control practices, such as mechanical cultivation, crop rotation, cover crops and weed-free crop seeds, as part of an integrated weed control program.
- Use good agronomic principles that enhance crop development and crop competitiveness.
- Thoroughly clean plant residues from equipment before leaving fields suspected to contain resistant weeds.
- Manage weeds in and around fields to reduce weed seed production.

### **Weed Resistance Management:**

~~Isoxaben, the active ingredient in this product, is a Group 21 herbicide based on the mode of action classification system of the Weed Science Society of America. Any weed population may contain or develop plants resistant to Group 21 herbicides. Resistant weeds may dominate the weed population if these herbicides are used repeatedly in the same field. Such resistant weed plants may not be effectively managed using Group 21 herbicides but may be effectively managed utilizing other herbicides alone or in mixtures from a different herbicide Groups that are labeled for control of these weeds and/or by using cultural or mechanical practices. However, an herbicide mode of action classification by itself may not adequately address specific weeds that are resistant to specific herbicides. Consult your local company representative, state cooperative extension service, professional consultants or other qualified authorities to determine appropriate actions for treating specific resistant weeds or to report herbicide failures.~~

### **Best Management Practices:**

~~Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is recommended. A diversified weed management program may include the use of multiple herbicides with different modes of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistant weeds. It is recommended to scout for weeds before Gallery Turf Fertilizer application for identification and growth stage and after application to facilitate the early identification of weed shifts and/or weed resistance and thus provide direction on future weed management practices. One of the best ways to contain resistant weed populations is to implement measures to avoid allowing weeds to reproduce by seed or to proliferate vegetatively. Cleaning equipment between sites and avoiding movement of plant material between sites will greatly aid in reducing the spread of resistant weed seed.~~

## **Uses**

### **Turfgrass**

Gallery Turf Fertilizer may be used as a preemergence treatment for control of certain broadleaf weeds in established cool season and warm season turfgrasses.

Apply Gallery Turf Fertilizer anytime prior to germination of target weeds.

**Note:** Refer to the Product Information section of this label for general use precautions and information on application rates and weeds controlled prior to using this product.

### **Use Restrictions:**

Apply Gallery Turf Fertilizer only on established turfgrass. Apply to newly seeded turfgrass (including overseeded turfgrass) **only** after seedlings are established (three leaf stage and tillering) and well rooted.

Do not overseed established turfgrass sooner than 60 days following an application of Gallery Turf Fertilizer.

- Do not apply Gallery Turf Fertilizer to golf course putting greens.
- Do not apply Gallery Turf Fertilizer to dichondra.
- Florida Applications: Do not exceed [2] [3] applications per year on bahiagrass in the north.
- Florida Applications: Do not exceed [2] [3] [4] applications per year on bahiagrass in [central Florida] [the south].
- Florida Applications: Do not exceed [3] [4] [5] applications per year on bermudagrass in the north.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on bermudagrass in central Florida.
- Florida Applications: Do not exceed [5] [6] [7] applications per year on bermudagrass in the south.
- Florida Applications: Do not exceed [1] [2] applications per year on centipedegrass in the north.
- Florida Applications: Do not exceed [2] [3] applications per year on centipedegrass in [central Florida] [the south].
- Florida Applications: Do not exceed [2] [3] [4] applications per year on St. Augustinegrass in the north.
- Florida Applications: Do not exceed [2] [3] [4] [5] applications per year on St. Augustinegrass in central Florida.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on St. Augustinegrass in the south.
- Florida Applications: Do not exceed [3] [4] [5] applications per year on zoysiagrass in the north.
- Florida Applications: Do not exceed [3] [4] [5] [6] applications per year on zoysiagrass in central Florida.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on zoysiagrass in the south.

**Gallery Turf Fertilizer may be used on the following turfgrass species:**

#### **Established Cool Season Turfgrass**

##### **Common Name**

bentgrass, colonial  
bentgrass, creeping  
bluegrass, Kentucky  
fescue, chewings  
fescue, creeping red  
fescue, sheep  
fescue, tall  
ryegrass, perennial

##### **Scientific Name**

*Agrostis tenuis*  
*Arostis stolonifera*  
*Poa pratensis*  
*Festuca rubra var. commutata*  
*Festuca rubra*  
*Festuca ovina*  
*Festuca arundinaceae*  
*Lolium perenne*

#### **Established Warm Season Turfgrass**

##### **Common Name**

bahiagrass  
bermudagrass  
centipedegrass  
fescue, tall  
St. Augustinegrass  
zoysiagrass  
zoysiagrass

##### **Scientific Name**

*Paspalum notatum var. Saurae parodi*  
*Cynodon dactylon*  
*Eremochloa ophiuroides*  
*Festuca arundinaceae*  
*Stenotaphrum secundatum*  
*Zoysia japonica*  
*Zoysia tenuifolia*

#### **Ornamental Plantings**

Gallery Turf Fertilizer may be used as a preemergence treatment for control of certain broadleaf weeds in landscape ornamentals, container grown ornamentals, and field grown ornamentals.

Apply Gallery Turf Fertilizer anytime prior to germination of target weeds or immediately after cultivation.

**Note:** Refer to the Product Information section of this label for general use precautions and information on application rates and weeds controlled prior to using this product.

**Use Restrictions:**

To avoid possible plant injury, **do not** apply Gallery Turf Fertilizer to:

- nursery: seedling beds, cutting beds, or transplant beds
- unrooted liners or cuttings that have been planted in pots for the first time
- pots less than six inches wide
- bedding plants or areas where bedding plants will be planted or transplanted within 1 year after application

**Do not** apply Gallery Turf Fertilizer to newly transplanted ornamentals or nursery stock until soil or potting media has been settled by packing and irrigation or rainfall and no cracks are present or plant injury may occur.

Applications of Gallery Turf Fertilizer over the top of plants with newly forming buds may cause injury. Possible plant injury may be avoided by applying to the soil surface beneath ornamental plants.

When planting into a site treated with Gallery Turf Fertilizer in the previous 8 months, use untreated soil as fill around roots when replacing plants or injury may occur.

Injury to certain ornamental plants has been observed following application of Gallery Turf Fertilizer. To avoid plant injury, do not use Gallery Turf Fertilizer for weed control in the following ornamental plant species:

*Ajuga* spp. (bugle)  
*Brassica* spp. (mustard)  
*Echinacea purpurea* (purple coneflower)  
*Euonymus alatus 'compacta'* (dwarf burning bush)  
*Euphorbia* spp. (spurge)  
*Hydrangea* spp. (hydrangea)  
*Iberis* spp. (candytuft)  
*Juniperus horizontalis 'Prince of Wales'* (prince of wales juniper)  
*Melaleuca quinquenervia* (cajeput tree)  
*Rhododendron caroliniaum* (Carolina rhododendron)  
*Rhododendron catawbiense 'Roseum elegans'* (roseum elegans rhododendron)  
*Sedum* spp. (stonecrop)  
*Yucca recurvifolia* (green yucca)

**Gallery Turf Fertilizer may be used on the following established plant species (Note: Limitations on treatment methods):**

**Trees**

**Treatment Method:**  
**C = Container Grown**  
**F = Field Grown**

Scientific Name	Common Name	
<i>Abies balsamea</i>	fir, balsam	C, F
<i>Abies concolor</i>	fir, white	F
<i>Abutilon hybridum</i>	albus-flowering maple	C, F
	luteus-flowering maple	C, F
	roseus-flowering maple	C, F
	tangerine-flowering maple	C, F
	vesuvius red-flowering maple	F
<i>Acer ginnala</i>	flame maple	F
<i>Acer rubrum</i>	red maple	F
	red sunset maple	F
<i>Acer saccharinum</i>	silver maple	C, F

<i>Alsophila australis</i>	Australian tree fern	C, F
<i>Araucaria heterophylla</i>	norfolk island pine	C, F
<i>Archontophoenix cunninghamiana</i>	king palm	C, F
<i>Areacastrum romanzoffianum</i>	queen palm	C, F
<i>Bauhinia galpinii</i>	red bauhinia	C, F
<i>Betula nigra</i>	birch, river	C, F
<i>Betula papyrifera</i>	paper birch	F
<i>Brachychiton populneus</i>	bottle tree	C, F
<i>Bucida buceras</i>	black olive	F
<i>Ceratonia siliqua</i>	carob	F
<i>Cercis canadensis</i>	redbud	C, F
<i>Chamaecyparis obtusa</i>	filicoides-fernspray cypress	F
	gracilis-slender hinoki cypress	F
<i>Chamaecyparis pisifera</i>	sawara-false cypress	F
	squarrosa-moss cypress	F
<i>Chamaedorea cataractarum</i>	cat palm	F
	palm	C, F
<i>Chamaedorea costaricana</i>	palm	C, F
<i>Chamaedorea elegans</i>	parlor palm	C, F
<i>Chamaerops humilis</i>	Mediterranean fan palm	C, F
<i>Cornus florida</i>	cloud nine dogwood	C, F
	dogwood, flowering	C, F
<i>Crataegus viridis</i>	green hawthorn	F
<i>Cryptomeria japonica</i>	cryptomeria, Japanese	C, F
<i>Cupaniopsis anacardioides</i>	carrot wood	F
<i>Cupressus glabra</i>	Arizona cypress	F
<i>Cupressocyparis 'Emerald Isle'</i>	emerald Isle leyland cypress	C, F
<i>Cupressocyparis leylandii</i>	blue leyland cypress	C, F
<i>Cupressus ariz 'Blue pyramid'</i>	blue pyramid cypress	C, F
<i>Cupressus sempervirens</i>	cypress tree	C, F
<i>Cycas revoluta</i>	sago palm	C, F
<i>Elaeagnus angustifolia</i>	Russian olive	C, F
<i>Elaeagnus x ebbengei 'Gilt edge'</i>	gilt edge silverberry	C, F
<i>Eucalyptus camaldulensis</i>	red gum eucalyptus	F
<i>Eucalyptus cinerea</i>	eucalyptus, mealy	F
	silver dollar eucalyptus	F
<i>Eucalyptus microtheca</i>	coolibah tree	C, F
<i>Eucalyptus sideroxylon</i>	eucalyptus, red ironbark	F
<i>Fagus sylvatica</i>	European beech	C, F
<i>Ficus benjamina</i>	fig	C, F
	mini ficus	C, F
<i>Fraxinus udhei</i>	shamel ash	C, F
<i>Ginkgo biloba</i>	ginkgo (maidenhair tree)	F
<i>Gleditsia triacanthos</i>	honey locust	F
	shademaster honey locust	F
<i>Heteromeles arbutiflora</i>	toyon	F
<i>Illicium floridanum</i>	Florida anise-tree	C, F
<i>Juniperus virginiana</i>	redcedar, eastern	C, F
<i>Leptospermum scoparium</i>	tea tree	C, F
<i>Liquidambar styraciflua</i>	sweetgum, American	F
<i>Magnolia grandiflora</i>	magnolia, southern	C, F
<i>Magnolia stellata</i>	royal star magnolia	C, F
<i>Morus alba</i>	white mulberry	F
<i>Musa aluminata</i>	banana	C, F
<i>Oxydendrum arboreum</i>	sourwood	C, F
<i>Picea abies</i>	pendula-weeping Norway spruce	C, F

	repens-spreading Norway spruce	C, F
	spruce, Norway	C, F
<i>Picea glauca</i>	conica-dwarf alberta spruce	C, F
<i>Picea glauca conica</i>	conica-dwarf alberta spruce	F
<i>Picea pungens</i>	dwarf globe blue spruce	C, F
	glauca-Colorado blue spruce	C, F
	hoopsii-hoop's blue spruce	C, F
	koster-koster blue spruce	F
<i>Pinus aristata</i>	bristlecone pine	F
<i>Pinus canariensis</i>	canary island pine	F
<i>Pinus contorta</i>	shore pine, beach pine	F
<i>Pinus eldarica</i>	eldarica pine	C, F
<i>Pinus leucodermis</i>	Bosnian pine	C, F
<i>Pinus mugo</i>	pumilio-shrubby swiss mountain pine	C, F
<i>Pinus nigra</i>	Austrian black pine	C, F
<i>Pinus radiata</i>	monterey pine	F
<i>Pinus strobus</i>	eastern white pine	C, F
	white pine	C, F
<i>Pinus sylvestris</i>	columnar Scotch pine	C, F
	Scotch pine	C, F
<i>Pinus thunbergiana</i>	Japanese black pine	C, F
<i>Platanus occidentalis</i>	American sycamore	F
<i>Platanus racemosa</i>	Californina sycamore	F
<i>Podocarpus spp.</i>	podocarpus	F
<i>Populus deltoides</i>	cottonwood	F
<i>Prosopis chilensis</i>	Chilean mesquite	C, F
<i>Prunus yedoensis</i>	yoshino flowering cherry	F
<i>Prunus caroliniana</i>	Carolina laurel cherry	C, F
<i>Prunus laurocerasus</i>	English laurel	C, F
<i>Quercus ilicifolia</i>	bear oak	F
<i>Quercus palustris</i>	pin oak	F
<i>Quercus phellos</i>	willow oak	C, F
<i>Quercus rubra</i>	red oak	C, F
<i>Quercus virginiana</i>	live oak	C, F
<i>Salix babylonica</i>	babylon weeping willow	F
	corkscrew willow	F
<i>Sequoiadendron giganteum</i>	giant sequoia	F
<i>Swietenia mahogani</i>	mahogany	F
<i>Tabebuia caraiba</i>	yellow tab	F
<i>Trachycarpus fortunei</i>	windmill palm	C, F
<i>Tsuga canadensis</i>	eastern hemlock	C, F
<i>Ulmus parvifolia</i>	Chinese elm	F
<i>Washingtonia robusta</i>	Mexican fan palm	F
<i>Zamia furfuracea</i>	cardboard palm	C, F

## Shrubs

**Treatment Method:**  
**C = Container Grown**  
**F = Field Grown**

Scientific Name	Common Name	
<i>Abelia grandiflora</i>	edward goucher abelia	C, F
	glossy abelia	C, F
	sunrise variegated abelia	C, F
<i>Acacia abyssinica</i>	abyssinica acacia	C, F
<i>Acacia redolens</i>	acacia, prostrate	C, F
<i>Acacia stenophylla</i>	shoestring acacia	C, F

<i>Acalypha wilkesiana</i>	copper leaf	C, F
<i>Acer ginnala</i>	amur maple	C, F
<i>Acer palmatum</i>	coral bark Japanese maple	C, F
<i>Acer palmatum</i>	dwarf Japanese maple	C, F
<i>Agave americana</i>	century plant	F
<i>Amelanchier alnifolia</i>	Saskatoon serviceberry	C, F
<i>Andromeda polifolia</i>	bog rosemary	C, F
<i>Anisodonte hypomandarum</i>	cape mallow	C, F
<i>Arctostaphylos uva-ursi</i>	bearberry	C, F
<i>Ardisia japonica</i>	chirimen marlberry	C, F
<i>Athyrium nipponicum</i>	Japanese painted fern	C, F
<i>Baccharis pilularis</i>	coyotebush	F
<i>Berberis gladywensii</i>	william penn barberry	C, F
<i>Berberis mentorensis</i>	mentor barberry	C, F
<i>Berberis thunbergii</i>	aurea-golden Japanese barberry	F
	crimson pygmy barberry	C, F
	rose glow barberry	C, F
	atropurea-redleaf Japanese barberry	F
	cherry bomb	C, F
<i>Bougainvillea</i> spp.	barbara karst	C, F
	California gold	C, F
	pink pixie	C, F
	scarlet o'hara	C, F
	temple fire	C, F
	Texas dawn	C, F
<i>Buxus microphylla japonica</i>	boxwood, Japanese	F
<i>Buxus microphylla Koreana</i>	Korean boxwood	F
<i>Buxus sempervirens</i>	boxwood, common	C, F
<i>Buxus x 'Green velvet'</i>	green velvet boxwood	C, F
<i>Callistemon citrinus</i>	bottlebrush, lemon	F
<i>Callistemon viminalis</i>	weeping bottlebrush	C, F
<i>Calluna vulgaris</i>	spring torch Scotch heather	C, F
<i>Camellia japonica</i>	camellia	C, F
<i>Caryopteris clandonensis</i>	blue mist bluebeard	C, F
<i>Caryopteris x clandonen 'Blk night'</i>	dark knight bluebeard	C, F
<i>Cassia artemisioides</i>	cassia, feathery	C, F
<i>Cassia, eremophila</i>	senna	C, F
<i>Ceanothus</i> spp.	wild lilac	F
<i>Cephalotaxus drupacae</i>	plum yew	C, F
<i>Cerastium tomentosum</i>	snow-in-summer	C, F
<i>Ceratostigma willmottianum</i>	Chinese plumbago	C, F
<i>Ceratostigma plumbaginoides</i>	dwarf plumbago	C, F
<i>Chaenomeles japonica</i>	orange flowering quince	C, F
<i>Chamaecyparis obtusa</i>	kosteri cypress	C, F
	nana-dwarf hinoki cypress	C, F
	torulosa cypress	C, F
<i>Chamaecyparis pisifera</i>	baileyi-dogwood	F
	flaviramea-dogwood	F
	sawara-false cypress	F
	squarrosa minima cypress	C, F
	filifera-thread cypress	C, F
<i>Chrysalidocarpus lutescens</i>	areca palm	F
<i>Clethra alnifolia</i>	summersweet	C, F
<i>Cleyera japonica</i>	cleyera, Japanese	C, F
<i>Colonema pulchrum</i>	pink breath of heaven	C, F
<i>Convolvulus mauritanicus</i>	ground morningglory	C, F

<i>Convolvus cneorum</i>	bush morning glory	C, F
<i>Cornus alba</i>	sibirica-siberian dogwood	C, F
<i>Cornus stolonifera</i>	baileyi-red-osier dogwood	F
	flaviramea-yellowtwig dogwood	F
<i>Corylus contorta</i>	walking stick	C, F
<i>Cotinus coggygria</i>	royal purple smoke tree	C, F
<i>Cotinus dammeri</i>	coral beauty smoke tree	C, F
	eichholz smoke tree	C, F
<i>Cotoneaster adpressus</i>	praecox-early cotoneaster	C, F
<i>Cotoneaster apiculatus</i>	cotoneaster, cranberry	C, F
<i>Cotoneaster congestus</i>	cotoneaster, pyrenees	C, F
<i>Cotoneaster dammeri</i>	cotoneaster, bearberry	C, F
<i>Cotoneaster himalayan</i>	himalayan cotoneaster	C, F
<i>Cotoneaster horizontalis</i>	cotoneaster, rock	C, F
<i>Cycas revoluta</i>	sago palm	C, F
<i>Cytisus praecox</i>	hollandia-warminster broom	C, F
<i>Cytisus scoparius</i>	lena-Scotch broom	C, F
<i>Cytisus spp.</i>	hollandia-Scotch broom	F
	allgold warminster broom	C, F
	lilac time broom	C, F
<i>Dalea greggii</i>	trailing indigo bush	C, F
<i>Daphne cneorum</i>	rose daphne	C, F
<i>Daphne odora</i>	fragrant daphne	C, F
<i>Deutzia crenata</i>	nakiana-dwarf deutzia	C, F
<i>Deutzia gracilis</i>	slender gracilis	C, F
<i>Dodonea viscosa</i>	hopseed bush	F
<i>Elaeagnus pungens</i>	fruitland silver berry	C, F
<i>Enkianthus companulatus</i>	red-veined enkianthus	C, F
<i>Erica cinerea</i>	purple bell heather	C, F
<i>Erica vagans</i>	cornish heather	C, F
<i>Erica x darleyensa</i>	Mediterranean pink heather	C, F
<i>Escalonia spp.</i>	escallonia	C, F
<i>Eugenia myrtifolia</i>	teenie genie brushcherry	C, F
<i>Eugenia myrtifolia 'Globulus'</i>	dwarf brush cherry	C, F
<i>Euonymus fortunei</i>	canadale gold euonymus	C, F
	emerald'n gold euonymus	F
	sunspot euonymus	C, F
<i>Euonymus japonica</i>	silver king euonymus	F
	chollipo euonymus	C, F
	gold spot euonymus	C, F
	silver princess euonymus	C, F
	variegated evergreen euonymus	C, F
<i>Euonymus kiatschovica</i>	spreading euonymus	C, F
<i>Euonymus vegetus</i>	bigleaf wintercreeper	C, F
<i>Fatsia japonica</i>	Japanese aralia	C, F
<i>Felicia amelloides</i>	blue marguerite	C, F
<i>Forsythia intermedia</i>	forsythia, border	C, F
<i>Forsythia x 'Spring glory'</i>	spring glory forsythia	C, F
<i>Fuchsia x 'Santa Claus'</i>	santa claus fuchsia	C, F
<i>Gardenia jasminoides</i>	august beauty gardenia	C, F
	gardenia	C, F
	miniature gardenia	C, F
	radican gardenia	C, F
<i>Gaultheria procumbens</i>	wintergreen	C, F
<i>Gaultheria shallon</i>	salal/lemon leaf	C, F
<i>Gelsemium sempervirens</i>	Carolina jessamine	C, F

<i>Genista pilosa</i>	woadwaxen	C, F
<i>Hamamelis virginiana</i>	common witch hazel	C, F
<i>Hardenbergia violacea</i>	lilac vine	C, F
<i>Hebe buxifolia</i>	boxleaf hebe	C, F
<i>Hibiscus rosa-sinensis</i>	ross estey-hibiscus	C, F
<i>Hibiscus syriacus</i>	rose of sharon, red bird	C, F
	rose of sharon, red heart	F
	rose of sharon, woodbridge	C, F
<i>Ilex aquifolium</i>	balkans holly	F
	gold coast holly	F
<i>Ilex aquipernyi</i>	san jose holly	C, F
<i>Ilex attenuata</i>	savannah holly	C, F
<i>Ilex cornuta</i>	burford holly	C, F
	dwarf burford holly	C, F
	needle point holly	C, F
	holly, Chinese	
<i>Ilex crenata</i>	compacta-dwarf Japanese holly	C, F
	convexa holly	C, F
	dwarf Chinese holly	C, F
	green luster holly	C, F
	helleri-heller's Japanese holly	C, F
	hetzii's Japanese holly	C, F
	stokesii Japanese holly	C, F
<i>Ilex glabra</i>	compacta-compact inkberry holly	C, F
<i>Ilex glabra</i>	nordica-inkberry holly	C, F
<i>Ilex meserveae</i>	blue boy holly	C, F
	blue girl holly	C, F
	China boy holly	C, F
	China girl holly	C, F
	ebony magic holly	F
<i>Ilex vomitoria</i>	nana-dwarf yaupon holly	C, F
	pendula-weeping yaupon holly	C, F
	yaupon holly	C, F
<i>Illicium annisatum</i>	mystery gardenia	C, F
<i>Itea ilicifolia</i>	henry garnet holly leaf sweetspire	C, F
<i>Ixora collinea</i>	ixora	C, F
<i>Juniperus chinensis</i>	hollywood juniper	C, F
	media-old gold juniper	C, F
	pfizer juniper	C, F
	pfizerana glauca-blue juniper	C, F
	pfizerana-pfizer juniper	C, F
	sea green juniper	F
	torulosa-hollywood juniper	C, F
<i>Juniperus conferta</i>	emerald sea shore juniper	C, F
	shore juniper	C, F
<i>Juniperus horizontalis</i>	andorra juniper	C, F
	bar harbor juniper	C, F
	blue chip juniper	C, F
	blue rug juniper	C, F
	creeping juniper	C, F
	dwarf andorra juniper	C, F
	huntington blue juniper	C, F
	plumosa-andorra juniper	C, F
	wiltonii-blue carpet juniper	C, F
<i>Juniperus procumbens</i>	nana-dwarf Japaneses garden juniper	C, F
<i>Juniperus prostrata</i>	prostrata juniper	C, F



<i>Juniperus sabina</i>	broadmoor juniper	C, F
	foemina-hicks juniper	C, F
	savin juniper	C, F
	tamariscifolia-tam juniper	C, F
<i>Juniperus scopulorum</i>	emerald green juniper	F
	wichita blue juniper	C, F
<i>Juniperus squamata</i>	blue juniper	C, F
	blue star juniper	C, F
	parsonii juniper	C, F
<i>Kalmia latifolia</i>	laurel, mountain	C, F
<i>Lagerstroemia indica</i>	crape myrtle	C, F
<i>Lantana</i> spp.	lantana	C, F
<i>Leucophyllum frutescens</i>	Texas sage	C, F
<i>Leucophyllum laevigatum</i>	chihuahuan sage	C, F
<i>Leucothoe axillaris</i>	leucothoe, coast	C, F
<i>Leucothoe fontanesiana</i>	leucothoe, drooping	C, F
<i>Ligustrum japonicum</i>	privet, Japanese	C, F
	wax ligustrum	C, F
	yellow tip ligustrum	C, F
<i>Ligustrum lucidum</i>	privet, glossy	C, F
<i>Ligustrum ovalifolium</i>	California privet	F
<i>Ligustrum texanum</i>	howardi privet	C, F
	wax leaf privet	C, F
<i>Ligustrum vicaryi</i>	privet, golden	F
	vicary golden privet	F
<i>Ligustrum vulgare</i>	lodense privet	C, F
<i>Livistona chinensis</i>	Chinese fountain palm	F
<i>Lonicera fragrantissima</i>	winter honeysuckle	C, F
<i>Lonicera periclymenum</i>	flowering woodbine	C, F
	serotina woodbine	C, F
<i>Lonicera sempervirens</i>	trumpet honeysuckle	C, F
<i>Loropetalum chin rubrum</i> 'Razzleb'	razzleberri fringe flower	C, F
<i>Loropetalum chinense</i>	sizzling pink fringe flower	C, F
<i>Mahonia aquifolium compacta</i>	dwarf Oregon grape	C, F
<i>Mahonia bealei</i>	leather leaf mahonia	C, F
<i>Mahonia repens</i>	creeping mahonia	C, F
<i>Michelia figo</i>	banana shrub	C, F
<i>Myrica cerifera</i>	wax myrtle	C, F
<i>Myoporum parvifolium</i>	putah creek	C, F
<i>Nandina domestica</i>	compacta-dwarf heavenly bamboo	C, F
	harbour dwarf-heavenly bamboo	C, F
	heavenly bamboo (nandina)	C, F
	nana compacta-heavenly bamboo	C, F
	nana purpurea-heavenly bamboo	C, F
	woods dwarf-heavenly bamboo	C, F
<i>Nerium oleander</i>	hardy red oleander	C, F
	oleander	C, F
	ruby lace oleander	C, F
<i>Osmanthus fortunei</i>	fortunes osmanthus	C, F
<i>Osmathus fragrans</i>	sweet olive	C, F
<i>Pennisetum setaceum</i> 'Rubrum'	purple fountain grass	C, F
<i>Phoenix roebelenii</i>	pigmy date palm	C, F
<i>Photinia fraseri</i>	fraser's photinia	C, F
<i>Physocarpus opulifolius</i>	dwarf ninebark	C, F
<i>Pieris japonica</i>	lily-of-the-valley	C, F
	mountain fire lily-of-the-valley	C, F

	snowdrift lily-of-the-valley	C, F
	temple bells lily-of-the-valley	C, F
	valley rose lily-of-the-valley	C, F
	valley valentine lily-of-the-valley	C, F
<i>Pieris japonica x forestii</i>	forest flame lily-of-the-valley	C, F
<i>Pinus mugo</i>	mugo-mugho pine	C, F
<i>Pittosporum tobira</i>	green pittosporum	C, F
	wheeler's dwarf pittosporum	C, F
<i>Plumbago ariculata</i>	blue cape plumbago	F
<i>Plumbago capensis</i>	plumbago	C, F
<i>Podocarpus macrophyllus</i>	yewpine	C, F
<i>Polygala fruticosa</i>	sweet pea shrub	C, F
<i>Polystichum polyblepharum</i>	tassel fern	C, F
<i>Potentilla fragiformis</i>	cinquefoil	F
<i>Potentilla fruticosa</i>	cinquefoil	C, F
	floppy disc cinquefoil	C, F
	gold drop potentilla	F
	goldfinger potentilla	C, F
	red ace potentilla	C, F
	sunset potentilla	C, F
	tangerine potentilla	C, F
<i>Potentilla spp.</i>	cinquefoil	
<i>Potentilla verna</i>	spring cinquefoil	C, F
<i>Prunus gladiosa</i>	dwarf pink flowering almond	C, F
<i>Prunus laurocerasus 'Otto luykens'</i>	otto luykens English laurel	C, F
<i>Psidium cattleianum</i>	strawberry guava	C, F
<i>Pyracantha fortuneana</i>	lolendei monrovia pyracantha	C, F
	monon pyracantha	F
	red elf hybrid pyracantha	F
	rutgers hybrid pyracantha	C, F
	santa cruz pyracantha	C, F
	victory pyracantha	F
<i>Raphiolepis indica</i>	charisma-monruce raphiolepis	C, F
	enchantress-moness raphiolepis	F
	raphiolepis (India hawthorn)	C, F
	springtime-monme raphiolepis	F
<i>Raphiolepis ovata</i>	roundleaf raphiolepis	C, F
<i>Rhododendron calendulaceum</i>	cannon's double azalea	C, F
	flame azalea	F
	golden flare azalea	C, F
	klondike azalea	C, F
<i>Rhododendron campylocarpum</i>	butterfly rhododendron	F
<i>Rhododendron carolinianum x daurium</i>	PJM rhododendron	C, F
<i>Rhododendron catawbiense</i>	catawba album rhododendron	C, F
	catawba rhododendron	C, F
	lord roberts rhododendron	C, F
	rocket rhododendron	C, F
<i>Rhododendron caucasicum x ponticum</i>	cunningham white rhododendron	C, F
<i>Rhododendron exbury</i>	cannon's double azalea	C, F
	golden flare azalea	C, F
	klondike azalea	C, F
<i>Rhododendron forrestii repens</i>	gomer waterer rhododendron	C, F
<i>Rhododendron forrestii x griersonianum</i>	elizabeth rhododendron	C, F
<i>Rhododendron griffithianum</i>	jean marie rhododendron	C, F
<i>Rhododendron hybrid spp.</i>	America rhododendron	C, F
	English roseum rhododendron	F

	nova zembla rhododendron	C, F
	scintillation rhododendron	C, F
<i>Rhododendron impeditum</i>	rhododendron	C, F
<i>Rhododendron indica</i>	formosa azalea	C, F
	waucabusa azalea	C, F
<i>Rhododendron kaempferi</i>	blue danube azalea	C, F
<i>Rhododendron kerume</i>	coral bells azalea	C, F
	hino crimson azalea	C, F
	hino pink azalea	C, F
	mildred azalea	C, F
	snow azalea	C, F
<i>Rhododendron maximum</i>	rhodie max (rosebay)	C, F
<i>Rhododendron mucronulatum</i>	rhododendron	F
<i>Rhododendron obtusum</i>	hino-crimsom azalea	C, F
<i>Rhododendron ponticum</i>	chioniodes rhododendron	C, F
	daphnoides rhododendron	C, F
<i>Rhododendron racemosum</i>	dwarf scarlet wonder rhododendron	C, F
	tribly rhododendron	C, F
	unique rhododendron	C, F
	vulcan rhododendron	C, F
<i>Rhododendron sassthigiatim x carolinianum</i>	ramapo rhododendron	C, F
<i>Rhododendron satuski</i>	gumpo pink azalea	C, F
	higasa azalea	F
	reijn azalea	C, F
<i>Rhododendron spp. hybrids</i>	carror azalea	C, F
	fashion azalea	C, F
	gerard christina azalea	F
	girard roberta azalea	C, F
	golden flare exbury azalea	F
	helmut vogel azalea	F
	henry's red azalea	F
	hot shot azalea	C, F
	hume azalea	F
	inga azalea	F
	irene koster azalea	C, F
	president clay azalea	C, F
	tradition azalea	C, F
<i>Rhus lancea</i>	sumac, African	C, F
<i>Rhus typhina</i>	staghorn sumac	C, F
<i>Rosa x 'Flower carpet'</i>	red groundcover rose	C, F
<i>Rosa rugosa</i>	ramanas rose	C, F
<i>Rosmarinus officinalis</i>	rosemary	F
<i>Senecio cineraria</i>	dusty-miller	
<i>Skimmia japonica</i>	Japanese skimmia	C, F
<i>Skimmia revesiana</i>	reeve's skimmia	C, F
<i>Solanum rantonetii 'Royal purple'</i>	Paraguay nightshade	C, F
<i>Spiraea bumalda</i>	anthony waterer spiraea	C, F
<i>Spiraea cinerea</i>	first snow spiraea	C, F
<i>Spiraea japonica</i>	dolchia spiraea	C, F
	Japanese alpine spiraea	C, F
	magic carpet spiraea	C, F
	neon flash spiraea	C, F
	shirobana spiraea	C, F
<i>Spiraea vanhouttei</i>	bridal wreath	C, F
<i>Streptosolen jamesonii</i>	marmalade bush	C, F

<i>Syringa rothomagensis</i>	Chinese lilac	C, F
<i>Syringa vulgaris</i>	lilac, common	F
<i>Taxus cuspidata</i>	yew, Japanese	F
<i>Tecomaria capensis</i>	cape honeysuckle	C, F
<i>Ternstroemia gymnanthera</i>	ternstroemia, Japanese	C, F
<i>Teucrium fruticans</i>	bush germander	C, F
<i>Thevetia nerifolia</i>	yellow oleander	C, F
<i>Thuja occidentalis</i>	emerald arborvitae	C, F
	globosa-globe arborvitae	C, F
	little giant-dwarf arborvitae	C, F
	nigra-dark American arborvitae	C, F
	pyramidalis arborvitae	C, F
	rheingold arborvitae	C, F
	techny arborvitae	F
	woodwardii arborvitae	C, F
<i>Thuja orientalis</i>	aureus nana-dwarf golden arborvitae	C, F
	minima glauca-dwarf arborvitae	C, F
<i>Tibouchina urvilleana</i>	princes flower	C, F
<i>Veitchia merrilli</i>	Christmas palm	F
<i>Viburnum bodnantense</i>	pink dawn viburnum	C, F
<i>Viburnum carlesii</i>	Koreanspice viburnum	C, F
<i>Viburnum davidii</i>	david viburnum	C, F
<i>Viburnum japonicum</i>	viburnum	F
<i>Viburnum judd (V. x juddii)</i>	viburnum	C, F
<i>Viburnum lantana</i>	wayfaring tree	F
<i>Viburnum opulus sterile</i>	common snowball viburnum	F
<i>Viburnum plicatum tomentosum</i>	doublefile viburnum	C, F
<i>Viburnum setigerum</i>	tea viburnum	F
<i>Viburnum tinus compactum</i>	spring bouquet viburnum	F
<i>Viburnum trilobum</i>	cranberry bush	C, F
<i>Viburnum trilobum compactum</i>	dwarf cranberry bush	C, F
<i>Viburnum x pragense</i>	viburnum	C, F
<i>Weigela florida</i>	bristol ruby weigela	C, F
	java red weigela	C, F
	minuet weigela	C, F
<i>Xylosma congestum</i>	xylosma	F
<i>Yucca filamentosa</i>	yucca	C, F

---

### Terms and Conditions of Use

---

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

---

### Warranty Disclaimer

---

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

### Inherent Risks of Use

---

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to

label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risks shall be assumed by Buyer.

---

### **Limitation of Remedies**

---

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

EPA accepted 10/10/17

**[Sub Label B: Residential Uses]**

(Base label):

<b>ISOXABEN</b>	<b>GROUP</b>	<b>21</b>	<b>HERBICIDE</b>
-----------------	--------------	-----------	------------------

**Gallery<sup>®</sup> Turf Fertilizer (R)****HERBICIDE**

A turf fertilizer containing Gallery, a preemergence herbicide for control of certain broadleaf weeds and suppression of annual grasses in landscape ornamentals, container grown ornamentals, and home lawns.

**Intended for use on residential sites**

In the state of Florida, do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden.

[XX lbs] Covers up to XX,000 sq ft (XXX' x XX')  
and/or

This bag covers XX,000 sq ft

**[Optional Label Claims]**

**Stops broadleaf weeds before they start**

**Controls over 40 of the toughest broadleaf weeds, as listed.**

**Contains "Gallery<sup>®</sup>" for a picture perfect lawn.**

**May provide season long control of Clover, Dandelion, Henbit, Oxalis, & Spurge**

**Prevents the Growth of Broadleaf Weeds, including Dandelion, for up to 8 Months**

**Long-Lasting, season long control.**

**Broad-Spectrum Broadleaf Weed Prevented**

**Water in to Establish a Broadleaf Weed Prevention Zone**

**Controls [tough] weeds [such as XX ] [ XX ] [ XX ] [ XX ] [ XX ]**

**Apply [spring, summer or fall]**

**Use for a [beautiful] [weed-free] lawn**

**Easy to use**

**Extended time release feeding of nitrogen lasts into summer**

**Effective weed control leads to greener, thicker lawns**

**All the fertilizer nutrients your lawns needs for spring**

**Feeds [greens] [strengthens] [thickens] your lawn for up to 3 months**

**Apply once for the whole season**

**Feeds [and strengthens] root systems without burning your lawn**

**Fertilize once for the whole season – Feeds [and greens] with slow release fertilizer for up to 3 months**

**Fertilizers to fill in bare spots [and provide plush green lawns]**

**Prevents weeds [and fertilizes the lawn]**

**Fertilizes to provide rich color [and vigorous growth]**

**Effective weed control – prevents weeds all season long**

**Season-long weed killer [and fertilizer]**

**Controls weeds longer**

**Fertilizes longer**

**For use on cool and warm season turfgrass**

**[Just] One application kills listed weeds**

**Better weed control than other brands**

**Contains the latest technology in weed control**

**University Tested & Researched.**

**Labeled for use around a wide variety of trees & shrubs.**

**For Use on all turfgrass types**

**Prevents weeds from germination.**

**Treat borders of lawns near streets, sidewalks and driveways where weeds appear most/heaviest.**

**Effective weed control/prevention assures thicker healthier lawns.**

**Apply in the fall for winter annual weeds.**

**Apply early spring or late fall**

**Growing a Better World**

**[Contains] Harrell's PROFertilizer**

**Harrell's Logo****SurfCote™ - Sustained Uniform Release Fertilizer Continuously Provides Nutrients****Your Prescription for a Picture Perfect Lawn****Turf Care Supply Corp – Manufacturing****Turf Care Supply Corp - TCS GrowStar Premier Turf Products****Premier Turf Products**

**Manufactured in the USA [Note: Only can use if all inerts used in formulation are produced in the US and the formulation was manufactured in USA]**

**If this product is being sold in a state requiring the publication of information regarding its metal content, information regarding the contents and levels of metals in this product is available at the following website: <http://www.aapfco.org/metals.html>**

**Information regarding the contents and levels of metals in this product is available on the Internet at:**

- **California Regulatory Site: <https://apps1.cdffa.ca.gov/FertilizerProducts/>**
- **Oregon Regulatory Site: [http://oda.state.or.us/dbs/heavy\\_metal/search.lasso](http://oda.state.or.us/dbs/heavy_metal/search.lasso)**
- **Washington Regulatory Site: <http://agr.wa.gov/> or <http://agr.wa.gov/PestFert/Fertilizers/ProductDatabase.aspx>**
- **<http://www.aapfco.org/metals.html>**
- **<http://www.regulatory-info-lpi.com>**

**It is a violation of law in Maryland and New Jersey to apply this product to turf at a rate that results in greater than 0.7 lb per 1,000 sq.ft. of readily available nitrogen or more than 0.9 lb per 1,000 sq.ft. of total nitrogen with a least 20% of which shall consist of slow release nitrogen per application. Additional restrictions apply to total nitrogen applied to turf per season. Other state and local restrictions may apply. [Maryland or New Jersey Applications]**

**No more than 0.7 lb per 1,000 sq.ft. of readily available nitrogen during any given 30 day period. No more than 0.9 lb (cool season grasses) or 1.0 lb (warm season grasses) per 1,000 sq.ft. of total nitrogen with a least 20% of which shall consist of slow release nitrogen during any given 30 day period. [Virginia Applications]**

**Do not apply near water storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back onto your lawn and garden. [Maryland Applications]**

**This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under §8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; 2) establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) reestablishing or repairing a turf area. [Maryland Applications which contain Phosphorus] Not for use on established lawns or grass in MD. [If phos >5% and is not classified as a “starter” fertilizer]**



In Virginia and New Jersey this lawn/turf fertilizer contains phosphorous and is only for nonagricultural use on (i) turf during its first growing season, (ii) on turf areas being repaired or renovated, and (iii) on turf where a soil test performed within the last 3 years indicates a phosphorous deficiency. This fertilizer is not for the routine maintenance of turf. [Virginia and New Jersey Applications which contain Phosphorus]

Apply only to actively growing turf. Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands in the driveway, sidewalk, or street, back onto your lawn. Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use. [FL Language for Specialty Products]

The maximum rate of application of this product must not exceed the lesser of the applicable rate determined from the [Table [XX]: Use Rates or Broadcast Application Rates], and the applicable rate determined in the State Nutrient Restrictions table.

**STATE NUTRIENT RESTRICTIONS**

State	Readily Available Nitrogen* lbs/1,000 sq ft	Total Nitrogen lbs/1,000 sq ft
MD, NH, VA	0.7	0.9
NJ	0.7	1.0
XXX	XXX	XXX

\* Nitrogen present beyond the amount shown as Readily Available must be in the form of Slow-Release nitrogen

Not for sale, distribution or use in New York State

<b>Group</b>	<b>21</b>	<b>HERBICIDE</b>
--------------	-----------	------------------

Active Ingredient:

isoxaben: N-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers.....0.38%

Other Ingredients .....99.62%

Total .....100.00%

Contains 0.09 lb active ingredient per 25 lb bag.

**Guaranteed Fertilizer Analysis: XX-XX-XX †**

Total Nitrogen (N)..... XX%

XX% Ammoniacal Nitrogen

XX% Nitrate Nitrogen

XX% Water-Insoluble Nitrogen\*

XX% Urea Nitrogen\*\*

XX% Slowly available water soluble Nitrogen.

XX% Other Water Soluble Nitrogen

Available Phosphate (P<sub>2</sub>O<sub>5</sub>)\*\*\* ..... XX%

Soluble Potash (K<sub>2</sub>O)..... XX%

Calcium (Ca) ..... XX%

Magnesium (Mg) ..... XX%

XX% Water Soluble Magnesium (Mg)  
 Sulfur\*\*\*\*  
 XX% Combined Sulfur (S)  
 Boron (B) ..... XX%  
 Chlorine (Cl) [Max, Not more than] ..... XX%  
 Cobalt (Co)..... XX%  
 Copper (Cu)..... XX%  
 Iron (Fe)..... XX%  
 XX% Chelated Iron (Fe)  
 XX% Water Soluble Iron (Fe) Manganese (Mn)XX%  
 Water Soluble Manganese (Mn) ..... XX%  
 Molybdenum (Mo) ..... XX%  
 Nickel (Ni)..... XX%  
 Sodium (Na)..... XX%  
 Zinc (Zn)..... XX%  
 XX% Chelated Zinc (Zn)

\*[XX]% Water insoluble nitrogen from biosolids.

\*\*[XX]% coated slow release Nitrogen (N)

\*\*[XX]% Slowly available [urea] nitrogen from [polycoated] [polymer/sulfur] coated urea

\*\*[XX]% Slowly available water soluble nitrogen

\*\*[XX]% Slowly available nitrogen from methylene ureas

\*\*[XX]% Slowly available nitrogen from ureaform

\*\*[XX]% Slowly available nitrogen from BioSolids

\*\*[XX]% Stabilized urea nitrogen or [urea] nitrogen stabilized with dicyandiamide

\*\*\*\*[XX]% Free Sulfur (S)

Derived From: [XX]<sup>†</sup>

Primary plant foods derived from: [XX]<sup>†</sup>

Secondary plant foods derived from [XX]<sup>†</sup>

Iron derived from: [XX]<sup>†</sup>

Density – XX lb./cu.ft.

**[Editor's Note: the following information may be added in the boxes above, this footnote must be deleted from the label after completion of the above Fertilizer Analysis data]**

<sup>†</sup> Urea, methylene urea, sulfur coated urea, polymer coated urea, poly[mer] coated sulfur coated urea, urea formaldehyde, Ureaform, ammonium sulfate, corn gluten, bio-solids (Milorganite, Nutri-Pel, composted poultry manure), activated sewage sludge, diammonium phosphate, monoammonium phosphate, magnesium ammonium phosphate hexa-hydrate, triple super phosphate, ammoniated phosphates, sulfate of potash, muriate of potash, sulfate of potash-magnesia, potassium nitrate, copper oxide, copper sulfate, iron/ferric oxide, iron/ferrous succrate, iron/ferrous oxy-sulfate, iron/ferrous sulfate, manganese oxide, molybdc oxide, manganese sulfate, manganese succrate, zinc oxide, zinc sulfate, sodium borate, boron, sulfur, sodium molybdate, stabilized nitrogen, potassium magnesium sulfate, ferrous sulfate monohydrate, magnesium sulfate or limestone.

**Keep Out Of Reach Of Children**

**CAUTION**

---

**Precautionary Statements**

**Hazards to Humans and Domestic Animals**

---

**Causes Eye Irritation • Harmful If Swallowed, Inhaled Or Absorbed Through The Skin**

**Avoid breathing dust and contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.**

### **First Aid**

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 [or 1-800-XXX-XXXX] for emergency medical treatment information.

### **Environmental Hazards**

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

**Ground Water Advisory:** This pesticide has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

**Surface Water Advisory:** This pesticide may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soil and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of isoxaben from runoff water and sediment.

### **Storage and Disposal**

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

**If empty:** Place in trash or offer for recycling if available.

**If partly filled:** Call your local solid waste agency or 1-800-992-5994 for disposal instructions. Never place unused product down any indoor or outdoor drain.

**Refer to label booklet for Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994 or [1-800-XXX-XXXX].

EPA Reg. No. 62719-178

EPA Est. No. \_\_\_\_\_

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

**Produced by/for:**  
**Manufactured by/for:**  
**Distributed by/for:**  
**Guaranteed by:**

**Dow AgroSciences LLC**  
**9330 Zionsville Road**  
**Indianapolis, IN 46268**

**NET WEIGHT** \_\_\_\_\_

(cover/shipping container):

<b>ISOXABEN</b>	<b>GROUP</b>	<b>21</b>	<b>HERBICIDE</b>
-----------------	--------------	-----------	------------------

# Gallery® Turf Fertilizer (R)

## HERBICIDE

A turf fertilizer containing Gallery, a preemergence herbicide for control of certain broadleaf weeds and suppression of annual grasses in landscape ornamentals, container grown ornamentals, and home lawns.

### Intended for use in residential sites

In the state of Florida, do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden.

[XX lbs] Covers up to XX,000 sq ft (XXX' x XX')  
and/or

This bag covers XX,000 sq ft

Not for sale, distribution or use in New York State

<b>Group</b>	<b>21</b>	<b>HERBICIDE</b>
--------------	-----------	------------------

Active Ingredient:

isoxaben: N-[3-(1-ethyl-1-methylpropyl)-  
5-isoxazolyl]- 2,6-dimethoxybenzamide  
and isomers .....0.38%

Other Ingredients .....99.62%

Total .....100.00%

Contains 0.09 lb active ingredient per 25 lb bag.

### Guaranteed Fertilizer Analysis: XX-XX-XX †

Total Nitrogen (N)..... XX%

XX% Ammoniacal Nitrogen

XX% Nitrate Nitrogen

XX% Water-Insoluble Nitrogen\*

XX% Urea Nitrogen\*\*

XX% Slowly available water soluble Nitrogen.

XX% Other Water Soluble Nitrogen

Available Phosphate (P<sub>2</sub>O<sub>5</sub>)\*\*\* ..... XX%

Soluble Potash (K<sub>2</sub>O)..... XX%

Calcium (Ca) ..... XX%

Magnesium (Mg) ..... XX%

XX% Water Soluble Magnesium (Mg)

Sulfur\*\*\*\*

XX% Combined Sulfur (S)

Boron (B) ..... XX%

Chlorine (Cl) [Max, Not more than] ..... XX%

Cobalt (Co)..... XX%  
 Copper (Cu)..... XX%  
 Iron (Fe)..... XX%  
     XX% Chelated Iron (Fe)  
     XX% Water Soluble Iron (Fe) Manganese (Mn)XX%  
     Water Soluble Manganese (Mn) ..... XX%  
 Molybdenum (Mo) ..... XX%  
 Nickel (Ni)..... XX%  
 Sodium (Na)..... XX%  
 Zinc (Zn)..... XX%  
     XX% Chelated Zinc (Zn)

\*[XX]% Water insoluble nitrogen from biosolids.  
 \*\*[XX]% coated slow release Nitrogen (N)  
 \*\*[XX]% Slowly available [urea] nitrogen from [polycoated] [polymer/sulfur] coated urea  
 \*\*[XX]% Slowly available water soluble nitrogen  
 \*\*[XX]% Slowly available nitrogen from methylene ureas  
 \*\*[XX]% Slowly available nitrogen from ureaform  
 \*\*[XX]% Slowly available nitrogen from BioSolids  
 \*\*[XX]% Stabilized urea nitrogen or [urea] nitrogen stabilized with dicyandiamide  
 \*\*\*\*[XX]% Free Sulfur (S)

Derived From: [XX]†  
 Primary plant foods derived from: [XX] †  
 Secondary plant foods derived from [XX] †  
 Iron derived from: [XX] †

Density – XX lb./cu.ft.

**[Editor’s Note: the following information may be added in the boxes above, this footnote must be deleted from the label after completion of the above Fertilizer Analysis data]**

† Urea, methylene urea, sulfur coated urea, polymer coated urea, poly[mer] coated sulfur coated urea, urea formaldehyde, Ureaform, ammonium sulfate, corn gluten, bio-solids (Milorganite, Nutri-Pel, composted poultry manure), activated sewage sludge, diammonium phosphate, monoammonium phosphate, magnesium ammonium phosphate hexa-hydrate, triple super phosphate, ammoniated phosphates, sulfate of potash, muriate of potash, sulfate of potash-magnesia, potassium nitrate, copper oxide, copper sulfate, iron/ferric oxide, iron/ferrous sucrate, iron/ferrous oxy-sulfate, iron/ferrous sulfate, manganese oxide, molybdc oxide, manganese sulfate, manganese sucrate, zinc oxide, zinc sulfate, sodium borate, boron, sulfur, sodium molbdate, stabilized nitrogen, potassium magnesium sulfate, ferrous sulfate monohydrate, magnesium sulfate or limestone.

**Keep Out Of Reach Of Children**  
**CAUTION**

**Refer to inside of label booklet for additional precautionary information including Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994 [or 1-800-XXX-XXXX].

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

**Produced by/for:**  
**Manufactured by/for:**  
**Distributed by/for:**  
**Guaranteed by:**

**Dow AgroSciences LLC**  
**9330 Zionsville Road**  
**Indianapolis, IN 46268**

**NET WEIGHT \_\_\_\_\_**

(Page 1 through end):

**If this product is being sold in a state requiring the publication of information regarding its metal content, information regarding the contents and levels of metals in this product is available at the following website: <http://www.aapfco.org/metals.html>**

**Information regarding the contents and levels of metals in this product is available on the Internet at:**

- **California Regulatory Site: <https://apps1.cdfa.ca.gov/FertilizerProducts/>**
- **Oregon Regulatory Site: [http://oda.state.or.us/dbs/heavy\\_metal/search.lasso](http://oda.state.or.us/dbs/heavy_metal/search.lasso)**
- **Washington Regulatory Site: <http://agr.wa.gov/> or <http://agr.wa.gov/PestFert/Fertilizers/ProductDatabase.aspx>**
- **<http://www.aapfco.org/metals.html>**
- **<http://www.regulatory-info-lpi.com>**

**It is a violation of law in Maryland and New Jersey to apply this product to turf at a rate that results in greater than 0.7 lb per 1,000 sq.ft. of readily available nitrogen or more than 0.9 lb per 1,000 sq.ft. of total nitrogen with a least 20% of which shall consist of slow release nitrogen per application. Additional restrictions apply to total nitrogen applied to turf per season. Other state and local restrictions may apply. [Maryland or New Jersey Applications]**

**No more than 0.7 lb per 1,000 sq.ft. of readily available nitrogen during any given 30 day period. No more than 0.9 lb (cool season grasses) or 1.0 lb (warm season grasses) per 1,000 sq.ft. of total nitrogen with a least 20% of which shall consist of slow release nitrogen during any given 30 day period. [Virginia Applications]**

**Do not apply near water storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back onto your lawn and garden. [Maryland Applications]**

**This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under §8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; 2) establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) reestablishing or repairing a turf area. [Maryland Applications which contain Phosphorus] Not for use on established lawns or grass in MD. [If phos >5% and is not classified as a "starter" fertilizer]**

**In Virginia and New Jersey this lawn/turf fertilizer contains phosphorous and is only for nonagricultural use on (i) turf during its first growing season, (ii) on turf areas being repaired or renovated, and (iii) on turf where a soil test performed within the last 3 years indicates a phosphorous deficiency. This fertilizer is not for the routine maintenance of turf. [Virginia and New Jersey Applications which contain Phosphorus]**

**Note: Supplemental applications of fertilizer are needed to achieve a minimum 1lb Nitrogen per 1,000 sq ft.**

**Apply only to actively growing turf. Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands in the driveway, sidewalk, or street, back onto your lawn. Check with your local Cooperative Extension Agency to obtain specific information on local turf best management**



practices. Check with your county or city government to determine if there are local regulations for fertilizer use. [FL Language for Specialty Products]

The maximum rate of application of this product must not exceed the lesser of the applicable rate determined from the [Table [XX]: Use Rates or Broadcast Application Rates], and the applicable rate determined in the State Nutrient Restrictions table.

#### STATE NUTRIENT RESTRICTIONS

State	Readily Available Nitrogen* lbs/1,000 sq ft	Total Nitrogen lbs/1,000 sq ft
MD, NH, VA	0.7	0.9
NJ	0.7	1.0
XXX	XXX	XXX

\* Nitrogen present beyond the amount shown as Readily Available must be in the form of Slow-Release nitrogen

---

### Precautionary Statements

---

#### Hazards to Humans and Domestic Animals

**Causes Eye Irritation • Harmful If Swallowed, Inhaled Or Absorbed Through The Skin**

**Avoid breathing dust and contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.**

#### First Aid

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 [or 1-800-XXX-XXXX] for emergency medical treatment information.

#### Environmental Hazards

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

**Ground Water Advisory:** This pesticide has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

**Surface Water Advisory:** This pesticide may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soil and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of isoxaben from runoff water and sediment.

---

## Directions for Use

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all directions carefully before applying this product.

### Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

**If empty:** Place in trash or offer for recycling if available.

**If partly filled:** Call your local solid waste agency or 1-800-992-5994 for disposal instructions. Never place unused product down any indoor or outdoor drain.

## Product Information

This turf fertilizer product contains Gallery<sup>®</sup>, a preemergence herbicide for control of certain broadleaf weeds and suppression of annual grasses in established turfgrass, landscape ornamentals, and container grown ornamentals. Apply in late summer to early fall, in early spring, or anytime prior to germination of target weeds.

## Precautions

Gallery Turf Fertilizer (R) controls weeds germinating from seed. Gallery Turf Fertilizer (R) does not control established weeds. Existing weeds should be controlled with postemergence herbicides. To ensure optimal fertilizer and herbicidal response, water treated areas as soon as possible after application. A single rainfall or sprinkler irrigation of 0.5 inch or more is required to activate this product.

## Use Restrictions

Do not apply more than 19 lb of Gallery Turf Fertilizer (R) per 1000 sq ft.

Not for sale, distribution or use in New York State.

Do not apply by air.

Do not apply Gallery Turf Fertilizer to turfgrass grown for seed.

## Treatment of Turfgrass or Ornamental Species not Listed on the Label for Gallery Turf Fertilizer (R)

Users who wish to use Gallery Turf Fertilizer (R) on a plant species not listed on this label must determine the suitability for such use by treating a small area or a small number of plants at a specified rate. Prior to

treating a larger area, observe the treated area/plants for any sign of herbicidal injury during 30 to 60 days of typical growing conditions.

## Application Instructions

Apply Gallery Turf Fertilizer (R) using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or plant injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. When using a drop-type spreader, a splash board mounted below the release point will provide more even granule distribution.

**[Editor's note: With available data Supplemental Distributor may add the following statement.]**  
**These settings were calibrated and field tested. However, age, condition of spreader speed of operations and evenness of terrain may require slightly different settings for desired coverage.**

Check equipment frequently to verify calibration and proper functioning.

**[Editor's note: Supplemental Distributor may add a table listing recommended application equipment and calibration settings.]**

Brand and Model of Spreader	Application Rates	Calibration Settings at Specified Rate					
	lb/1000 sq ft	4.6	XX	XX	XX	XX	5.7
	lb/acre	200	XX	XX	XX	XX	250
	lb nitrogen / 1000 sq ft	XX	XX	XX	XX	XX	XX
XXXXXXX		XX	XX	XX	XX	XX	XX

[AND/OR]

Calibration Settings			
Rate	lbs. Product per 1,000 sq ft	lbs. Nitrogen per 1,000 sq ft	lbs A.I per acre
Low	4.6	XX	XX
Medium	XX	XX	XX
High	5.7	XX	XX

[AND/OR]

Brand and Model of Spreader	Calibration Settings (lb/1000 sq ft)*		
	Low	Medium	High
XXXXXXX	XX	XX	XX
XXXXXXX	XX	XX	XX
XXXXXXX	XX	XX	XX

[\*These settings are approximate for the application rate shown per 1,000 square feet. You may need to adjust the spreader setting depending on walking speed, spreader condition and product]

## Broadcast Application Rates

Apply Gallery Turf Fertilizer (R) at a broadcast rate of 4.6 to 5.7 lb per 1000 sq ft. A minimum rate of 4.6 lb per 1000 sq ft will provide optimum results. Use the higher rate in the rate range where heavy weed infestation is anticipated.

## Weeds Controlled or Suppressed

### Weeds controlled when applied at 4.6 to 5.7 lb per 1000 sq ft

burweed, lawn	henbit	speedwell, thymeleaf
buttercup	knotweed, prostrate	spurge, prostrate
celery, wild	mallow, little	spurge, spotted
chickweed, common	parsley - piert	spurry, corn
chickweed, mouseear	pennywort, lawn	woodsorrel, creeping
foolsparsely	pineappleweed	woodsorrel, yellow
geranium, Carolina		

### Weeds partially controlled by a single application or controlled by repeat applications when applied at 4.6 to 5.7 lb per 1000 sq ft

clover, spp. <sup>1</sup>	plantain, broadleaf
clover, white <sup>1</sup>	plantain, buckhorn
dandelion <sup>1</sup>	plantago, slender
medic, black	

<sup>1</sup>From seed only

### Weeds suppressed or partially controlled when applied at 4.6 to 5.7 lb per 1000 sq ft

bluegrass, annual  
crabgrass  
ryegrass, annual (Italian)

### **Weed Resistance Management:**

~~Isoxaben, the active ingredient in this product, is a Group 21 herbicide based on the mode of action classification system of the Weed Science Society of America. Any weed population may contain or develop plants resistant to Group 21 herbicides. Resistant weeds may dominate the weed population if these herbicides are used repeatedly in the same field. Such resistant weed plants may not be effectively managed using Group 21 herbicides but may be effectively managed utilizing other herbicides alone or in mixtures from a different herbicide Groups that are labeled for control of these weeds and/or by using cultural or mechanical practices. However, an herbicide mode of action classification by itself may not adequately address specific weeds that are resistant to specific herbicides. Consult your local company representative, state cooperative extension service, professional consultants or other qualified authorities to determine appropriate actions for treating specific resistant weeds or to report herbicide failures.~~

### **Best Management Practices:**

~~Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is recommended. A diversified weed management program may include the use of multiple herbicides with different modes of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistant weeds. It is recommended to scout for weeds before Gallery Turf Fertilizer application for identification and growth stage and after application to facilitate the early identification of weed shifts and/or weed resistance and thus provide direction on future weed management practices. One of the best ways to contain resistant weed populations is to implement measures to avoid allowing weeds to reproduce by seed or to proliferate~~

~~vegetatively. Cleaning equipment between sites and avoiding movement of plant material between sites will greatly aid in reducing the spread of resistant weed seed.~~

## Uses

### Turfgrass

Gallery Turf Fertilizer (R) may be used as a preemergence treatment for control of certain broadleaf weeds in established cool season and warm season turfgrass.

Apply Gallery Turf Fertilizer (R) anytime prior to germination of target weeds.

**Note:** Refer to the Product Information section of this label for general use precautions and information on application rates and weeds controlled prior to using this product.

#### Use Restrictions:

Apply Gallery Turf Fertilizer (R) only on established turfgrass. Apply to newly seeded turfgrass (including overseeded turfgrass) **only** after seedlings are established (three leaf stage and side shoots are visible) and well rooted. Do not overseed established turfgrass sooner than 60 days following an application of Gallery Turf Fertilizer (R).

- Do not apply Gallery Turf Fertilizer (R) to dichondra.
- Florida Applications: Do not exceed [2] [3] applications per year on bahiagrass in the north.
- Florida Applications: Do not exceed [2] [3] [4] applications per year on bahiagrass in [central Florida] [the south].
- Florida Applications: Do not exceed [3] [4] [5] applications per year on bermudagrass in the north.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on bermudagrass in central Florida.
- Florida Applications: Do not exceed [5] [6] [7] applications per year on bermudagrass in the south.
- Florida Applications: Do not exceed [1] [2] applications per year on centipedegrass in the north.
- Florida Applications: Do not exceed [2] [3] applications per year on centipedegrass in [central Florida] [the south].
- Florida Applications: Do not exceed [2] [3] [4] applications per year on St. Augustinegrass in the north.
- Florida Applications: Do not exceed [2] [3] [4] [5] applications per year on St. Augustinegrass in central Florida.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on St. Augustinegrass in the south.
- Florida Applications: Do not exceed [3] [4] [5] applications per year on zoysiagrass in the north.
- Florida Applications: Do not exceed [3] [4] [5] [6] applications per year on zoysiagrass in central Florida.
- Florida Applications: Do not exceed [4] [5] [6] applications per year on zoysiagrass in the south.

**Gallery Turf Fertilizer (R) may be used for use on the following turfgrass species:**

#### Established Cool Season Turfgrass

bentgrass, colonial	fescue, creeping red
bentgrass, creeping	fescue, sheep
bluegrass, Kentucky	fescue, tall
fescue, chewings	ryegrass, perennial

#### Established Warm Season Turfgrass

bahiagrass	fescue, tall
Bermudagrass	St. Augustinegrass
centipedegrass	zoysiagrass

### Ornamental Plantings

Gallery Turf Fertilizer (R) may be used as a preemergence treatment for control of certain broadleaf weeds in landscape and container grown ornamentals.

Apply Gallery Turf Fertilizer (R) anytime prior to germination of target weeds or immediately after preparing the soil.

**Note:** Refer to the Product Information section of this label for general use precautions and information on application rates and weeds controlled prior to using this product.

#### **Use Restrictions:**

To avoid possible plant injury, **do not** apply Gallery Turf Fertilizer (R) to:

- unrooted liners or cuttings that have been planted in pots for the first time
- pots less than six inches wide
- bedding plants or areas where bedding plants will be planted or transplanted within 1 year after application

**Do not** apply Gallery Turf Fertilizer (R) to newly transplanted ornamentals until soil or potting media has been settled by packing and irrigation or rainfall and no cracks are present or plant injury may occur.

Applications of Gallery Turf Fertilizer (R) over the top of plants with newly forming buds may cause injury. Possible plant injury may be avoided by applying to the soil surface beneath ornamental plants.

When planting into a site treated with Gallery Turf Fertilizer (R) in the previous 8 months, use untreated soil as fill around roots when replacing plants or injury may occur.

Injury to certain ornamental plants has been observed following application of Gallery Turf Fertilizer (R). To avoid plant injury, do not use Gallery Turf Fertilizer (R) for weed control in the following ornamental plant species:

bugle	mustard
cajeput tree	prince of wales juniper
candytuft	purple coneflower
Carolina rhododendron	roseum elegans rhododendron
dwarf burning bush	spurge
green yucca	stonecrop
hydrangea	

**Gallery Turf Fertilizer (R) may be used on the following established plant species:**

#### **Trees**

albus-flowering maple	Japanese black pine
Australian tree fern	king palm
Austrian black pine	live oak
banana	luteus-flowering maple
birch, river	magnolia, southern
blue leyland cypress	Mediterranean fan palm
blue pyramid cypress	mini ficus
bosnian pine	norfolk island pine
bottle tree	palm
cardboard palm	parlor palm
Carolina laurel cherry	pendula-weeping Norway spruce
Chilean mesquite	pumilio-shrubby swiss mountain pine
cloud nine dogwood	queen palm
columnar Scotch pine	red baubinia
conica-dwarf alberta spruce	red oak

coolibah tree  
 cryptomeria, Japanese  
 cypress tree  
 dogwood, flowering  
 dwarf globe blue spruce  
 eastern hemlock  
 eastern white pine  
 eldarica pine  
 emerald Isle leyland cypress  
 English laurel  
 European beech  
 ficus  
 fir, balsam  
 Florida anise-tree  
 gilt edge silverberry  
 glauca-Colorado blue spruce  
 hoopsii-hoop's blue spruce

redbud  
 redcedar, eastern  
 repens-spreading Norway spruce  
 roseus-flowering maple  
 royal star magnolia  
 Russian olive  
 sago palm  
 Scotch pine  
 shamel ash  
 silver maple  
 sourwood  
 spruce, Norway  
 tangerine-flowering maple  
 tea tree  
 white pine  
 willow oak  
 windmill palm

## Shrubs

abyssinica acacia  
 acacia, prostrate  
 allgold warminster broom  
 America rhododendron  
 amur maple  
 andorra juniper  
 anthony waterer spiraea  
 august beauty gardenia  
 aureus nana-dwarf golden arborvitae  
 banana shrub  
 bar harbor juniper  
 barbara karst  
 bearberry  
 bigleaf wintercreeper  
 blue boy holly  
 blue chip juniper  
 blue danube azalea  
 blue girl holly  
 blue juniper  
 blue marguerite  
 blue mist bluebeard  
 blue rug juniper  
 blue star juniper  
 bog rosemary  
 boxleaf hebe  
 boxwood, common  
 bridal wreath  
 bristol ruby weigela  
 broadmoor juniper  
 burford holly  
 bush germander  
 bush morning glory  
 California gold  
 camellia  
 canadale gold euonymus  
 cannon's double azalea  
 cape honeysuckle

formosa azalea  
 forsythia, border  
 fortunes osmanthus  
 fragrant daphne  
 fraser's photinia  
 fruitland silver berry  
 gardenia  
 girard roberta azalea  
 globosa-globe arborvitae  
 glossy abelia  
 gold spot euonymus  
 golden flare azalea  
 goldfinger potentilla  
 gomer waterer rhododendron  
 green luster holly  
 green pittosporum  
 green velvet boxwood  
 ground morningglory  
 gumpo pink azalea  
 harbour dwarf-heavenly bamboo  
 hardy red oleander  
 heavenly bamboo (nandina)  
 helleri-heller's Japanese holly  
 henry garnet holly leaf sweetspire  
 hetzii's Japanese holly  
 himalayan cotoneaster  
 hino crimson azalea  
 hino pink azalea  
 hollandia-warminster broom  
 hollywood juniper  
 hot shot azalea  
 howardi privet  
 huntington blue juniper  
 irene koster azalea  
 ixora  
 Japanese alpine spiraea  
 Japanese aralia

purple fountain grass  
 putah creek  
 pyramidalis arborvitae  
 radican gardenia  
 ramanas rose  
 ramapo rhododendron  
 razzleberri fringe flower  
 red ace potentilla  
 red groundcover rose  
 red-veined enkianthus  
 reeve's skimmia  
 reijn azalea  
 raphiolepis (India hawthorn)  
 rheingold arborvitae  
 rhodie max (rosebay)  
 rhododendron  
 rocket rhododendron  
 rose daphne  
 rose glow barberry  
 rose of sharon, red bird  
 rose of sharon, woodbridge  
 ross estey-hibiscus  
 roundleaf raphiolepis  
 royal purple smoke tree  
 ruby lace oleander  
 rutgers hybrid pyracantha  
 sago palm  
 salal/lemon leaf  
 san jose holly  
 santa claus fuchsia  
 santa cruz pyracantha  
 Saskatoon serviceberry  
 savannah holly  
 savin juniper  
 scarlet o'hara  
 scintillation rhododendron  
 senna

cape mallow	Japanese painted fern	serotina woodbine
Carolina jessamine	Japanese skimmia	shirobana spiraea
carror azalea	java red weigela	shoestring acacia
cassia, feathery	jean marie rhododendron	shore juniper
catawba album rhododendron	klondike azalea	sibirica-siberian dogwood
catawba rhododendron	Koreanspice vibunum	silver princess euonymus
charisma-monruce raphiolepis	kosteri cypress	sizzling pink fringe flower
cherry bomb	lantana	slender gracilis
chihuahuan sage	laurel, mountain	snow azalea
China boy holly	leather leaf mahonia	snowdrift lily-of-the-valley
China girl holly	lena-Scotch broom	snow-in-summer
Chinese lilac	leucothoe, coast	spreading euonymus
Chinese plumbago	leucothoe, drooping	spring cinquefoil
chionoides rhododendron	lilac time broom	spring glory forsythia
chirimen marlberry	lilac vine	spring torch Scotch heather
chollipo euonymus	lily-of-the-valley	squarrosa minima cypress
cinquefoil	little giant-dwarf arborvitae	staghorn sumac
cleyera, Japanese	lodense privet	stokesii Japanese holly
common witch hazel	lolendei monrovia pyracantha	strawberry guava
compacta-compact inkberry holly	lord roberts rhododendron	sumac, African
compacta-dwarf heavenly bamboo	magic carpet spiraea	summersweet
compacta-dwarf Japanese holly	marmalade bush	sunrise variegated abelia
convexa holly	media-old gold juniper	sunset potentilla
copper leaf	Mediterranean pink heather	sunspot euonymus
coral bark Japanese maple	mentor barberry	sweet olive
coral beauty smoke tree	mildred azalea	sweet pea shrub
coral bells azalea	miniature gardenia	tamariscifolia-tam juniper
cornish heather	minima glauca-dwarf arborvitae	tangerine potentilla
cotoneaster, bearberry	minuet weigela	tassel fern
cotoneaster, cranberry	mountain fire lily-of-the-valley	teenie genie brushcherry
cotoneaster, pyrenees	mugo-mugho pine	temple bells lily-of-the-valley
cotoneaster, rock	mystery gardenia	temple fire
cranberry bush	nakiana-dwarf deutzia	ternstroemia, Japanese
crape myrtle	nana compacta-heavenly bamboo	Texas dawn
creeping juniper	nana purpurea-heavenly bamboo	Texas sage
creeping mahonia	nana-dwarf hinoki cypress	torulosa cypress
crimson pygmy barberry	nana-dwarf Japaneses garden juniper	torulosa-hollywood juniper
cunninham white rhododendron	nana-dwarf yaupon holly	tradition azalea
daphnoides rhododendron	needle point holly	trailing indigo bush
dark knight bluebeard	neon flash spiraea	tribly rhododendron
david viburnum	nigra-dark American arborvitae	trumpet honeysuckle
dolchia spiraea	nordica-inkberry holly	unique rhododendron
doublefile viburnum	nova zembla rhododendron	valley rose lily-of-the-valley
dwarf andorra juniper	oleander	valley valentine lily-of-the-valley
dwarf brush cherry	orange flowering quince	variegated evergreen euonymus
dwarf burford holly	otto luykens English laurel	viburnum
dwarf Chinese holly	Paraguay nightshade	vulcan rhododendron
dwarf cranberry bush	parsonii juniper	walking stick
dwarf Japanese maple	pendula-weeping yaupon holly	waucabusa azalea
dwarf ninebark	pfitzer juniper	wax leaf privet
dwarf Oregon grape	pfitzerana glauca-blue juniper	wax ligustrum
dwarf pink flowering almond	pfitzerana-pfitzer juniper	wax myrtle
dwarf plumbago	pigmy date palm	weeping bottlebrush
dwarf scarlet wonder rhododendron	pink breath of heaven	wheeler's dwarf pittosporum
edward goucher abelia	pink dawn viburnum	wichita blue juniper
eichholz smoke tree	pink pixie	william penn barberry



elizabeth rhododendron	PJM rhododendron	wiltonii-blue carpet juniper
emerald arborvitae	plum yew	winter honeysuckle
emerald sea shore juniper	plumbago	wintergreen
escallonia	plumosa-andorra juniper	woadwaxen
false spiraea	praecox-early cotoneaster	woods dwarf-heavenly bamboo
fashion azalea	president clay azalea	woodwardii arborvitae
filifera-thread cypress	princes flower	yaupon holly
first snow spiraea	privet, glossy	yellow oleander
floppy disc cinquefoil	privet, Japanese	yellow tip ligustrum
flowering woodbine	prostrata juniper	yewpine
foemina-hicks juniper	purple bell heather	yucca
forest flame lily-of-the-valley		

---

### Terms and Conditions of Use

---

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

---

### Warranty Disclaimer

---

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

### Inherent Risks of Use

---

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risks shall be assumed by Buyer.

---

### Limitation of Remedies

---

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

EPA accepted 10/10/17