

PM 23 62719-184

Review
File 10/16

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division

OCT 15 1993

Douglas M. Roby
DOWELANCO
QUAD IV 9002 PURDUE RD
INDIANAPOLIS IN 46268

Subject: Label Amendment Submission of 06/28/93
to Comply with WPS Labeling Requirements
EPA Reg Nr. 52719-184
SONALAN 10G

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is accepted subject to the comments listed below.

Based on your certification, the Agency has accepted only those changes to your labeling which are necessary to comply with PR Notices 93-7 and 93-11, which reflect the WPS labeling requirements of 40 CFR part 156, subpart K. Any other labeling changes submitted in connection with this amendment application and not directly related to compliance with PR Notice 93-7 or 93-11 have neither been reviewed nor accepted by the Agency. If you wish to make any such changes, you must submit a separate amendment application proposing them. If your product registration is currently suspended, acceptance of this labeling amendment does not affect the suspension in any way.

A copy of your proposed labeling stamped, "Accepted with comments", is attached. Make any required changes described below and send three copies of final labeling as soon as it is available to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (H-7504C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, DC 20460-0001

BEST AVAILABLE COPY

20716

DOWELANCO
EPA Reg. Number: 62719-184
SONALAN 10G
Application date: 06/28/93
Original Submission

Hand or courier deliveries of final labels may be made to:

Document Processing Desk (FIN-LABEL)
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

The Agricultural Use Requirements referral statement in your proposed label does not reflect the language prescribed on page 43 of the Main Labeling Guidance.

Your label contains redundant WPS phrases that should be eliminated. Please see the attached comments.

Please correct the typographical errors circled on the draft before printing final labeling.

In your final labeling, the "Agricultural Use Requirements" text must be contained in a clearly separate box. This box may be set apart by a line, by another graphical device, by a different color background, or in any other way that clearly distinguishes it from surrounding text.

Sincerely,



Registration Support Branch
Registration Division

BEST AVAILABLE COPY

30416

8/9/93

1

WPS Labeling

(Editor's Note: Deleted text indicated by strike-through and added text underlined)

[Bag - Sonalan 10G]
[Front of Bag]

[Developed per EPA copy of 2/8/93 (Federal Copy Approved)]
[This bag supersedes 900-001838 (SLN for North Dakota)]

(logo) DowElanco

Sonalan* 10G

A selective herbicide for preemergence control of certain annual grasses and broadleaf weeds

Active Ingredient:

ethalfuralin: N-ethyl-N-(2-methyl-2-propenyl)-2,6-dinitro-4-(trifluoromethyl)benzenamine . 10%

Inert Ingredients . 90%

Total . 100%

Contains 5 pounds of active ingredient per 50 pound bag.

ACCEPTED
with COMMENTS
in EPA Letter Dated

OCT 15 1993

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

62719-184

Keep Out of Reach of Children

DANGER

PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

Refer to back panel of bag for precautionary information and Directions for Use including STORAGE AND DISPOSAL.

Notice: Read the entire label. Use only according to label directions. Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.

In case of emergency endangering health or the environment involving this product, call collect 517-636-4400.

Agricultural Chemical: Keep away from food, feedstuffs and water supplies. Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-184

EPA Est. 464-MI-1

*Trademark of DowElanco
DowElanco • Indianapolis, IN 46268, U.S.A.

Herbicide

Net Weight 50 lb

BEST AVAILABLE COPY

[Back of Bag]

Precautionary Statements

Hazards to Humans and Domestic Animals

DANGER

~~Precaucion al usuario: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.~~

~~Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.~~

~~(If you do not understand the label, find someone to explain it to you in detail.)~~

PELIGRO

Corrosive • Causes Irreversible Eye Damage • Harmful If Swallowed, Inhaled Or Absorbed Through The Skin

~~Wear goggles, face shield or safety glasses. Do not get in eyes. Avoid breathing dust and contact with skin, eyes or clothing. Use protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.~~

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If in eyes: Hold eyelids open and flush with a gentle stream of water for 15 minutes. Call a physician, preferably an ophthalmologist) or Poison Control Center immediately.

If swallowed: Do not induce vomiting. Call a physician or Poison Control Center. If available, administer activated charcoal (6-8 heaping teaspoons) with a large quantity of water. Do not give anything by mouth to an unconscious person. Immediately transport to a medical care facility and see a physician.

If inhaled: Remove individual to fresh air. Get medical attention if breathing difficulty occurs. If not breathing, give artificial respiration, preferably cardiopulmonary resuscitation, and get medical attention immediately.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation develops.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing equipment washwaters. Runoff or erosion from treated areas may be hazardous to fish in neighboring areas.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use before applying this product.

BEST AVAILABLE COPY

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container only. In case of spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, in accordance with applicable regulations, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Refer to label booklet in pocket of bag for additional precautionary information and Directions for Use.

[EDITOR NOTE: Insert here a fat, squat arrow pointing down]

Directions Enclosed [EDITOR NOTE: Turned at a 20° angle inside a box which has the top, right and bottom sides as bold dotted lines and the left side as a solid line]

20410

BEST AVAILABLE COPY

8/9/93

60416

[Right Gusset of Bag]

(logo) DowElanco

Sonalan* 10G

00010602
PI 3451960

BEST AVAILABLE COPY

70116

8/9/93

5

[Left Gusset of Bag]

(logo) DowEianco

Sonalan* 10G

00010602
PI 3451960

900-001916

8/9/93

6

70716

[Literature Booklet -- Sonalan 10G]
[Front Cover of Booklet]

[Developed per EPA copy of 2/8/93 (Federal Copy Approved)]

(logo) DowElanco

Sonalan* 10G

A selective herbicide for preemergence control of certain annual grasses and broadleaf weeds

EPA Reg. No. 62719-184
20410

900-001917

*Trademark of DowElanco
DowElanco • Indianapolis, IN 46268, U.S.A.

Herbicide

[Page 2 of Booklet]

Active Ingredient:

ethalfuralin: N-ethyl-N-(2-methyl-2-propenyl)-2,6-dinitro-4-(trifluoromethyl)benzenamine . 10%

Inert Ingredients . 90%

Total . 100%

Contains 5 pounds of active ingredient per 50 pound bag.

Precautionary Statements

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

DANGER PELIGRO:

~~Precaucion al usuario: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.~~

Corrosive • Causes Irreversible Eye Damage • Harmful If Swallowed, Inhaled Or Absorbed Through The Skin

~~Wear goggles, face shield or safety glasses. Do not get in eyes. Avoid breathing dust and contact with skin, eyes or clothing. Use protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.~~

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If in eyes: Hold eyelids open and flush with a gentle stream of water for 15 minutes. Call a physician, preferably an ophthalmologist) or Poison Control Center immediately.

If swallowed: Do not induce vomiting. Call a physician or Poison Control Center. If available, administer activated charcoal (6-8 heaping teaspoonfuls) with a large quantity of water. Do not give anything by mouth to an unconscious person. Immediately transport to a medical care facility and see a physician.

If inhaled: Remove individual to fresh air. Get medical attention if breathing difficulty occurs. If not breathing, give artificial respiration, preferably cardiopulmonary resuscitation, and get medical attention immediately.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation develops.

BEST AVAILABLE COPY

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing equipment washwaters. Runoff or erosion from treated areas may be hazardous to fish in neighboring areas.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use before applying this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exemptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container only. In case of spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, in accordance with applicable regulations, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

BEST AVAILABLE COPY

110716

General Information

Sonalan 10G is a selective herbicide for preemergence control of certain annual grasses and broadleaf weeds. Apply and incorporate Sonalan 10G before planting. Sonalan 10G does not control established weeds.

General Use precautions

Poor weed control and/or crop injury may result if directions are not followed. Uneven application or improper incorporation can result in erratic weed control or injury to the treated or rotational crops.

Do not graze or use treated crop for feed or forage including silage.

Do not apply Sonalan 10G to soils that are wet, cloddy, or subject to prolonged periods of flooding as poor weed control or crop injury may result.

Apply only in areas and to crops specified on this label.

Rotational Crop Restrictions

Replanting: If replanting is required, plant only those crops approved for treatment with Sonalan 10G or Sonalan E.C.

Sugar Beets or Red Beets: Do not plant sugar beets or red beets within 13 months following an application of 11 or more pounds per acre of Sonalan 10G. Where less than 11 pounds per acre of Sonalan 10G have been applied, sugar beets or red beets may be planted 8 months after an application of Sonalan 10G provided that the treated area is moldboard plowed to a depth of 12 inches prior to planting.

Spinach or Oats: In California and Arizona do not plant spinach or oats within 8 months following an application of 11 pounds or more of Sonalan 10G.

Application Directions

Soil Preparation

If necessary, use tillage to reduce ground cover (weeds or crop residue) to a level that will not interfere with the uniform application and soil incorporation of Sonalan 10G. The soil surface condition should allow Sonalan 10G to be thoroughly and uniformly mixed into the top 2 to 3 inches of soil. If this is not possible the soil should be tilled prior to application.

Application

Apply Sonalan 10G only with equipment capable of accurate calibration and uniform application of herbicide granules. Apply at recommended rate for soil texture. Follow calibration directions provided by the equipment manufacturer.

Soil Incorporation

Two incorporation passes are necessary for Sonalan 10G. The first should occur as soon as possible after application and should not be delayed more than 48 hours after application. For best weed control results the second incorporation should be delayed for 3 to 5 days after the first. Use incorporation equipment capable of thoroughly and uniformly mixing Sonalan 10G into the top 2 to 3 inches of the seedbed.

Cultivation After Planting

Sonalan 10G treated areas may be shallowly cultivated without loss of weed control activity. Avoid deep cultivation that could bring untreated soil to the soil surface resulting in loss of weed control.

BEST AVAILABLE COPY

2016

Soil Texture Guide for Application Rates

The amount of Sonalan 10G required for optimum weed control will vary, depending on soil texture and organic matter content. A fine-textured soil will require a higher application rate than a coarse-textured soil. Sonalan 10G application rates are based on the "Soil Texture Groups" (coarse, medium or fine) which correspond to the field soil textures to be treated. Using the table below, determine if the soil to be treated is coarse, medium or fine and use the application rate recommended for coarse, medium or fine soils.

Soil Texture Group	Soil Texture of the Soil to be Treated
Coarse soils (Light)	Sand, Loamy sand, Sandy loam
Medium soils	Loam, Silty clay loam†, Silt loam, Silt, Sandy clay loam†
Fine Soils (Heavy)	Clay, clay loam, Silty clay loam†, Silty clay, Sandy clay, Sandy clay loam†

†Silty clay loam and sandy clay loam soils are transitional soils and may be classified as either medium or fine textured soils. If silty clay loam or sandy clay loam soils are predominantly sand or silt, they are usually classified as medium textured soils. If they are predominantly clay, they are usually classified as fine textured soils.

Application Rate Ranges: Where a rate range is shown, use the lower rate for coarser textured soils or soils with low organic matter content. Use the higher rate in the rate range for finer textured soils and on soils containing more than 5% organic matter. Where soil texture is variable within the same field, use the lower recommended Sonalan 10G rate.

Weeds Controlled by Preplant Incorporated Treatment with Sonalan 10G**Grasses:**

<u>Common Name</u>	<u>Scientific Name</u>
annual bluegrass	<i>Poa annua</i>
barnyardgrass	<i>Echinochloa crus-galli</i>
crabgrass	<i>Digitaria</i> spp.
(large crabgrass)	
(smooth crabgrass)	
cupgrass, woolly	<i>Eriochloa villosa</i>
foxtail	<i>Setaria</i> spp.
(giant foxtail)	
(green foxtail)	
(foxtail millet)	
(pigeongrass)	
(yellow foxtail)	
goosegrass	<i>Eleusine indica</i>
johnsongrass (from seed)	<i>Sorghum halepense</i>
jungerice	<i>Echinochloa colonum</i>
oat, wild†	<i>Avena fatua</i>
panicum, fall	<i>Panicum dichotomiflorum</i>
panicum, Texas	<i>Panicum texanum</i>
(buffalograss)	
(Coloradograss)	
ryegrass, Italian	<i>Lolium multiflorum</i>
(annual ryegrass)	
sandbar, field	<i>Cenchrus incertus</i>
shattercane	<i>Sorghum bicolor</i>
(wild cane)	
signalgrass, broadleaf	<i>Brachiaria platyphylla</i>
(brachiaria)	
witchgrass	<i>Panicum capillare</i>

BEST AVAILABLE COPY

Broadleaf Weeds:

<u>Common Name</u>	<u>Scientific Name</u>
buckwheat, wild	<i>Polygonum convolvulus</i>
carpetweed	<i>Mollugo verticillata</i>
catchfly, conical	<i>Silene conica</i>
chickweed, common	<i>Stellaria media</i>
fiddleneck tarweed	<i>Amsinckia lycopsoides</i>
groundcherry, annual†† (lanceleaf groundcherry)	<i>Physalis lanceifolia</i>
henbit	<i>Lamium amplexicaule</i>
kochia	<i>Kochia scoparia</i>
lambsquarters, common	<i>Chenopodium album</i>
nightshade, annual†† (black nightshade) (eastern black nightshade) (hairy nightshade)	<i>Solanum</i> spp.
pigweed (prostrate pigweed) (redroot pigweed) (smooth pigweed) (tumble pigweed)	<i>Amaranthus</i> spp.
purslane, common	<i>Portulaca oleracea</i>
pursley, Florida	<i>Richardia scabra</i>
rockpurslane, redmaids	<i>Calandrinia caulescens</i>
thistle, Russian	<i>Salsola iberica</i>

†Sonalan 10G will provide wild oat control, except for seeds germinating below the zone treated with Sonalan 10G.

††See special use rates for control in dry bean.

Approved Crops

DRY BEANS

Apply Sonalan 10G as a preplant incorporated treatment. Follow soil preparation, application and incorporation procedures recommended in the "General Information" section of this label.

Broadcast Application Rates Per Acre: General weed control, except groundcherry and nightshade.

Soil Texture	Sonalan 10G (pounds)
Coarse	5.5 - 7.5
Medium	7.5 - 9.5
Fine	9.5 - 11.5

Broadcast Application Rates Per Acre: General weed control, including groundcherry and nightshade.

Precaution: Do not use these Sonalan 10G rates for garden bean, green bean, snap bean, or string bean.

Soil Texture	Sonalan 10G (pounds)
Coarse	11.5 - 13.0
Medium	13.0 - 15.0
Fine	15.0 - 17.0

BEST AVAILABLE COPY

Overlay Treatments Following Application of Sonalan 10G

For broader spectrum weed control, Sonalan 10G preplant incorporated treatment may be followed by preemergence overlay treatments using products registered for use on dry beans. Consult the manufacturer's labels for additional weeds controlled, directions for use, cautions and limitations before use.

Postemergence Treatments Following Application of Sonalan 10G

For broader spectrum weed control, Sonalan 10G preplant incorporated treatment may be followed by postemergence or layby treatments using products registered for use on dry beans. Consult the manufacturer's labels for additional weeds controlled, directions for use, cautions and limitations before use.

DRY PEAS

(Do Not Use In California)

Apply Sonalan 10G as a preplant incorporated treatment. Follow soil preparation, application and incorporation procedures recommended in the "General Information" section of this label.

Broadcast Application Rates Per Acre: General weed control, except groundcherry and night shade.

Soil Texture	Sonalan 10G (pounds)
Coarse	5.5 - 7.5
Medium	7.5

Precaution: Do not exceed recommended rates for Sonalan 10G in dry peas or crop injury may occur. Not all varieties of peas have been tested for tolerance to Sonalan 10G. Consult your seed contractor before applying Sonalan 10G to determine the tolerance of a particular pea variety. Application of Sonalan 10G to non-tolerant varieties may result in crop injury and reduced yield. Do not graze or forage crop grown in treated soil or cut for hay or silage.

PEANUTS

Apply Sonalan 10G as a preplant incorporated treatment. Follow soil preparation, application and incorporation procedures recommended in the "General Information" section of this label.

Broadcast Application Rates Per Acre:

Soil Texture	Sonalan 10G (pounds)
Coarse	5.5 - 7.5
Medium	7.5 - 9.5
Fine	9.5 - 11.5

Precaution: Do not graze or forage crop grown in treated soil or cut for hay or silage.

BEST AVAILABLE COPY

SOYBEANS

Apply Sonalan 10G as a preplant incorporated treatment. Follow soil preparation, application and incorporation procedures recommended in the "General Information" section of this label.

Broadcast Application Rates Per Acre: General weed control, except groundcherry and nightshade.

Soil Texture	Sonalan 10G (pounds)
Coarse	5.5 - 7.5
Medium	7.5 - 9.5
Fine	9.5 - 11.5

Broadcast Application Rates Per Acre: General weed control, including partial control of groundcherry and nightshade.

Sonalan 10G should be applied as a preplant incorporated treatment for partial control of groundcherry and nightshade on medium and fine textured soils. Follow recommended soil preparation and application procedures for Sonalan 10G.

Soil Texture	Sonalan 10G (pounds)
Medium	11.5
Fine	13.0

Precautions:

Soybeans should be planted no more than 2 inches deep after early season adverse weather conditions have passed, especially when using higher rate programs. Excessively deep planting or cool wet weather early in the growth cycle causes additional stress to soybean plants. These factors may result in a reduced stand, delayed maturity and reduced yield.

Overlay Treatments Following Sonalan 10G Application (Do Not Use in California)

For broader spectrum weed control, Sonalan 10G preplant incorporated treatment may be followed by preemergence overlay applications of Canopy, Dual, Gemini, Lasso, Lexone, Lorox, Lorox Plus, Preview, or Sensor. Consult these product labels for additional weeds controlled, directions for use, cautions and limitations before use.

Postemergence Treatments Following Sonalan 10G Application

For broader spectrum weed control, Sonalan 10G preplant incorporated treatment may be followed by postemergence or layby applications of Basagran, Blazer, Classic, Cobra, Pinnacle, Pursuit[†], Reflex, Storm or Tackle. Consult these product labels for additional weeds controlled, directions for use, cautions and limitations before use.

[†]Use of Pursuit is limited to certain states. Use Pursuit as a postemergence treatment following Sonalan 10G only in states specified on the label for Pursuit.

BEST AVAILABLE COPY

SUNFLOWERS

Apply Sonalan 10G as a preplant incorporated treatment. Follow soil preparation, application and incorporation procedures recommended in the "General Information" section of this label.

Broadcast Application Rates Per Acre: General weed control, except groundcherry and nightshade.

Soil Texture	Sonalan 10G (pounds)
Coarse	5.5 - 7.5
Medium	7.5 - 9.5
Fine	9.5 - 11.5

Overlay Treatments Following Sonalan 10G Application

For broader spectrum weed control, Sonalan 10G preplant incorporated treatment may be followed by preemergence overlay treatments using products registered for use on sunflowers. Consult the manufacturer's labels for additional weeds controlled, directions for use, cautions and limitations before use.

Warranty Disclaimer

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. DOWELANCO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of DowElanco or the seller. All such risks shall be assumed by Buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at DowElanco's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

DowElanco shall not be liable for losses or damages resulting from handling or use of this product unless DowElanco is promptly notified of such loss or damage in writing. In no case shall DowElanco be liable for consequential or incidental damages or losses.

The terms of the "Warranty Disclaimer" above and this "Limitation of Remedies" cannot be varied by any written or verbal statements or agreements. No employee or sales agent of DowElanco or the seller is authorized to vary or exceed the terms of the "Warranty Disclaimer" or this "Limitation of Remedies" in any manner.

Copyright© 1993 DowElanco

Label text is based on EPA accepted copy dated 02/08/93
Revised 06/10/93

Amendments:

- 1) Added labels statements required by the Worker Protections Standard (PR Notice 93-7).

BEST AVAILABLE COPY