18

Supplemental Labeling

ACCEPTED

MAY 2 2 1996

Under the Federal Insecticid Fungicide, and Rodenticide as omended, for the pectic registered under EPA Rec. 10.627/9 -184

DowElanco

DowElanco

9330 Zionsville Road

Indianapolis, Indiana 46268-1054 USA

Sonalan* 10G EPA Reg. No. 62719-184

For Distribution and Use Only in Minnesota, North Dakota and South Dakota

Fall or Spring Application to Ground with Standing Crop Stubble or Crop
Residue for Weed Control in Following Crop

- Special Use Directions for Minimum Tillage Situations -

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Sonalan 10G before applying. Carefully follow all
 precautionary statements and applicable use directions.
- Use of Sonalan 10G according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Sonalan 10G.

Directions for Use

Sonalan* 10G herbicide is a preemergence incorporated product that controls weeds as they germinate. Sonalan 10G will not control established weeds.

Application Timing

Sonalan 10G may be applied in fall or spring to ground with standing crop stubble or crop residue for weed control in the following crop. When applied in this manner, the first incorporation of Sonalan 10G must occur within 72 hours after application. The second incorporation must be delayed a minimum of 5 days after the first and may occur any time up to planting. This application is limited to use on ground that has not had a primary tillage treatment following crop harvest. Following application, plant only crops to which Sonalan 10G may be applied as a preplant incorporated treatment.

Broadcast Application Rates

Refer to the product label for Sonalan 10G for use directions for specific crops, including application rates and other use information. Apply at the recommended broadcast application rate for the crop, soil texture, and target weed species. When fall applied, use the highest rate in the rate range allowed for the soil texture to be treated.

Incorporation

Incorporate once using minimum tillage equipment such as the rotary harrow, field cultivator, under cutter chisel plow or other suitable tillage implement. It is desirable to loosen and mix surface soil sufficiently to trap Sonalan 10G, but maintain a surface cover of stubble and crop residue sufficient for erosion approtection. The second incorporation may occur any time prior to or in conjunction with tillage operations associated with planting.

			, , , , ,
*Trademark of DowElanco	•	Amendments:	· ()
:		1. Adds application to standing ເທຍ ຮັເບຸ່ນໃ	ole or crop '
(Label Code) Approved / /		residue and 72 hour incorporation Gela	у. '
Initial printing			