

Product Bulletin



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Sonalan* 10G

EPA Reg. No. 62719-184

Weed Control in Canola and Crambe

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Sonalan 10G before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Sonalan 10G according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Sonalan 10G, except as described below.

Directions for Use

Apply and incorporate Sonalan* 10G herbicide before planting. Application may be made in the fall or in the spring prior to planting canola or crambe. See instructions for fall application, application directions and soil-incorporation directions in the "General Information" section of the product label for Sonalan 10G.

Use the correct rate for each soil type to avoid crop injury. Prolonged wet or cold soil, deep seed placement, soil crusting, or application overlaps may combine to cause slowed or reduced emergence.

Broadcast Application Rates:

Soil Texture	Sonalan 10G (lb/acre) †
Coarse	5.5 – 7.5
Medium	7.5 – 9.5
Fine	9.5 – 11.5


† Use the higher rate in rate range where difficult-to-control weeds or high weed populations are anticipated.

Overlay Recommendations

For broader spectrum weed control, other products registered for use in canola or crambe may be applied as a sequential overlay treatment following application of Sonalan 10G. Consult the manufacturer's label for additional weeds controlled, directions for use, cautions and limitations before use.

Precautions and Restrictions

- Do not graze or forage crop grown in treated soil.
- **In Montana and Wyoming:** Plant only rapeseed (canola), crambe, safflower or sunflower as rotational crops in the crop year following the crop treated with 11.5 pounds per acre of Sonalan 10G. In addition to rapeseed (canola), crambe, safflower or sunflower, spring seeded barley grown under irrigated conditions can be planted in the crop year following the crop treated with a 7.5 pounds per acre rate of Sonalan 10G. If one of these specified crops is not planted, the land should be left idle or fallow for the entire year following the crop treated with Sonalan 10G.



*Trademark of Dow AgroSciences LLC
T1L / Sonalan 10G / Prop Sec 3 Canola & Crambe / 01-12-00

EPA-accepted: __/__/__
Initial printing.