

PM 23

62719-188

12-15-98

10418



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
62719-188

Date of Issuance:
DEC 15 1998

NOTICE OF PESTICIDE:
Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance:
Unconditional

Name of Pesticide Product:
Sonalan HFP

Name and Address of Registrant (include ZIP Code):

Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Based on your response to the Reregistration Eligibility Document, EPA has reregistered the product listed above. The following revisions must be incorporated into the labeling before the product is released for shipment:

- 1) Revise the sentence reading "Causes burns and irreversible eye damage" in the Precautionary Statements to read "Causes skin burns and irreversible eye damage."
- 2) Revise the sentence reading "Do not induce vomiting or give anything by mouth to an unconscious person" within the oral statement of the Statement of Practical Treatment to read "Do not induce vomiting."

Enclosed is a copy of your label stamped "Accepted with Comments". This action is taken under the authority of section 4(g)(2)(C) of the Federal Insecticide, Fungicide and Rodenticide Act, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain registration of your product.

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

Signature of Approving Official:

Date:

DEC 15 1998

DK: 305-7546: HB/PM 23
7505C
D. KENNY
12 15 1998

2/18

(Base Label):

(Logo) Dow AgroSciences LLC

Sonalan* HFP

A selective herbicide for preemergence control of certain annual grasses and broadleaf weeds.

Active Ingredient:

ethalfluralin: *N*-ethyl-*N*-(2-methyl-2-propenyl)-2, 6-dinitro-4-(trifluoromethyl)benzenamine 31.5%

Inert Ingredients 68.5%

Total 100.0%

ACCEPTED
with COMMENTS
In EPA Letter Dated

DEC 15 1998

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

62719-188

Contains 3 pounds active ingredient per gallon.

Keep Out of Reach of Children

DANGER PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Corrosive • Causes Burns And Irreversible Eye Damage • Harmful If Swallowed Or Inhaled • Prolonged or Frequently Repeated Skin Contact May Cause Allergic Reactions in Some Individuals

Do not get in eyes, on skin or on clothing. Avoid breathing vapor or spray mist.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category "H" on an EPA chemical resistance category selections chart.

Applicators and other handlers must wear:

- Coveralls worn over long-sleeved shirt and long pants
- Chemical-resistant gloves such as Barrier Laminate, or Viton
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing, or loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides, 40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eyelids open and flush with a gentle, steady stream of water for 15 minutes. Get medical attention.

If on skin: Wash with plenty of soap and water. Get medical attention.

If swallowed: Call a doctor or get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person. Drink promptly a large quantity of milk, egg white, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol.

If inhaled: Remove victim to fresh air. Get medical attention if breathing difficulty occurs. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.

Note to physician: Probable mucosal damage may contraindicate the use of gastric lavage.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Runoff or erosion from treated areas may be hazardous to fish in neighboring areas.

Physical or Chemical Hazards

Flammable. Do not use or store near heat sources or open flame.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Personal Protective Equipment (PPE), and User Safety Recommendations, and Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Avoid Freezing -- Store Above 40°F

EPA Reg. No. 62719-188

EPA Est. 1471-IN-2

*Trademark of Dow AgroSciences LLC
Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Herbicide**Net Contents 2.5 gal**

(Datapack Cover):

(Logo) Dow AgroSciences LLC

Sonalan* HFP

A selective herbicide for preemergence control of certain annual grasses and broadleaf weeds.

Active Ingredient:

ethalfluralin: *N*-ethyl-*N*-
(2-methyl-2-propenyl)-2, 6-dinitro-
4-(trifluoromethyl)benzenamine 31.5%

Inert Ingredients 68.5%

Total 100.0%

Contains 3 pounds active ingredient per gallon.

Keep Out of Reach of Children

DANGER PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Personal Protective Equipment (PPE), and User Safety Recommendations, and Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Avoid Freezing -- Store Above 40°F

EPA Reg. No. 62719-188

EPA Est. 1471-IN-2

*Trademark of Dow AgroSciences LLC
Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Herbicide

Net Contents 2.5 gal

5 of 18

(Page 1 through end):

Table of Contents	Page
Precautionary Statements:	
Hazards to Humans and Domestic Animals.....	
Personal Protective Equipment.....	
User Safety Recommendations.....	
First Aid.....	
Environmental Hazards.....	
Directions for Use:	
Agricultural Use Requirements.....	
Storage and Disposal.....	
General Information	
General Use Precautions.....	
Rotational Crop Restrictions.....	
Mixing Directions:	
Sonalan HFP - Alone.....	
Sonalan HFP in Tank Mix.....	
Mixing Order.....	
Application Directions:	
Ground Application.....	
Application with Liquid Fertilizer.....	
Liquid Fertilizer Mixing Instructions.....	
Testing for Tank Mix Compatibility in Liquid Fertilizers.....	
Application with Dry Bulk Fertilizer.....	
Limitation.....	
Impregnation.....	
Compliance with Individual State Regulations.....	
Soil Preparation.....	
Application Timing.....	
Soil-Incorporation Directions.....	
Soil Texture Guide for Application Rates.....	
Grasses and Broadleaf Weeds Controlled by Sonalan HFP.....	
Approved Crops	
Dry Beans.....	
Dry Peas.....	
Peanuts.....	
Soybeans.....	
Sunflower.....	
Warranty Disclaimer.....	
Inherent Risks of Use.....	
Limitation of Remedies.....	

60418

Precautionary Statements

Hazards to Humans and Domestic Animals

DANGER

Corrosive • Causes Burns And Irreversible Eye Damage • Harmful If Swallowed Or Inhaled • Prolonged or Frequently Repeated Skin Contact May Cause Allergic Reactions in Some Individuals

Do not get in eyes, on skin or on clothing. Avoid breathing vapor or spray mist.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category "H" on an EPA chemical resistance category selections chart.

Applicators and other handlers must wear:

- Coveralls worn over long-sleeved shirt and long pants
- Chemical-resistant gloves such as Barrier Laminate, or Viton
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing, or loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides, 40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eyelids open and flush with a gentle, steady stream of water for 15 minutes. Get medical attention.

If on skin: Wash with plenty of soap and water. Get medical attention.

If swallowed: Call a doctor or get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person. Drink promptly a large quantity of milk, egg white, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol.

If inhaled: Remove victim to fresh air. Get medical attention if breathing difficulty occurs. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.

Note to physician: Probable mucosal damage may contraindicate the use of gastric lavage.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water

7/2/18

when disposing of equipment washwaters or rinsate. Runoff or erosion from treated areas may be hazardous to fish in neighboring areas.

Physical or Chemical Hazards

Flammable. Do not use or store near heat sources or open flame.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of ~~12~~ 24 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls ~~over a long-sleeved~~ short-sleeved shirt and ~~long~~ short pants
- Chemical-resistant gloves such as Barrier Laminate, or Viton
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Storage: Avoid freezing. Store above 40°F (5°C). If frozen, poor weed control may result. Do not use or store near heat or open flame. Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Plastic Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Metal Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

8048

Bulk/Mini-bulk Tank Handling: Tanks that have been used for phenoxy-type herbicides should not be used for Sonalan HFP. To clean, triple rinse (or equivalent) and wash with appropriate cleaners before reusing.

General Information

Sonalan* HFP is a preemergence herbicide that controls many annual grasses and broadleaf weeds. Sonalan HFP controls weeds as they germinate. Sonalan HFP will not control established weeds.

General Use Precautions

Poor weed control and/or crop injury may result if directions are not followed. Uneven application, over application, or improper incorporation can result in erratic weed control or injury to the treated or rotational crops.

Do not graze or forage crop grown in treated soil or cut for hay or silage.

Do not apply Sonalan HFP to soils which are wet, cloddy, or subject to prolonged periods of flooding because poor weed control or crop injury may result.

Do not apply to any area not specified on the label.

Chemigation: Do not apply this product through any type of irrigation system, unless otherwise specified by a Dow AgroSciences product bulletin or supplemental label.

Rotational Crop Restrictions

Replanting

If replanting is required, replant only crops listed on this label.

Sugar Beets or Red Beets

Do not plant sugar beets or red beets within 13 months following an application of three (3) pints per acre or more of Sonalan HFP. Where less than three (3) pints per acre of Sonalan HFP have been applied, sugar beets or red beets may be planted no earlier than eight (8) months after an application and then only if the soil treated with Sonalan HFP is moldboard plowed to a depth of at least 12 inches prior to planting.

Spinach or Oats

In California and Arizona, do not plant spinach or oats within eight (8) months following an application of three (3) pints per acre or more of Sonalan HFP.

Special Rotational Crop Restrictions for the States of Montana and Wyoming

Plant only rapeseed (canola), safflower or sunflower as rotational crops in the crop year following the crop treated with 3 pints per acre of Sonalan HFP. In addition to rapeseed (canola), safflower or sunflower, spring seeded barley grown under irrigated conditions can be planted in the crop year following the crop treated with a 2 pint per acre rate of Sonalan HFP. If one of these specified crops is not planted, the land should be left idle or fallow for the entire year following the crop treated with Sonalan HFP.

Mixing Directions

Sonalan HFP - Alone

Start with a clean spray tank. Fill sprayer one-third (1/3) to one half (1/2) full with clean water or liquid fertilizer. Start agitation. Add correct quantity of Sonalan HFP, continue agitation and finish filling the tank.

9/2/18

Sonalan HFP in Tank Mix

Vigorous continuous agitation is required for all tank mixes. Sparger pipe agitators generally provide the best agitation in spray tanks.

Mixing Order: Fill the tank one-fourth (1/4) to one-third (1/3) full with clean water or liquid fertilizer. (See page X for additional liquid fertilizer mixing instructions). Start the agitation. Add different formulation types to the spray tank in the following order: dry flowables (DF), wettable powders (WP), aqueous suspensions (AS), flowables (F), or liquids (L). Allow each product type to completely disperse before adding another. Allow additional mixing and dispersion time when using dry flowable (DF) products. Continue agitation and fill tank to three-fourths (3/4) full, add the Sonalan HFP or other emulsifiable concentrates (EC) and mix thoroughly. Then add any solution (S) formulations, agitate and finish filling. Maintain agitation during filling and through application. If spraying and agitation must be stopped before the tank is empty, the materials may settle to the bottom. In this case, it is important to resuspend all of the material in the bottom of the tank before continuing the spray application. A sparger agitator is particularly useful for this purpose. Sometimes it is more difficult to resuspend settled material than it is to suspend it originally.

Carefully follow all mixing instructions for each material added to the tank. Initial dispersion of dry or flowable formulations can be improved by mixing with a small amount of water (slurrying) and pouring the slurry through a 20 or 35 mesh wetting screen in the top of the spray tank. Line screens in the tank should be no finer than 50 mesh (100 mesh is finer than 50 mesh).

If a buildup of material on the walls of the spray tank is observed, wash the tank with soapy water between fillings. Rinse and continue the spraying operation. Clean the tank, lines and screens thoroughly after use. Check the sprayer daily to ensure proper calibration and uniform application. Do not apply Sonalan HFP under wind conditions which may cause spray drift to nontarget areas or non-uniform application.

Application Directions

Ground Application: Apply Sonalan HFP in at least five (5) gallons of water or liquid fertilizer per acre (broadcast basis). Use any properly calibrated, low pressure herbicide sprayer that will apply the spray uniformly.

Application with Liquid Fertilizer

Sonalan HFP may be mixed with most liquid fertilizer materials. Combinations of Sonalan HFP with solutions and suspension type fertilizers provide weed and grass control equal to the same rates of Sonalan HFP applied in water. For application in liquid fertilizer, follow label recommendations regarding crops, application rates, incorporation directions and precautions for Sonalan HFP alone.

Compliance with individual state regulations relating to liquid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company offering the fertilizer and chemical mixture for sale.

Liquid Fertilizer Mixing Instructions: Emulsifiable concentrates, such as Sonalan HFP, can be mixed with liquid fertilizers. In all cases, continuous agitation is required to prevent the Sonalan HFP from rising to the surface as an oily layer. When necessary, (see "Testing for Tank Mix Compatibility in Liquid Fertilizers") a compatibility agent (a phosphate ester type surfactant designed to be used with liquid fertilizers) can be used to ensure that the Sonalan HFP emulsifies properly (i.e. forms a uniform mixture with a milky appearance rather than an oily layer). Compatibility agents can be used at rates as low as one and one-half (1 1/2) to two (2) pints per ton of liquid fertilizer and should be mixed well with the fertilizer before adding any emulsifiable concentrate product. The use of compatibility agents is especially important when tank mixing emulsifiable concentrates (EC) with dry flowables (DF), wettable powders (WP), flowables (F), liquids (L), aqueous suspensions (AS), or solutions (S) in liquid fertilizer. If the emulsion is not properly formed and the EC rises to the surface of the fertilizer as an oil ("oils out"), the oil may combine with the wettable powder, flowable, or suspension to form oily curds (viscous phase) which are difficult to disperse.

10/18

Testing for Tank Mix Compatibility in Liquid Fertilizers: Emulsifiable concentrates alone or in tank mixture with dry flowables (DF), wettable powders (WP), liquids (L), flowables (F), aqueous suspensions (AS), or solutions (S) may not combine properly with some fluid fertilizer materials. Small quantities should always be tested before full-scale mixing. This will determine whether a compatibility agent is needed and which agent does the best job. Use only a phosphate ester type surfactant designed to be used with liquid fertilizers.

Use the following test to select the correct agent for your mixture.

1. Add one (1) pint of the liquid fertilizer to a quart jar.
2. Add one (1) to four (4) teaspoonful(s) of the DF, WP, L, F, or AS formulations (depending on the recommended rate per acre) to the liquid fertilizer. Close jar and agitate until dispersed evenly in the fertilizer. If the materials do not disperse well, it may be necessary to slurry the chemicals in water before adding to the fertilizer.
3. After dispersing the materials (Step 2), add three (3) to four (4) teaspoonfuls of the Sonalan HFP to the jar and shake well. Add solution herbicides to the mixture last and agitate. Observe the jar for about 10 minutes. If the materials rise to the surface and form a thick layer (oily curds) which will not disperse when agitated, a compatibility agent is needed. If the mixture is easily dispersed to its original state with slight agitation, no agent is needed, but good agitation must be provided in the fertilizer spray tank.
4. If the need for a compatibility agent is shown in Step 3, use a clean quart jar, and starting at Step 1 above, add one-half (1/2) teaspoonful of the compatibility agent to the liquid fertilizer. Mix well. Repeat Steps 2 and 3.

An effective compatibility agent will cause the mixture to remain uniformly mixed with little or no separating or oil rising to the surface for one-half (1/2) hour or longer. If slight separation does occur, two (2) to three (3) inversions of the jar should give a uniform remix. If oily curds form which will not disperse, more agent or another agent should be tried.

Use a clean jar for each test. The compatible mixture will have a uniform appearance and will be relatively easy to keep mixed with gentle agitation of the jar.

Application with Dry Bulk Fertilizer

Dry bulk fertilizer may be impregnated or coated with Sonalan HFP. Application of dry bulk fertilizer impregnated with Sonalan HFP provides weed and grass control equal to the same rates of Sonalan HFP applied in liquid carriers. Follow label recommendations regarding crops, application rates, and precautions for Sonalan HFP alone. Spread the fertilizer/chemical mixture with properly calibrated application equipment. Be certain the material is applied uniformly to the soil surface. When impregnated on dry bulk fertilizer, Sonalan HFP must be incorporated 2 times (see Soil-Incorporation Directions).

Limitations: Apply a minimum of 200 pounds per acre of dry fertilizer impregnated with Sonalan HFP at the recommended rate. Most dry fertilizers can be used for impregnation with Sonalan HFP. When coated ammonium nitrate and/or limestone are used alone, do not impregnate with Sonalan HFP. These materials will not absorb the herbicide. Blends containing a mixture of these materials can be impregnated.

Impregnation: Use any closed drum, belt, ribbon or other commonly used dry bulk fertilizer blender. Nozzles used to spray the Sonalan HFP onto the fertilizer should be placed to provide uniform spray coverage. Add water to the Sonalan HFP to give a total volume of at least six (6) pints per ton of fertilizer.

Compliance with individual state regulations relating to dry bulk fertilizer blending, registration, labeling and application are the responsibility of the individual and/or company offering the fertilizer and chemical mixture for sale.

Soil Preparation

Factors influencing the uniform application and incorporation of Sonalan HFP:

Crop Residues or Existing Weeds - Excessive amounts of crop residues or existing weeds may interfere with the incorporation of Sonalan HFP into the soil. A manageable level of such ground cover will allow Sonalan HFP to be uniformly mixed into the top two (2) to three (3) inches of the final seedbed. If the level of the ground cover is such that this cannot be done, the soil must be tilled prior to the application of Sonalan HFP.

Soil Roughness - The soil surface should be smooth enough so that you can operate a sprayer and incorporation equipment without wheel slippage and at speeds sufficient to ensure uniform application and incorporation of Sonalan HFP.

General Soil Conditions - To assure uniform incorporation of Sonalan HFP, soil moisture conditions should be such that large clods can be broken up during the incorporation process.

Application Timing

Spring Application

Apply and incorporate Sonalan HFP any time after January 1 when soil can be worked and is in a condition which allows thorough mixing to insure uniform incorporation.

Fall Application

When applied with dry bulk fertilizer, Sonalan HFP may be applied and incorporated in the fall prior to a spring planting of dry beans or sunflowers. Apply Sonalan HFP with dry bulk fertilizer anytime between October 1 and December 31 at the recommended rates for crops and soil textures listed in the Approved Crops section of this label. Fall application of Sonalan HFP is not recommended on fields which remain wet or are subject to periods of flooding.

Soil-Incorporation Directions

Use incorporation equipment that thoroughly and uniformly mixes Sonalan HFP into the top two (2) to three (3) inches of the final seedbed, or erratic weed control and/or crop injury may result. Incorporation should occur as soon as possible after application and, to prevent loss of herbicidal activity, should not be delayed more than 48 hours after application.

Incorporation in Bedded Culture

Application prior to bedding: Apply Sonalan HFP and mix thoroughly into the top two (2) to three (3) inches of soil. The bedding operation provides additional mixing. Do not expose untreated soil during post-bedding operations.†

Application after bedding: Knock off beds to planting height before applying Sonalan HFP. Apply Sonalan HFP and mix thoroughly with equipment that will conform to the bed shape. Do not expose untreated soil during the incorporation process.†

†Avoid removal of treated soil from the seedbed before or during the planting operation. Exposure of untreated soil during planting will allow weeds to germinate in the drill row.

Incorporation When Applied with Dry Bulk Fertilizer: When impregnated on dry bulk fertilizer, Sonalan HFP must be incorporated 2 times. The first incorporation should occur as soon as possible after application, but not delayed more than 48 hours after application. For best weed control results, the second incorporation should be delayed at least 5 days after the first and occur before planting. When fall applied, the second incorporation may occur in the spring as a part of tillage associated with seedbed preparation. The second incorporation should be made in a different direction, and to avoid bringing untreated soil to the surface, should not be deeper than the first.

2018

Optional Methods of Incorporation for Soybeans, Dry Beans and Peanuts on Medium and Coarse Textured Soils: Sonalan HFP may be applied as a surface application and incorporated by rainfall or sprinkler irrigation. Soil should be in good tilth and free of clods. Make applications to tilled land and/or standing or chopped stubble from the previous season's crop. Conditions should permit thorough movement into the top 2-3 inches of the final seedbed. Application should be made immediately after tillage and after planting, but prior to rainfall or irrigation. Rainfall or sprinkler irrigation prior to application will tend to consolidate and seal the soil surface and prevent the downward movement of Sonalan HFP that is expected under porous, open, recently tilled conditions.

If continuous rainfall in the amount of at least one-half to one-inch occurs within 2 days after application of Sonalan HFP, no further incorporation is required. If the prescribed amount of rainfall is not anticipated, supplemental overhead sprinkler irrigation of at least one-half to one-inch of water may be applied within 2 days after application. Supplemental irrigation can be applied through a center pivot, solid set or hand moved sprinkler system. Furrow irrigation is prohibited. Mechanical incorporation methods should be used to incorporate Sonalan HFP if the required amount of rainfall or supplemental irrigation does not occur within two days after application. If mechanical incorporation is used, adjust equipment so as to not disturb planted seed.

Note: For crops with application rates that exceed 3.7 pints per acre, do not apply more than 3.7 pints (1.4 lb a.i.)/acre, regardless of the application rate in the specific use directions, if incorporation is to be accomplished by supplemental overhead sprinkler irrigation. Do not apply more than 2.0 pints (0.75 lb a.i.)/acre, regardless of the application rate in the specific use directions, if incorporation is to be accomplished by rainfall only.

Cultivation after Planting

Soil treated with Sonalan HFP may be shallowly cultivated without reducing the weed control activity of Sonalan HFP. Do not cultivate deeper than the treated soil since weeds may germinate where untreated soil is brought to the surface.

Soil Texture Guide for Application Rates

Use rates of Sonalan HFP for specific crops are based on the "Soil Texture Class" (coarse, medium, or fine) of the soil to be treated, and organic matter content. A fine-textured soil (clay loam) requires a higher use rate than a coarse-textured soil (loamy sand). Field soil textures, grouped according to Soil Texture Class, are listed in the table below and should be used to determine if the soil to be treated is coarse, medium or fine. Do not exceed recommended use rates for specific crops or target weed species.

Soil Texture Class	Field Soil Texture Group
Coarse soils: (Light)	Sand, loamy sand, sandy loam
Medium soils:	Loam, silty clay loam [†] , silt loam, silt, sandy clay loam [†]
Fine soils: (Heavy)	Clay, clay loam, silty clay loam [†] , silty clay, sandy clay, sandy clay loam [†]

[†]Silty clay loam and sandy clay loam soils are transitional soils and may be classified as either medium or fine textured soils. If silty clay loam or sandy clay loam soils are predominantly sand or silt they are usually classified as medium textured soils. If they are predominantly clay, they are usually classified as fine textured soils.

Application Rate Ranges: Where a rate range is shown, use the lower rate for more coarse-textured soils in the Field Soil Texture Group or soils with low organic matter content. Use the higher rate in the rate range for more fine-textured soils in the Field Soil Texture Group and soils containing more than 5% organic matter. Where soil texture is variable within the same field, use the lower recommended rate for

Sonalan HFP. Sonalan HFP is not recommended for use on soils containing more than 10% organic matter.

Grasses and Broadleaf Weeds Controlled by Sonalan HFP

Sonalan HFP applied preplant soil incorporated will control the following weeds:

Grasses:

Common Name

Scientific Name

annual bluegrass
 barnyardgrass
 crabgrass
 (large crabgrass)
 (small crabgrass)
 cupgrass, woolly
 foxtail
 (giant foxtail)
 (green foxtail)
 (foxtail millet)
 (pigeongrass)
 (yellow foxtail)
 johnsongrass (from seed)
 junglerice
 oat, wild†
 panicum, fall
 panicum, Texas
 (buffalograss)
 (Colorado grass)
 ryegrass, Italian
 (annual ryegrass)
 sandbur, field
 shattercane
 (wild cane)
 signalgrass, broadleaf
 (brachiaria)
 witchgrass

Poa annua
Echinochloa crus-galli
Digitaria spp.

Eriochloa villosa
Setaria spp.

Sorghum halepense
Echinochloa colonum
Avena fatua
Panicum dichotomiflorum
Panicum texanum

Lolium multiflorum

Cenchrus incertus
Sorghum bicolor

Brachiaria platyphylla

Panicum capillare

Broadleaf Weeds:

Common Name

Scientific Name

buckwheat, wild
 carpetweed
 catchfly, conical
 chickweed, common
 fiddleneck tarweed
 groundcherry†† (annual)
 (lanceleaf
 groundcherry)
 (Wright's
 groundcherry)
 henbit
 kochia
 lambsquarters, common
 nightshade†† (annual)
 (black nightshade)
 (eastern black
 nightshade)†††

Polygonum convolvulus
Mollugo verticillata
Silene conica
Stellaria media
Amsinckia lycopsoides

Physalis lanceifolia

Physalis wrightii

Lamium amplexicaule
Kochia scoparia
Chenopodium album
Solanum spp.

140/18

(hairy nightshade)	
pigweed	<i>Amaranthus</i> spp.
(prostrate pigweed)	
(redroot pigweed)	
(smooth pigweed)	
(tumble pigweed)	
purslane, common	<i>Portulaca oleracea</i>
pusley, Florida	<i>Richardia scabra</i>
rockpurslane, redmaids	<i>Calandrinia caulescens</i>
thistle, Russian	<i>Salsola kali</i>

†Sonalan HFP will provide wild oat control except those germinating below the zone treated with Sonalan HFP.

††See special instructions for control in dry bean.

†††See special instructions for partial control in soybean.

Approved Crops

DRY BEANS

(Does not include soybean)

Sonalan HFP - Alone

Apply and incorporate Sonalan HFP in the spring before planting or in the fall. Fall application may be made only with dry bulk fertilizer. See instructions for fall application of Sonalan HFP under the heading "Application Timing" in the General Information section of this label. Follow soil preparation, application, and incorporation procedures recommended in the "General Information" section of this label.

Broadcast Application Rates Per Acre: General weed control, except groundcherry and nightshade.

Soil Texture	Sonalan HFP
	(pints)
Coarse	1 1/2-2
Medium	2-2 1/2
Fine	2 1/2-3

Broadcast Application Rates Per Acre: General weed control, including groundcherry and nightshade from seed. Two incorporation passes are required for control of ground cherry and eastern black nightshade.

Soil Texture	Sonalan HFP
	(pints)
Coarse	3-3 1/2
Medium	3 1/2-4
Fine	4-4 1/2

Precaution: When high rates for control of nightshade and groundcherry are applied, rotate only to crops listed on this label.

15 of 18

Tank Mix, Overlay or Postemergence Recommendations

For broader spectrum weed control, other products registered for use in dry beans may be applied in tank mix combination or as a sequential overlay or postemergence treatment following application of Sonalan HFP. Consult the manufacturer's label for additional weeds controlled, directions for use, cautions and limitations before use. See detailed information for tank mixing in the General Information section of this label.

DRY PEAS

(Not for Use in California)

General weed control - except groundcherry and nightshade: Apply and incorporate Sonalan HFP in the spring before planting or in the fall. Fall application may be made only with dry bulk fertilizer. See instructions for fall application of Sonalan HFP under the heading "Application Timing" in the General Information section of this label. Follow soil preparation, application, and incorporation procedures recommended in the "General Information" section of this label.

Broadcast Rates/Acre

Soil Texture	Sonalan HFP
	(pints)
Coarse	1 1/2-2
Medium	2

Precaution: Do not exceed these rates of Sonalan HFP for weed control in dry peas or crop injury may occur. Not all varieties of peas have been tested for tolerance to Sonalan HFP. Consult your seed contractor before applying Sonalan HFP to determine the tolerance of a particular pea variety. Application of Sonalan HFP to non-tolerant pea varieties may result in crop injury and reduced yields. Do not graze or forage crop grown in treated soil or cut for hay or silage.

PEANUTS

Sonalan HFP - Alone

Apply Sonalan HFP in the spring as a preplant incorporated treatment. Follow soil preparation, application, and incorporation procedures recommended in the "General Information" section of this label.

Broadcast Rates/Acre

Soil Texture	Sonalan HFP
	(pints)
Coarse	1 1/2-2
Medium	2-2 1/2
Fine	2 1/2-3

Do not graze or forage crop grown in treated soil or cut for hay or silage.

Tank Mix or Overlay Recommendations

For broader spectrum weed control, other products registered for use in peanuts may be applied in tank mix combination or as at-cracking, sequential or overlay treatments following application of Sonalan HFP. Consult the manufacturer's label for additional weeds controlled, directions for use, cautions and

limitations before use. See detailed information for tank mixing in the General Information section of this label.

SOYBEANS

Sonalan HFP - Alone

Apply and incorporate Sonalan HFP in the spring before planting or in the fall. Fall application may be made only with dry bulk fertilizer. See instructions for fall application of Sonalan HFP under the heading "Application Timing" in the General Information section of this label. Follow soil preparation, application, and incorporation procedures recommended in the "General Information" section of this label.

Broadcast Application Rates Per Acre: General weed control, except groundcherry and nightshade.

Soil Texture	Sonalan HFP
	(pints)
Coarse	1 1/2 - 2
Medium	2 - 2 1/2
Fine	2 1/2 - 3

Broadcast Application Rates Per Acre: General weed control, including partial control of eastern black nightshade and groundcherry. Two incorporation passes are required for partial control of ground cherry and eastern black nightshade.

Soil Texture	Sonalan HFP
	(pints)
Medium	3
Fine	3 1/2

Precautions:

- When high rates for control of nightshade and groundcherry are applied, rotate only to crops listed on this label.
- Soybeans should be planted no more than two (2) inches deep after early season adverse weather conditions have passed, especially when using higher rate programs. Excessively deep planting or cool, wet weather early in the growth cycle causes additional stress to the soybean plant. These factors may result in reduced stands, delayed maturity and reduced yields.

Tank Mix, Overlay or Postemergence Recommendations

For broader spectrum weed control, other products registered for use in soybean may be applied in tank mix combination or as a sequential overlay or postemergence treatment following application of Sonalan HFP†. Consult the manufacturer's label for additional weeds controlled, directions for use, cautions and limitations before use. See detailed information for tank mixing in the General Information section of this label.

†Do not use Scepter in tank mix combination with Sonalan HFP or as a sequential overlay or postemergence application following Sonalan HFP.

SUNFLOWER

Sonalan HFP - Alone

Apply and incorporate Sonalan HFP in the spring before planting or in the fall. Fall application may be made only with dry bulk fertilizer. See instructions for fall application of Sonalan HFP under the heading "Application Timing" in the General Information section of this label. Follow soil preparation, application, and incorporation procedures recommended in the "General Information" section of this label.

Broadcast Application Rates Per Acre: General weed control, except groundcherry and nightshade.

Soil Texture	Sonalan HFP
	(pints)
Coarse	1 1/2-2
Medium	2-2 1/2
Fine	2 1/2-3

Broadcast Application Rates Per Acre: General weed control, including groundcherry and nightshade from seed.

Soil Texture	Sonalan HFP
	(pints)
Coarse	3-3 1/2
Medium	3 1/2-4
Fine	4-4 1/2

Precautions

- Do not graze or forage crop grown in treated soil or cut for hay or silage.
- When high rates for control of nightshade and groundcherry are applied, rotate only to crops listed on this label.

Tank Mix or Overlay Recommendations

For broader spectrum weed control, other products registered for use in sunflowers may be applied in tank mix combination or as a sequential overlay treatment following application of Sonalan HFP. Consult the manufacturer's label for additional weeds controlled, directions for use, cautions and limitations before use. See detailed information for tank mixing in the General Information section of this label.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

*Trademark of Dow AgroSciences LLC

EPA Accepted: / /