

Product Bulletin



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Sonalan* HFP

EPA Reg. No. 62719-188

Weed Control in Canola and Crambe

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Sonalan HFP before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Sonalan HFP according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Sonalan HFP, except as described below.

Directions for Use

Sonalan HFP - Alone


Apply and incorporate Sonalan* HFP herbicide before planting. Application may be made in the fall or in the spring prior to planting canola or crambe. See instructions for fall application, application directions and soil-incorporation directions in the "General Information" section of the product label for Sonalan HFP.

Use the correct rate for each soil type to avoid crop injury. Prolonged wet or cold soils, deep seed placement, soil crusting, or application overlaps may combine to cause slowed or reduced emergence.

Broadcast Application Rates:

Soil Texture	Sonalan HFP (pt/acre) †
Coarse	1 1/2-2
Medium	2-2 1/2
Fine	2 1/2-3

† Use the higher rate in rate range where difficult-to-control weeds or high weed populations are anticipated.



Tank Mix or Overlay Recommendations

For broader spectrum weed control, other products registered for use in canola or crambe may be applied in tank mix combination or as a sequential overlay treatment following application of Sonalan HFP. Consult the manufacturer's label for additional weeds controlled, directions for use, cautions and limitations before use. See detailed information for tank mixing in the "General Information" section of the product label for Sonalan HFP.

Precautions and Restrictions

- Do not graze or forage crop grown in treated soil.
- **In Montana and Wyoming:** Plant only rapeseed (canola), crambe, safflower or sunflower as rotational crops in the crop year following the crop treated with 3 pints per acre of Sonalan HFP. In addition to rapeseed (canola), crambe, safflower or sunflower, spring seeded barley grown under irrigated conditions can be planted in the crop year following the crop treated with a 2 pints per acre rate of Sonalan HFP. If one of these specified crops is not planted, the land should be left idle or fallow for the entire year following the crop treated with Sonalan HFP.

*Trademark of Dow AgroSciences LLC
T1L / Sonalan HFP / Prop Sec 3 Canola & Crambe / 01-17-00

EPA-accepted: / /
Initial printing.