

(Logo) DowElanco

Turf Fertilizer - Contains Gallery* plus Team*

A Balanced Turf Fertilizer Containing GALLERY and TEAM, Selective Preemergence Herbicides for Control of Certain Broadleaf Weeds and Annual Grasses in Established Perennial Turfgrasses.

Active Ingredients:

isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers.....	0.29%
trifluralin: α,α,α -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine.....	0.27%
benefin: <i>N</i> -butyl- <i>N</i> -ethyl- α,α,α -trifluoro-2,6-dinitro- <i>p</i> -toluidine.....	0.53%
Inert Ingredients.....	98.91%
Total.....	100.00%

GUARANTEED FERTILIZER ANALYSIS	
Total Nitrogen.....	18.0%
Available Phosphoric Acid (P2O5).....	5.0%
Soluble Potash (K2O).....	9.0%

Contains 0.2725 pounds active ingredient per 25 pound bag.

Keep Out of Reach of Children CAUTION

See back or side panel of container for additional precautionary statements.

Notice: Read the entire label. Use only according to label directions. Before buying this product, read Warranty Disclaimer and Limitation of Remedies Sections elsewhere on this label.

In case of emergency endangering health or the environment involving this product, call collect 517-636-4400

Do Not Ship or Store with Food, Feeds, Drugs, or Clothing.


E.P.A. Reg. No. 62719-192

E.P.A. Est. _____

*Trademark of DowElanco

DowElanco • Indianapolis, Indiana 46268, U.S.A.

**Herbicide
Net Contents - 25 Pounds**



1 of 6

Precautionary Statements

Keep Out of Reach of Children**Hazards to Humans and Domestic Animals****CAUTION****Causes Eye Irritation • Harmful If Swallowed or Inhaled**

Avoid breathing dust and contact with eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

First Aid

If In eyes: Flush eyes with plenty of water. Get medical attention if irritation persists.

If swallowed: Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or inadvertent application to areas adjacent to treated turfgrasses may result in reduced germination or emergence of nontarget plants. Do not contaminate water when disposing of equipment washwaters.

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read all directions carefully before applying this product.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

General Information

This turf fertilizer product contains GALLERY and TEAM, preemergence herbicides for control of certain broadleaf weeds and annual grasses in established turfgrasses. Apply in late summer or early fall to control fall approximately 3 weeks prior to germination of winter annual weeds, or in early spring, 3 weeks prior to germination of weeds which emerge in spring or early summer.

This product will not control established weeds. Existing weeds should be controlled with postemergence herbicides. To insure optimal fertilizer and herbicidal response, water treated areas as soon as possible after application. A single rainfall or sprinkler irrigation of 0.5 inch or more is required to activate this product.

Important Use Precautions:

- Do not apply to non-turf areas such as flower beds or gardens. Take precautions to prevent inadvertent application from rotary-type applicators to such areas. Herbicides in this product prevent seed emergence and may injure bedding plants.
- This product may injure turf that is weakened due to severe winter weather or is not well established.
- Do not apply this product in the spring and summer to tall fescue turfgrass seeded the previous fall.
- Do not apply in the spring to Bermudagrass which has been overseeded with winter annual grasses since thinning may occur.
- Do not apply to newly sprigged or hydromulched areas of Bermudagrass, St. Augustinegrass, centipedegrass or zoysiagrass until turf is well established.
- Do not apply to dichondra.
- Do not apply to golf course putting greens.
- Any cultural practices that disturb the soil, such as aerification or verticutting should be done prior to application of this product.
- Do not overdose or overlap during application in such a way as to apply in excess of labeled rates as turf injury may occur.
- **Reseeding:** Do not reseed treated areas for a minimum of 8 weeks after a single application of 275 pounds per acre, or 12 to 16 weeks after a single application of 375 pounds per acre.
- Users who wish to use this product on turfgrass species not recommended on this label may determine the suitability for such uses by treating a small area at a recommended rate. Prior to treating a larger area, the treatment should be observed for any sign of herbicidal injury for 30 to 60 days to determine if the treatment is safe to the target turf species. **THE USER ASSUMES RESPONSIBILITY FOR ANY PLANT DAMAGE OR OTHER LIABILITY RESULTING FROM USE OF GALLERY ON TURF SPECIES NOT RECOMMENDED ON THIS LABEL.**

This Product Is Recommended for Use on the Following Established Warm and Cool Season Perennial Turfgrass Species:

Established Cool Season Grasses

Common Name	Scientific Name
Bentgrass, colonial	<i>Agrostis tenuis</i>
Bentgrass, creeping	<i>Agrostis stolonifera</i>
Bluegrass, Kentucky	<i>Poa pratensis</i>
Fescue, chewings	<i>Festuca rubra</i> var. <i>commutata</i>
Fescue, creeping red	<i>Festuca rubra</i>
Fescue, sheep	<i>Festuca ovina</i>
Fescue, tall	<i>Festuca arundinaceae</i>
Ryegrass, perennial	<i>Lolium perenne</i>

11/2/92

Established Warm Season Grasses

Common Name	Scientific Name
Bahiagrass	<i>Paspalum notatum</i> var. <i>Saurae parodi</i>
Bermudagrass	<i>Cynodon dactylon</i>
Centipedegrass	<i>Eremochloa ophiuroides</i>
Fescue, tall	<i>Festuca arundinaceae</i>
St. Augustinegrass	<i>Stenotaphrum secundatum</i>
Zoysiagrass	<i>Zoysia japonica</i>
Zoysiagrass	<i>Zoysia tenuifolia</i>

Application

Application Equipment and Application Techniques

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to manufacturer's directions. Check frequently to be sure equipment is operating properly and distributing product uniformly. Avoid use of spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turf injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. A splash board mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

Broadcast Application Rates

Apply GALLERY plus TEAM Turf Fertilizer at a rate of 250 to 375 pounds per acre. A minimum of 250 pounds per acre is recommended for optimum results.

	Cool Season Turfgrasses	Warm Season Turfgrasses
Summer Annual Grass Weeds (Spring Application)	250 lbs/acre 5.75 lbs/1000 sq ft	250 to 375 lbs/acre 5.75 to 8.6 lbs/1000 sq ft
Winter Annual Grass Weeds (Fall application??)	275 lbs/acre 5.75 lbs/1000 sq ft	250 to 375 lbs/acre 5.75 to 8.6 lbs/1000 sq ft

Weeds Controlled

Weeds controlled by GALLERY plus TEAM Turf Fertilizer applied prior to weed germination at 250 to 375 pounds per acre (5.75 to 8.6 lbs/1000 sq ft)

Common Name	Scientific Name
Annual bluegrass	<i>Poa annua</i>
Barnyardgrass	<i>Echinochloa crus galli</i>
Burweed, lawn	<i>Soliva pterosperma</i>
Buttercup	<i>Ranunculus</i> spp.
Carpetweed	<i>Mollugo verticillata</i>
Celery, wild	<i>Apium leptophyllum</i>
Chickweed, common	<i>Stellaria media</i>
Chickweed, mouseear	<i>Cerastium vulgatum</i>
Crabgrass	<i>Digitaria</i> spp.
(Large crabgrass)	
(Smooth crabgrass)	
Foolsparsley	<i>Aethusa cynapium</i>
Foxtails	<i>Setaria</i> spp.
(Bottlegrass)	

5.86

(Bristlegrass)	
(Giant foxtail)	
(Green foxtail)	
(Pigeongrass)	
(Robust foxtail)	
Geranium, Carolina	<i>Geranium carolinianum</i>
Goosegrass	<i>Eleusine indica</i>
Henbit	<i>Lamium amplexicaule</i>
Knotweed, prostrate	<i>Polygonum aviculare</i>
Mallow, little	<i>Malva parviflora</i>
Parsley - pier	<i>Alchemilla arvensis</i>
Pennywort, lawn	<i>Hydrocotyle sibthorpioides</i>
Pineappleweed	<i>Matricaria matricarioides</i>
Speedwell, thymeleaf	<i>Veronica serpyllifolia</i>
Spurge, prostrate	<i>Euphorbia humistrata</i>
Spurge, spotted	<i>Euphorbia maculata</i>
Spurry, corn	<i>Spergula arvensis</i>
Woodsorrel, creeping	<i>Oxalis corniculata</i>
Woodsorrel, yellow	<i>Oxalis stricta</i>

Weeds suppressed or partially controlled by a single application; or, controlled by repeat applications of GALLERY plus TEAM Turf Fertilizer applied prior to weed germination at 250 to 375 pounds per acre (5.75 to 8.6 lbs/1000 sq ft):

Common Name	Scientific Name
Clover spp. ¹	<i>Trifolium spp.</i>
Clover, white ¹	<i>Trifolium repens</i>
Dandelion ¹	<i>Taraxacum officinale</i>
Groundsel, common	<i>Senecio vulgaris</i>
Medic, black	<i>Medicago lupulina</i>
Plantain, broadleaf	<i>Plantago major</i>
Plantain, buckhorn	<i>Plantago lanceolata</i>
Plantain, slender	<i>Plantago elongata</i>

¹From seed only

WARRANTY DISCLAIMER

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. DowElanco makes no other express or implied warranty of merchantability or fitness for a particular purpose or any other express or implied warranty.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to the label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of DowElanco or the seller. All such risks shall be assumed by the buyer.

BEST AVAILABLE COPY

2.46
1

LIMITATION OF REMEDIES

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at DowElanco's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

DowElanco shall not be liable for losses or damages resulting from handling or use of this product unless DowElanco is promptly notified of such loss or damage in writing. In no case shall DowElanco be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of DowElanco or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Copyright© - DowElanco - 1991