

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

SEP 2 5 2003

Ms. Kimberly S. Gilbert Regulatory Manager Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268

Dear Ms. Gilbert:

Subject: TURF FERTILIZER -- Contains Gallery* Plus Team*
Registration No. 62719-192
Application and Letter Dated July 28, 2003, Request To
Amend Registration with Revised Labeling as Described
on the Application and in the Letter; and Resubmission
of Labeling in Response To Agency's Letter Dated
August 7, 2003 (Split Labels for Residential and

The resubmitted labeling associated with your request to amend the subject pesticide product registration has been reviewed and found acceptable under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, provided that you:

1. Revise the "Non-Residential" label as follows:

Non-Residential Markets)

- a. Delete the claim for "Commercial" turf, or clearly describe the use sites. "Non-residential" turf may be used if there are no use sites that should be excluded.
- b. Delete the descriptor "General" in the "General Use Precautions and Restrictions" section.
- c. Under the "Broadcast Application Rates" section describe when to use each dose within the range of doses given. This comment applies any range of doses given on this labeling, i.e., under the "Weeds Controlled" section.
- d. Delete the statement that reads: "This productlimestone on page 2. If you wish to use this statement, you must revise the Confidential Statement of Formula (CSF) of this product by qualifying and quantifying the ingredients named in that statement.

2. Submit one copy of each of the final printed labels prior to shipment under the enclosed stamped revised labels (one residential and one non-residential).

If these conditions are not complied with, this registrations will be subject to cancellation in accordance with FIFRA, section 6(e). Your release for shipment of this product under this labeling constitutes acceptance of these conditions.

A copy of each of the two labels are enclosed for your records.

Sincerely yours,

Joanne I. Miller Product Manager (23)

Herbicide Branch

Registration Division (7505C)

Enclosure (2 Stamped Labels)

Page 1

[Editors note: Supplemental distributor label may include fertility and customer service information sections within this label.]

(Logo) Dow AgroSciences LLC

Dow AgroSciences Turf Fertilizer - Contains Gallery* plus Team*

A turf fertilizer containing Gallery and Team, preemergence herbicides for control of annual grasses and broadleaf weeds in established home lawns.

[Editor's note: One or more of the following statements may be included on the label.]

Lawn weed preventer and fertilizer

and/or

Stops weeds before they start

and/or

Prevents broadleaf weeds from germinating all year

and/or

Controls broad spectrum of weeds

and/or

Prevents dandellons, crabgrass, clover, chickweed, plantain, spurge, and henbit

One easy application

and/or

Greens and thickens lawns

Active Ingredients:

isoxaben: N-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-	
2,6-dimethoxybenzamide and isomers	0.29%
trifluralin: α,α,α-trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl-p-toluidine	
benefin: N-butyl-N-ethyl-a,a,a-trifluoro-2,6-dinitro-p-toluidine	0.53%
Inert Ingredients	
Total	

Guaranteed Fertilizer Analysis: XX-XX-XX[†]

Water-Insoluble Nitrogen	XX%
Total Nitrogen (N)	
Ammoniacal Nitrogen	
Urea Nitrogen	
Available Phosphoric Acid (P2O5)	
Soluble Potash (K ₂ O)	
Calcium	
Magnesium (Mg)	
Copper (Cu)	
Iron (Fe)	XX%
Manganese (Mn)	XX%
Zinc (Zn)	XX%

ACCEPTED
with COMMENTS
In EPA Letter Dated

SEP 2.5 2003
Under the Federal Insecticide, Fundicide, and Rodenticide Act as amended, for the posticide registered under EPA Res. No. 6.21(9-192)

[†]This product may be formulated with one or more of the following fertilizer materials: urea, methylene urea, sulfur coated urea, ammoniated phosphates, sulfate of potash, muriate of potash, copper oxide, iron oxide, manganese oxide, zinc oxide or limestone.

Contains 0.2725^{††} lb active ingredient per 25 lb bag.

[**Editor's note: Amount of active ingredient may be changed according to package size chosen by supplemental distributor.]

Keep Out of Reach of Children CAUTION

See back or side panel of container for additional precautionary statements.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-192

EPA Est. _____

*Trademark of Dow AgroSciences LLC

Dow AgroSciences LLC Indianapolis, Indiana 46268 U.S.A.

Herbicide

Net Weight - XX lb/XX.X kg

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if Swallowed or inhaled Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid breathing dust. Protective clothing such as a long-sleeved shirt, long pants, chemical-resistant gloves made of any waterproof material such as polyvinyl chloride or polyethylene, and socks and shoes must be worn when applyting this product. Users should remove protective clothing immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Application Restrictions: Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Entry Restrictions: Do not allow people or pets to enter the treated areas until dusts have settled. If watering-in is required after the application, do not enter or allow others to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your solid waste agency or 1-800-992-5994 for disposal instructions. Never place unused product down any indoor or outdoor drain.

How This Product Works

This turf fertilizer product contains Gallery* and Team* preemergence herbicides for control of certain broadleaf weeds and annual grasses in established residential turf. This product prevents certain annual grasses and broadleaf weeds in lawns. This product also feeds the lawn with a fertilizer designed to provide prolonged, even greening without excess top growth.

This product will not control existing weeds. Existing weeds should be controlled with a foliar applied post-emergence herbicide designed to be used on lawns, applied either before or after application of this product.

When to Apply

This product can be applied at any time of the spring or fall. The type of weeds controlled will vary depending on time of application so other weed control products may be needed to provide a total annual weed control program.

Early spring (before forsythia blooms): If applied in early spring, this product will prevent crabgrass and other annual grasses (including Poa annua) and broadleaf weeds, such as dandelion, clover, plantain, and spurge, that normally germinate in the spring and summer. When used in this manner, this product may be substituted for a multi-step weed and feed program. For your area, consult the local extension service for optimum timing for control of crabgrass.

Mid-late spring (after foreythia blooms, before Memorial Day): This product will prevent annual grasses (including Poa annua) and broadleaf weeds, such as dandelion, clover, plantain, and spurge, that have yet to germinate in the spring and summer, ensuring season-long weed control.

Early fall (before average night temperatures fall below 60°F): This product will prevent establishment of winter annual grasses (including Poa annua) and broadleaf weeds, such as chickweed, henbit, and speedwell, that normally germinate in the fall and winter and broadleaf weeds that germinate the following spring season. A supplemental application of a crabgrass pre-emergence herbicide may be needed in the spring for spring and summer weed control..

Late fall: This product will prevent establishment of broadleaf weeds, such as chickweed, henbit, and speedwell, that may germinate over winter and broadleaf weeds that germinate the following spring season. A supplemental application of a crabgrass pre-emergence herbicide may be needed in the spring for spring and summer weed control.

To ensure optimal fertilizer and weed control, time the application to take advantage of predicted rainfall. Otherwise, sprinkler irrigate with 0.5 inch or more to activate this product.

General Use Precautions and Restrictions:

- Apply this product only to established turf. Apply to newly seeded turf (including overseeded turf) only after seedlings are established and well rooted (generally after 3 mowings)
- Reseeding: Do not reseed treated areas for a minimum of 8 weeks following application of this product.
- · Do not apply to non-turf areas such as flower beds or gardens. Take precautions to prevent inadvertent application from rotary-type applicators to such areas. The herbicides in this product prevent seed emergence and may injure bedding plants.
- This product may injure turf that is weakened due to severe winter weather or is not well established.
- · Do not apply this product in the spring and summer to tall fescue turfgrass seeded the previous fall.
- Do not apply in the spring to Bermudagrass that has been overseeded with winter annual grasses since thinning may occur.
- Do not apply to newly sprigged or hydromulched areas of Bermudagrass, St. Augustinegrass, centipedegrass or zoysiagrass until turf is well established.
- Do not apply to dichondra.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.



- Do not overdose or overlap during application in such a way as to apply in excess of labeled rates as turf injury may occur.
- Apply this product only to the grass species listed on this label. The user assumes responsibility for any plant damage or other liability resulting from use of Gallery on turf species not recommended on this label.

This Product is Recommended for Use on the Following Established Turfgrass Species:

Established Cool Season Grasses

Common Name

Bentgrass, colonial Bentgrass, creeping Bluegrass, Kentucky Fescue, chewings Fescue, creeping red Fescue, sheep Fescue, tall Ryegrass, perennial

Established Warm Season Grasses

Common Name

Bahiagrass
Bermudagrass
Centipedegrass
Fescue, tall
St. Augustinegrass
Zoysiagrass
Zoysiagrass

How to Apply

This product may be applied to either dry or wet turf. Incorporate fertilizer and herbicide into soil by watering in. A single rainfall or sprinkler irrigation of 0.5 inch or more is required to activate this product.

Application Equipment and Application Techniques

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate. Apply this product uniformly over the lawn. A more uniform application may be achieved by spreading half of the required amount over the area and then applying the remaining half at right angles to the previous direction. Check equipment frequently to ensure it is functioning properly and applying the product uniformly.

Broadcast Application Rates

Apply this product to cool season grasses at a rate of 5.75 lb per 1000 sq ft and to warm season grasses at a rate of 7.2 lb per 1000 sq ft.

Weeds Controlled

This product prevents the following weeds when applied prior to their germination at the specified rate:

Common Name

Annual bluegrass

Barnyardgrass Burweed, lawn Buttercup Carpetweed Celery, wild Chickweed, common Chickweed, mouseear Crabgrass (Large crabgrass) (Smooth crabgrass) Field sandbur (field sandspur) Foolsparsley **Foxtails** (Bottlegrass) (Bristlegrass) (Giant foxtail) (Green foxtail) (Pigeongrass) (Robust foxtail) Geranium, Carolina Goosegrass Henbit Knotweed, prostrate Mallow, little Parsley - piert Pennywort, lawn Pineappleweed Speedwell, thymeleaf Spurge, prostrate Spurge, spotted Spurry, corn Woodsorrel, creeping Woodsorrel, yellow

Weeds suppressed or partially controlled by a single application or controlled by repeat applications 8 weeks apart of this product applied prior to weed germination:

Common Name

Clover spp. ¹
Clover, white ¹
Dandelion ¹
Groundsel, common
Medic, black
Plantain, broadleaf
Plantain, buckhorn
Plantain, slender

¹From seed only

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

- 1. Refund of purchase price paid by buyer or user for product bought, or
- 2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

*Trademark of	Dow	AgroSciences	LLC
EPA-accepted:	: /	Ī	

[Editors note: Supplemental distributor label may include fertility and customer service information sections within this label.]

(Logo) Dow AgroSciences LLC

Turf Fertilizer - Contains Gallery* plus Team*

A turf fertilizer containing Gallery and Team, preemergence herbicides for control of annual grasses and broadleaf weeds in established perennial turfgrasses such as golf courses, parks, cemeteries, and commercial and residential turf.

This product is for use by professional applicators only.

Active	ıngrea	ents:

isoxaben: N-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-	
2,6-dimethoxybenzamide and isomers	0.29%
trifluralin: α,α,α-trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl-p-toluidine	
benefin: N-butyl-N-ethyl-a,a,a-trifluoro-2,6-dinitro-p-toluidine	
Inert Ingredients	
Total	

Water-Insoluble Nitrogen	XX%
Total Nitrogen (N)	
Ammoniacal Nitrogen	
Urea Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	
Soluble Potash (K ₂ O)	XX%
Calcium	XX%
Magnesium (Mg)	XX%
Copper (Cu)	XX%
Iron (Fe)	XX%
Manganese (Mn)	XX%
Zinc (Zn)	XX%

ACCEPTED with COMMENTS In EPA Letter Dated

SEP 2.5 2003

Under the Federal Insecticide, Fundicide, and Rodenticide Act as amended, for the pesticide registered ander EPA

^TThis product may be formulated with one or more of the following fertilizer materials: urea, methylene urea, sulfur coated urea, ammoniated phosphates, sulfate of potash, muriate of potash, copper oxide, iron oxide, manganese oxide, zinc oxide or limestone.

Contains 0.2725^{††} lb active ingredient per 25 lb bag.

[ftEditor's note: Amount of active ingredient may be changed according to package size chosen by supplemental distributor.]

Keep Out of Reach of Children

CAUTION

See back or side panel of container for additional precautionary statements.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.

Page 2

11/16

In case of emergency endangering health or the environment involving this pro	oduct, call 1-800-992-5 <mark>99</mark> 4
Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.	
EPA Reg. No. 62719-192	EPA Est.

*Trademark of Dow AgroSciences LLC Dow AgroSciences LLC Indiana 46268 U.S.A.

Herbicide

Net Weight - XX lb/XX.X kg

Precautionary Statements

Hazards to Humans and Domestic Animais

CAUTION

Harmful if Swallowed or Inhaled Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions in Some Individuals

Avoid breathing dust. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Personal Protective Equipment (PPE):

Applicators and other handlers must wear:

Long-sleeved shirt and long pants

Chemical resistant gloves (≥ 14 mils) such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethelene, polyvinyl chloride, or viton Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner which will directly exposed canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Application Restrictions: Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Entry Restrictions: Do not enter or allow others to enter treated area until dusts have settled. If soil incorporation is required after application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

General Information

This turf fertilizer product contains Gallery* and Team* preemergence herbicides for control of certain broadleaf weeds and annual grasses in established turfgrasses such as golf courses, parks, cemeteries, and commercial and residential turf. Apply in late summer or early fall, approximately 3 weeks prior to germination of winter annual weeds, or in early spring, 3 weeks prior to germination of weeds which emerge in spring or early summer.

This product will not control existing weeds. Existing weeds should be controlled with a foliar applied post-emergence herbicide, applied either before or after application of this product.

To ensure optimal fertilizer and herbicidal response, water treated areas as soon as possible after application. A single rainfall or sprinkler irrigation of 0.5 inch or more is required to activate this product.

General Use Precautions and Restrictions:

- Apply this product only to established turf. Apply to newly seeded turf (including overseeded turf) only
 after seedlings are established and well rooted (generally after 3 mowings)
- Reseeding: Do not reseed treated areas for a minimum of 8 weeks after a single application of 275 lb per acre, or 12 to 16 weeks after a single application of 375 lb per acre.
- Do not apply to non-turf areas such as flower beds or gardens. Take precautions to prevent inadvertent
 application from rotary-type applicators to such areas. The herbicides in this product prevent seed
 emergence and may injure bedding plants.
- This product may injure turf that is weakened due to severe winter weather or is not well established.
- Do not apply this product in the spring and summer to tall fescue turgrass seeded the previous fall.
- Do not apply in the spring to Bermudagrass that has been overseeded with winter annual grasses since thinning may occur.
- Do not apply to newly sprigged or hydromulched areas of Bermudagrass, St. Augustinegrass, centipedegrass or zovsiagrass until turf is well established.
- Do not apply to dichondra.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.
- Do not overdose or overlap during application in such a way as to apply in excess of labeled rates as turf injury may occur.



 Apply this product only to the grass species listed on this label. The user assumes responsibility for any plant damage or other liability resulting from use of Gallery on turf species not recommended on this label.

This Product is Recommended for Use on the Following Established Turfgrass Species:

Established Cool Season Grasses

Common NameScientific NameBentgrass, colonialAgrostis tenuisBentgrass, creepingAgrostis stolonifera

Bluegrass, Kentucky Poa pratensis

Fescue, chewings Festuca rubra var. commutata

Fescue, creeping red Festuca rubra Fescue, sheep Festuca ovina

Fescue, tall Festuca arundinaceae

Ryegrass, perennial Lolium perenne

Established Warm Season Grasses

Common Name Scientific Name

Bahiagrass Paspalum notatum var. Saurae parodi

Bermudagrass Cynodon dactylon
Centipedegrass Eremochloa ophiuroides
Fescue, tall Festuca arundinaceae
St. Augustinegrass Stenotaphrum secundatum

Zoysiagrass Zoysia japonica Zoysiagrass Zoysia tenuifolia

Application Directions

Application Equipment and Application Techniques

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate. Apply this product uniformly over the lawn. A more uniform application may be achieved by spreading half of the required amount over the area and then applying the remaining half at right angles to the previous direction. Check equipment frequently to ensure it is functioning properly and applying the product uniformly.

Broadcast Application Rates

Apply this product at a rate of 250 to 375 lb per acre. A minimum of 250 lb per acre is recommended for optimum results.

	Cool Season Turfgrasses	Warm Season Turfgrasses
Summer Annual Grass Weeds	250 lb/acre	250 to 375 lb/acre
(Spring Application)	5.75 lb/1000 sq ft	5.75 to 8.6 lb/1000 sq ft
Winter Annual Grass Weeds	275 lb/acre	250 to 375 lb/acre
(Fall Application)	6.3 lb/1000 sq ft	5.75 to 8.6 lb/1000 sq ft

Weeds Controlled

This product prevents the following weeds when applied prior to weed germination at 250 to 375 lb per acre (5.75 to 8.6 lb/1000 sq ft):

Common Name

Scientific Name

Annual bluegrass

Poa annua

Barnyardgrass Burweed, lawn Buttercup

Carpetweed

Echinochloa crus-galli Soliva pterosperma Ranunculus spp. Mollugo verticillata Apium leptophyllum

Celery, wild Chickweed, common Chickweed, mouseear

Stellaria media Cerastium vulgatum

Crabgrass

Digitaria spp.

(Large crabgrass)
(Smooth crabgrass)

Field sandbur (field sandspur)
Foolsparsley

Cenchrus incertus Aethusa cynapium Setaria spp.

Foxtails
(Bottlegrass)
(Bristlegrass)
(Giant foxtail)
(Green foxtail)
(Pigeongrass)

(Robust foxtail)
Geranium, Carolina

Geranium carolinianum

Goosegrass

Eleusine indica

Henbit Knotweed, prostrate Mallow, little Parsley - piert Pennywort, lawn Lamium amplexicaule Polygonum aviculare Malva parviflora Alchemilla arvensis

Pineappleweed
Speedwell, thymeleaf
Spurge, prostrate
Spurge, spotted
Spurry, corn
Woodsorrel, creeping

Woodsorrel, yellow

Hydrocotyle sibthorpioides Matricaria matricarioides Veronica serpyllifolia Euphorbia humistrata Euphorbia maculata Spergula arvensis Oxalis corniculata Oxalis stricta

Weeds suppressed or partially controlled by a single application or controlled by repeat applications 8 weeks apart of this product applied prior to weed germination at 250 to 375 lb per acre (5.75 to 8.6 lb/1000 sq ft):

Common Name

Scientific Name

Clover spp. 1 Clover, white 1 Dandelion 1

Trifolium spp.
Trifolium repens
Taraxacum officinale
Senecio vulgaris

Groundsel, common Medic, black Plantain, broadleaf Plantain, buckhorn

Medicago lupulina Plantago major Plantago lanceolata

Plantain, bucknor Plantain, slender Plantago lanceolata
Plantago elongata

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid.

¹From seed only

Page 7

Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

- 1. Refund of purchase price paid by buyer or user for product bought, or
- 2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

*Trademark of			LLC
EPA-accepted:	/_	_/	