

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

MAY 21 2008

Ms. Kimberly Gilbert  
Regulatory Leader  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, IN 46268

RE: Notification of Minor Label Change per PR Notice 98-10  
EPA Registration Number: 62719-192  
Date of Submission: April 9, 2008

Dear Ms. Gilbert:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated, April 9, 2008, for the product, Turf Fertilizer – Contains Gallery Plus Team. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped “Notification” and will be placed in our records.

If you have any questions, please me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda Arrington".

Linda Arrington  
Notifications & Minor Formulations Team Leader  
Registration Division (7505P)  
Office of Pesticide Programs



United States  
**Environmental Protection Agency**  
 Washington, DC 20460

Registration  
 Amendment  
 Other

OPP Identifier Number

**Application for Pesticide - Section I**

1. Company/Product Number Dow AgroSciences LLC/62719-192	2. EPA Product Manager Joanne I. Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Dow AgroSciences LLC/Turf Fert-Contains Gallerv plus Team	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

**Section - II**

Amendment - Explain below.  Final printed labels in response to Agency letter dated \_\_\_\_\_

Resubmission in response to Agency letter dated \_\_\_\_\_  "Me Too" Application.

Notification - Explain below.  Other - Explain below.

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

**Section - III**

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container

3. Location of Net Contents Information  
 Label  Container

4. Size(s) Retail Container

5. Location of Label Directions

6. Manner in Which Label is Affixed to Product  
 Lithograph Paper glued Stenciled  Other \_\_\_\_\_

**Section - IV**

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Kimberly S. Gilbert	Title Regulatory Manager	Telephone No. (Include Area Code) (317) 337-4685 (fax: 317-337-4649)
-----------------------------	-----------------------------	-------------------------------------------------------------------------

**Certification**  
 I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

2. Signature <i>Kimberly S. Gilbert</i>	3. Title Regulatory Manager	6. Date Application Received (Stamped)
4. Typed Name Kimberly S. Gilbert (ksgilbert@dow.com)	5. Date April 9, 2008	

# Turf Fertilizer - Contains Gallery<sup>®</sup> plus Team<sup>®</sup>

EPA Reg. No. 62719-192

The master label for this product consists of separate, self-contained sub labels for professional use and residential use, respectively.


## Registration Notes:

Source label text based on EPA accepted copy dated September 10, 2007 and notification coded "Q8B / Turf Fertilizer – Contains Gallery Plus Team / MSTR Notif / 10-05-07."

### Changes by notification:

1. Updated trademark symbol throughout.
2. Added to list of optional label claims.

®Trademark of Dow AgroSciences LLC



**[Sub Label A: Professional Uses]**

(Base label for rigid containers 5 gal or less):

# Turf Fertilizer - Contains Gallery® plus Team®

## Herbicide

A fertilizer containing Gallery and Team, preemergence herbicides for control of annual grasses and broadleaf weeds in established turfgrass such as golf courses, parks, cemeteries, and non-residential and residential turf.

This product is for use by professional applicators only.

and/or

Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristlegrass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [field sandbur] [field sandspur] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain] ✓

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [chickweed] [henbit] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control/[prevention] than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass [and other weeds] from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types [and/or] including bentgrass

and/or

For use on cool season and warm season turfgrass

and/or

Just one application needed

and/or


More than a crabgrass preventer [also a barnyardgrass preventer]

and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as]

[crabgrass] [goosegrass] [barnyardgrass]

and/or



University tested [and researched]

and/or

Never see crabgrass again

and/or

Labeled for over 40 different [grassy] [and] [broadleaf] weeds

Active Ingredients:

isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers .....	0.29%
trifluralin: $\alpha, \alpha, \alpha$ -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine .....	0.27%
benefin: <i>N</i> -butyl- <i>N</i> -ethyl- $\alpha, \alpha, \alpha$ -trifluoro-2,6-dinitro- <i>p</i> -toluidine .....	0.53%
Other Ingredients .....	98.91%
Total .....	100.00%

Contains 0.2725 lb active ingredient per 25 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

**Guaranteed Fertilizer Analysis: XX-XX-XX**

Water-Insoluble Nitrogen .....	XX%
Total Nitrogen (N) .....	XX%
Ammoniacal Nitrogen .....	XX%
Urea Nitrogen .....	XX%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	XX%
Soluble Potash (K <sub>2</sub> O) .....	XX%
Calcium .....	XX%
Magnesium (Mg) .....	XX%
Copper (Cu) .....	XX%
Iron (Fe) .....	XX%
Manganese (Mn) .....	XX%
Zinc (Zn) .....	XX%

Keep Out of Reach of Children

**CAUTION**

**Precautionary Statements**

**Hazards to Humans and Domestic Animals**

Harmful If Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid breathing dust.

**Personal Protective Equipment (PPE)**

All loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Socks plus shoes

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**

6/37

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

**First Aid**

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control centre or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

**Environmental Hazards**

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

**Storage and Disposal**

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refer to label booklet for Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.



(Base label for nonrigid containers any size):

# Turf Fertilizer - Contains Gallery<sup>®</sup> plus Team<sup>®</sup>

## Herbicide

A fertilizer containing Gallery and Team, preemergence herbicides for control of annual grasses and broadleaf weeds in established turfgrass such as golf courses, parks, cemeteries, and non-residential and residential turf.

This product is for use by professional applicators only.

and/or

Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristleglass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [field sandbur] [field sandspur] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]

and/or

Season long crabgrass control

and/or

Season long crabgrass prevention

and/or

Easy to use

and/or

Apply [early spring] [spring] [late spring]

and/or

Apply in the fall for [annual bluegrass] [chickweed] [henbit] [and other winter weeds]

and/or

Use for a [beautiful] lawn

and/or

Better crabgrass control[/prevention] than other leading brands

and/or

Effective crabgrass control/prevention leads to [thicker] [healthier] lawns

and/or

Prevents crabgrass [and other weeds] from germinating

and/or

[Prevents] [Controls] crabgrass

and/or

For use on all turfgrass types [and/or] including bentgrass

and/or

For use on cool season and warm season turfgrass

and/or

Just one application needed

and/or

More than a crabgrass preventer [also a barnyardgrass preventer]


and/or

Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]

and/or

University tested [and researched]

and/or





**Never see crabgrass again**

and/or

**Labeled for over 40 different [grassy] [and] [broadleaf] weeds**

**Active Ingredients:**

isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers .....	0.29%
trifluralin: $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine .....	0.27%
benefin: <i>N</i> -butyl- <i>N</i> -ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>p</i> -toluidine .....	0.53%
Other Ingredients .....	98.91%
Total .....	100.00%

Contains 0.2725 lb active ingredient per 25 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

**Guaranteed Fertilizer Analysis: XX-XX-XX**

Water-Insoluble Nitrogen .....	XX%
Total Nitrogen (N) .....	XX%
Ammoniacal Nitrogen .....	XX%
Urea Nitrogen .....	XX%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	XX%
Soluble Potash (K <sub>2</sub> O) .....	XX%
Calcium .....	XX%
Magnesium (Mg) .....	XX%
Copper (Cu) .....	XX%
Iron (Fe) .....	XX%
Manganese (Mn) .....	XX%
Zinc (Zn) .....	XX%

**Keep Out of Reach of Children**

**CAUTION**

---

**Precautionary Statements**

---

**Hazards to Humans and Domestic Animals**

**Harmful If Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals**

**Avoid breathing dust.**

**Personal Protective Equipment (PPE)**

**All loaders, applicators, and other handlers must wear:**

- Long-sleeved shirt and long pants
- Socks plus shoes

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

10/37

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control centre or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

### Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Refer to label booklet for Directions for Use.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-192

EPA Est. \_\_\_\_\_

®Trademark of Dow AgroSciences LLC

Produced for

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, Indiana 46268

Net Weight \_\_\_\_\_

11/37

(Base label for refillable rigid containers greater than 5 gal):

# Turf Fertilizer - Contains Gallery® plus Team®

## Herbicide

**A fertilizer containing Gallery and Team, preemergence herbicides for control of annual grasses and broadleaf weeds in established turfgrass such as golf courses, parks, cemeteries, and non-residential and residential turf.**

**This product is for use by professional applicators only.**

and/or

**Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristleglass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [field sandbur] [field sandspur] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]**

and/or

**Season long crabgrass control**

and/or

**Season long crabgrass prevention**

and/or

**Easy to use**

and/or

**Apply [early spring] [spring] [late spring]**

and/or

**Apply in the fall for [annual bluegrass] [chickweed] [henbit] [and other winter weeds]**

and/or

**Use for a [beautiful] lawn**

and/or

**Better crabgrass control[/prevention] than other leading brands**

and/or

**Effective crabgrass control/prevention leads to [thicker] [healthier] lawns**

and/or

**Prevents crabgrass [and other weeds] from germinating**

and/or

**[Prevents] [Controls] crabgrass**

and/or

**For use on all turfgrass types [and/or] including bentgrass**

and/or

**For use on cool season and warm season turfgrass**

and/or

**Just one application needed**

and/or

**More than a crabgrass preventer [also a barnyardgrass preventer]**

and/or

**Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]**

and/or

**University tested [and researched]**

and/or

12/37

**Never see crabgrass again**  
and/or

**Labeled for over 40 different [grassy] [and] [broadleaf] weeds**

**Active Ingredients:**

isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers .....	0.29%
trifluralin: $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine .....	0.27%
benefin: <i>N</i> -butyl- <i>N</i> -ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>p</i> -toluidine .....	0.53%
Other Ingredients .....	98.91%
Total .....	100.00%

Contains 0.2725 lb active ingredient per 25 lb bag.

**[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]**

**Guaranteed Fertilizer Analysis: XX-XX-XX**

Water-Insoluble Nitrogen.....	XX%
Total Nitrogen (N).....	XX%
Ammoniacal Nitrogen .....	XX%
Urea Nitrogen .....	XX%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ).....	XX%
Soluble Potash (K <sub>2</sub> O).....	XX%
Calcium.....	XX%
Magnesium (Mg).....	XX%
Copper (Cu).....	XX%
Iron (Fe).....	XX%
Manganese (Mn).....	XX%
Zinc (Zn).....	XX%

**Keep Out of Reach of Children**

**CAUTION**

---

**Precautionary Statements**

**Hazards to Humans and Domestic Animals**

**Harmful If Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals**

**Avoid breathing dust.**

**Personal Protective Equipment (PPE)**

**All loaders, applicators, and other handlers must wear:**

- Long-sleeved shirt and long pants
- Socks plus shoes

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

13/  
27

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control centre or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

### Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Reuse:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

### Refer to label booklet for Directions for Use.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-192

EPA Est. \_\_\_\_\_

®Trademark of Dow AgroSciences LLC

Produced for  
Dow AgroSciences LLC



(Base label for nonrefillable rigid containers larger than 5 gal):

# Turf Fertilizer - Contains Gallery® plus Team®

## Herbicide

**A fertilizer containing Gallery and Team, preemergence herbicides for control of annual grasses and broadleaf weeds in established turfgrass such as golf courses, parks, cemeteries, and non-residential and residential turf.**

**This product is for use by professional applicators only.**

and/or

**Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristlegrass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [field sandbur] [field sandspur] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]**

and/or

**Season long crabgrass control**

and/or

**Season long crabgrass prevention**

and/or

**Easy to use**

and/or

**Apply [early spring] [spring] [late spring]**

and/or

**Apply in the fall for [annual bluegrass] [chickweed] [henbit] [and other winter weeds]**

and/or

**Use for a [beautiful] lawn**

and/or

**Better crabgrass control/[prevention] than other leading brands**

and/or

**Effective crabgrass control/prevention leads to [thicker] [healthier] lawns**

and/or

**Prevents crabgrass [and other weeds] from germinating**

and/or

**[Prevents] [Controls] crabgrass**

and/or

**For use on all turfgrass types [and/or] including bentgrass**

and/or

**For use on cool season and warm season turfgrass**

and/or

**Just one application needed**

and/or

**More than a crabgrass preventer [also a barnyardgrass preventer]**

and/or

**Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]**

and/or

**University tested [and researched]**

and/or

16/37

**Never see crabgrass again**

and/or

**Labeled for over 40 different [grassy] [and] [broadleaf] weeds**

**Active Ingredients:**

isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers .....	0.29%
trifluralin: $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine .....	0.27%
benefin: <i>N</i> -butyl- <i>N</i> -ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>p</i> -toluidine .....	0.53%
Other Ingredients .....	98.91%
Total .....	100.00%

Contains 0.2725 lb active ingredient per 25 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

**Guaranteed Fertilizer Analysis: XX-XX-XX**

Water-Insoluble Nitrogen.....	XX%
Total Nitrogen (N) .....	XX%
Ammoniacal Nitrogen .....	XX%
Urea Nitrogen .....	XX%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ).....	XX%
Soluble Potash (K <sub>2</sub> O).....	XX%
Calcium.....	XX%
Magnesium (Mg).....	XX%
Copper (Cu).....	XX%
Iron (Fe).....	XX%
Manganese (Mn).....	XX%
Zinc (Zn).....	XX%

**Keep Out of Reach of Children**

**CAUTION**

---

**Precautionary Statements**

---

**Hazards to Humans and Domestic Animals**

**Harmful If Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals**

**Avoid breathing dust.**

**Personal Protective Equipment (PPE)**

**All loaders, applicators, and other handlers must wear:**

- Long-sleeved shirt and long pants
- Socks plus shoes

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.



- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

### First Aid

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control centre or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

### Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

### Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refer to label booklet for Directions for Use.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.





20/37

**Never see crabgrass again**

and/or

**Labeled for over 40 different [grassy] [and] [broadleaf] weeds**

**Active Ingredients:**

isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers .....	0.29%
trifluralin: $\alpha, \alpha, \alpha$ -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine .....	0.27%
benefin: <i>N</i> -butyl- <i>N</i> -ethyl- $\alpha, \alpha, \alpha$ -trifluoro-2,6-dinitro- <i>p</i> -toluidine .....	0.53%
Other Ingredients .....	98.91%
Total .....	100.00%

Contains 0.2725 lb active ingredient per 25 lb bag.

**[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]**

**Guaranteed Fertilizer Analysis: XX-XX-XX**

Water-Insoluble Nitrogen.....	XX%
Total Nitrogen (N).....	XX%
Ammoniacal Nitrogen .....	XX%
Urea Nitrogen .....	XX%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ).....	XX%
Soluble Potash (K <sub>2</sub> O).....	XX%
Calcium.....	XX%
Magnesium (Mg).....	XX%
Copper (Cu).....	XX%
Iron (Fe).....	XX%
Manganese (Mn).....	XX%
Zinc (Zn) .....	XX%

**Keep Out of Reach of Children**

**CAUTION**

**Refer to inside of label booklet for additional precautionary information including Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.


EPA Reg. No. 62719-192

EPA Est. \_\_\_\_\_

®Trademark of Dow AgroSciences LLC

**Produced for  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, Indiana 46268**

**Net Weight** \_\_\_\_\_



21/37

(Page 1 through end):

**Precautionary Statements****Hazards to Humans and Domestic Animals****CAUTION**

**Harmful If Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals**

**Avoid breathing dust.**

**Personal Protective Equipment (PPE)**

**All loaders, applicators, and other handlers must wear:**

- Long-sleeved shirt and long pants
- Socks plus shoes

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

**First Aid**

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control centre or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

**Environmental Hazards**

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

**Directions for Use**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.  
Read all directions carefully before applying this product.

**Application Restrictions:** Do not apply this product in a way that will contact workers or other persons (or pets), either directly or through drift. Only protected handlers may be in the area during application.

**Entry Restrictions:** Do not enter or allow other people (or pets) to enter treated area until dusts have settled. If soil incorporation is required after application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

**Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.**

### **Storage and Disposal**

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

#### **Nonrefillable rigid containers 5 gallons or less:**

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

#### **Nonrefillable nonrigid containers:**

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

#### **Refillable rigid containers larger than 5 gal:**

**Container Reuse:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

#### **Nonrefillable rigid containers larger than 5 gal:**

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container

upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**General Information**

This product is a granular fertilizer containing Gallery® and Team®, preemergence herbicides for application to established turfgrass such as golf courses, parks, cemeteries, and non-residential and residential turf for control of certain broadleaf weeds and annual grasses. This product will not control existing weeds. Existing weeds should be controlled with a foliarly applied postemergence herbicide, either before or after application of this product.

**General Use Precautions and Restrictions**

- Do not apply to dichondra.
- Do not apply to newly sprigged or hydromulched areas of Bermudagrass, St. Augustinegrass, centipedegrass or zoysiagrass until turf is well established.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.
- Do not make more than 2 applications of this product per year.
- Apply this product **only** to established turf. Apply to newly seeded turf (including overseeded turf) only after seedlings are established (three leaf stage and tillering) and well rooted (generally after 3 mowings).
- **Reseeding:** Do not reseed treated areas for at least 8 weeks after a single application of 275 lb of product per acre, or 12 to 16 weeks after a single application of 375 lb of product per acre.
- Do not apply to non-turf areas such as flower beds or gardens. Take precautions to prevent inadvertent application from rotary-type applicators to such areas. The herbicides in this product prevent seed emergence and may injure bedding plants.
- This product may injure turf that is weakened due to severe winter weather or is not well established.
- Do not apply this product in the spring and summer to tall fescue turfgrass seeded the previous fall.
- Do not apply in the spring to Bermudagrass that has been overseeded with winter annual grasses since thinning may occur.
- Do not overdose or overlap during application in such a way as to apply in excess of labeled rates as turf injury may occur.
- Apply this product only to the grass species listed on this label. **The user assumes responsibility for any plant damage or other liability resulting from use of this product on turf species not recommended on this label.**
- **Maximum Single Application Rate:** Cool season turf – 275 lb of Turf Fertilizer – Contains Gallery plus Team per acre; warm season turf – 375 lb of Turf Fertilizer – Contains Gallery plus Team per acre.
- **Maximum Yearly Rate:** Cool season turf – 550 lb of Turf Fertilizer – Contains Gallery plus Team per acre; warm season turf – 750 lb of Turf Fertilizer – Contains Gallery plus Team per acre.
- **Minimum Re-Treatment Interval:** Cool season turf – 56 to 70 days; warm season turf – 70 to 84 days.
- This product must be watered in as soon as possible after application.

**This product is recommended for use on the following established turfgrass species:**

<b>Established Cool Season Turfgrass</b>	
<b>Common Name</b>	<b>Scientific Name</b>
bentgrass, colonial	<i>Agrostis tenuis</i>
bentgrass, creeping	<i>Agrostis stolonifera</i>
bluegrass, Kentucky	<i>Poa pratensis</i>
fescue, chewings	<i>Festuca rubra var. commutata</i>
fescue, creeping red	<i>Festuca rubra</i>
fescue, sheep	<i>Festuca ovina</i>
fescue, tall	<i>Festuca arundinaceae</i>
ryegrass, perennial	<i>Lolium perenne</i>

24/37

**Established Warm Season Turfgrass**

Common Name	Scientific Name
bahiagrass	<i>Paspalum notatum var. Saurae parodi</i>
Bermudagrass	<i>Cynodon dactylon</i>
centipedegrass	<i>Eremochloa ophiuroides</i>
fescue, tall	<i>Festuca arundinaceae</i>
St. Augustinegrass	<i>Stenotaphrum secundatum</i>
zoysiagrass	<i>Zoysia japonica</i>
zoysiagrass	<i>Zoysia tenuifolia</i>

**Weeds Controlled**

**Weeds controlled when applied at 250 to 375 lb per acre (5.75 to 8.6 lb/1000 sq ft):**

Common Name	Scientific Name
annual bluegrass	<i>Poa annua</i>
barnyardgrass	<i>Echinochloa crus-galli</i>
burweed, lawn	<i>Soliva pterosperma</i>
buttercup	<i>Ranunculus spp.</i>
carpetweed	<i>Mollugo verticillata</i>
celery, wild	<i>Apium leptophyllum</i>
chickweed, common	<i>Stellaria media</i>
chickweed, mouseear	<i>Cerastium vulgatum</i>
crabgrass	<i>Digitaria spp.</i>
large crabgrass	
smooth crabgrass	
field sandbur (field sandspur)	<i>Cenchrus incertus</i>
foolsparsley	<i>Aethusa cynapium</i>
foxtails	<i>Setaria spp.</i>
bottlegrass	
bristlegrass	
giant foxtail	
green foxtail	
pigeongrass	
robust foxtail	
geranium, Carolina	<i>Geranium carolinianum</i>
goosegrass	<i>Eleusine indica</i>
henbit	<i>Lamium amplexicaule</i>
knotweed, prostrate	<i>Polygonum aviculare</i>
mallow, little	<i>Malva parviflora</i>
parsley - piert	<i>Alchemilla arvensis</i>
pennywort, lawn	<i>Hydrocotyle sibthorpioides</i>
pineappleweed	<i>Matricaria matricarioides</i>
speedwell, thymeleaf	<i>Veronica serpyllifolia</i>
spurge, prostrate	<i>Euphorbia humistrata</i>
spurge, spotted	<i>Euphorbia maculata</i>
spurry, corn	<i>Spergula arvensis</i>
woodsorrel, creeping	<i>Oxalis corniculata</i>
woodsorrel, yellow	<i>Oxalis stricta</i>

**Weeds controlled by a repeat application 8 weeks after the first application when applied at 250 to 375 lb per acre (5.75 to 8.6 lb/1000 sq ft):**

Common Name	Scientific Name
-------------	-----------------





clover spp. <sup>1</sup>	<i>Trifolium</i> spp.
clover, white <sup>1</sup>	<i>Trifolium repens</i>
dandelion <sup>1</sup>	<i>Taraxacum officinale</i>
groundsel, common	<i>Senecio vulgaris</i>
medic, black	<i>Medicago lupulina</i>
plantain, broadleaf	<i>Plantago major</i>
plantain, buckhorn	<i>Plantago lanceolata</i>
plantain, slender	<i>Plantago elongata</i>

<sup>1</sup>From seed only

**Application Directions**

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

**Rates and Timing of Application**

**Broadcast Application Rates**

	Cool Season Turfgrass	Warm Season Turfgrass
summer annual grass weeds (spring application)	250 lb/acre (5.75 lb/1000 sq ft)	250 to 375 lb/acre (5.75 to 8.6 lb/1000 sq ft)
winter annual grass weeds (fall application)	275 lb/acre (6.3 lb/1000 sq ft)	250 to 375 lb/acre (5.75 to 8.6 lb/1000 sq ft)

Apply this product at a rate of 250 to 375 lb per acre. A minimum of 250 lb per acre is recommended for optimum results. Apply in late summer or early fall, approximately 3 weeks prior to germination of winter annual weeds, or in early spring, 3 weeks prior to germination of weeds which emerge in spring or early summer. Use the higher rate in the rate range where heavy weed infestation is anticipated.

**Terms and Conditions of Use**

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

**Warranty Disclaimer**

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT PERMITTED BY LAW, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

**Inherent Risks of Use**

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes),



27/37

**[Sub Label B: Residential Uses]**

(Base label):

# **Dow AgroSciences Turf Fertilizer - Contains Gallery<sup>®</sup> plus Team<sup>®</sup>**

## **Herbicide**

**A fertilizer containing Gallery and Team, preemergence herbicides for control of annual grasses and broadleaf weeds in established home lawns.**

and/or

**Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristleglass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [field sandbur] [field sandspur] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]**

and/or

**Season long crabgrass control**

and/or

**Season long crabgrass prevention**

and/or

**Easy to use**

and/or

**Apply [early spring] [spring] [late spring]**

and/or

**Apply in the fall for [annual bluegrass] [chickweed] [henbit] [and other winter weeds]**

and/or

**Use for a [beautiful] lawn**

and/or

**Better crabgrass control[/prevention] than other leading brands**

and/or

**Effective crabgrass control/prevention leads to [thicker] [healthier] lawns**

and/or

**Prevents crabgrass [and other weeds] from germinating**

and/or

**[Prevents] [Controls] crabgrass**

and/or

**For use on all turfgrass types [and/or] including bentgrass**

and/or

**For use on cool season and warm season turfgrass**

and/or

**Just one application needed**

and/or


**More than a crabgrass preventer [also a barnyardgrass preventer]**

and/or

**Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]**

and/or

**University tested [and researched]**



28/37

and/or

**Never see crabgrass again**

and/or

**Labeled for over 40 different [grassy] [and] [broadleaf] weeds**

**Active Ingredients:**

isoxaben: N-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]- 2,6-dimethoxybenzamide and isomers .....	0.29%
trifluralin: $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine .....	0.27%
benefin: N-butyl-N-ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-p-toluidine .....	0.53%
Other Ingredients .....	98.91%
Total .....	100.00%

Contains 0.2725 lb active ingredient per 25 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

**Guaranteed Fertilizer Analysis: XX-XX-XX**

Water-Insoluble Nitrogen .....	XX%
Total Nitrogen (N) .....	XX%
Ammoniacal Nitrogen .....	XX%
Urea Nitrogen .....	XX%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	XX%
Soluble Potash (K <sub>2</sub> O) .....	XX%
Calcium .....	XX%
Magnesium (Mg) .....	XX%
Copper (Cu) .....	XX%
Iron (Fe) .....	XX%
Manganese (Mn) .....	XX%
Zinc (Zn) .....	XX%

**Keep Out of Reach of Children**

**CAUTION**

**Precautionary Statements**

**Hazards to Humans and Domestic Animals**

**Harmful If Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals**

**Avoid breathing dust.**

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**First Aid**

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.



30/37

(Label booklet cover):

# Dow AgroSciences Turf Fertilizer - Contains Gallery® plus Team®

## Herbicide

**A fertilizer containing Gallery and Team, preemergence herbicides for control of annual grasses and broadleaf weeds in established home lawns.**

and/or

**Controls [tough] weeds [such as annual bluegrass] [barnyardgrass] [lawn burweed] [bottlegrass] [bristleglass] [buttercup] [carpetweed] [wild celery] [common, mouseear chickweed] [large, smooth crabgrass] [field sandbur] [field sandspur] [foolsparsley] [giant, green, robust foxtail] [pigeongrass] [Carolina geranium] [goosegrass] [henbit] [prostrate knotweed] [little mallow] [parsley-piert] [lawn pennywort] [pineappleweed] [thymeleaf speedwell] [prostrate, spotted spurge] [corn spurry] [creeping, yellow woodsorrel] [clover] [white clover] [dandelion] [common groundsel] [black medic] [broadleaf, buckhorn, slender plantain]**

and/or

**Season long crabgrass control**

and/or

**Season long crabgrass prevention**

and/or

**Easy to use**

and/or

**Apply [early spring] [spring] [late spring]**

and/or

**Apply in the fall for [annual bluegrass] [chickweed] [henbit] [and other winter weeds]**

and/or

**Use for a [beautiful] lawn**

and/or

**Better crabgrass control[/prevention] than other leading brands**

and/or

**Effective crabgrass control/prevention leads to [thicker] [healthier] lawns**

and/or

**Prevents crabgrass [and other weeds] from germinating**

and/or

**[Prevents] [Controls] crabgrass**

and/or

**For use on all turfgrass types [and/or] including bentgrass**

and/or

**For use on cool season and warm season turfgrass**

and/or

**Just one application needed**

and/or

**More than a crabgrass preventer [also a barnyardgrass preventer]**

and/or


**Two applications can be used 6 to 8 weeks apart for extended prevention of weeds [such as] [crabgrass] [goosegrass] [barnyardgrass]**

and/or

**University tested [and researched]**

and/or

**Never see crabgrass again**



31/37

and/or

**Labeled for over 40 different [grassy] [and] [broadleaf] weeds**

Active Ingredients:

isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers .....	0.29%
trifluralin: $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine .....	0.27%
benefin: <i>N</i> -butyl- <i>N</i> -ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>p</i> -toluidine .....	0.53%
Other Ingredients .....	98.91%
Total .....	100.00%

Contains 0.2725 lb active ingredient per 25 lb bag.

[Editor's note: Supplemental distributor will add fertilizer analysis and derivation statement for fertilizer carrier used.]

**Guaranteed Fertilizer Analysis: XX-XX-XX**

Water-Insoluble Nitrogen .....	XX%
Total Nitrogen (N) .....	XX%
Ammoniacal Nitrogen .....	XX%
Urea Nitrogen .....	XX%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) .....	XX%
Soluble Potash (K <sub>2</sub> O) .....	XX%
Calcium .....	XX%
Magnesium (Mg) .....	XX%
Copper (Cu) .....	XX%
Iron (Fe) .....	XX%
Manganese (Mn) .....	XX%
Zinc (Zn) .....	XX%

**Keep Out of Reach of Children**

**CAUTION**

Refer to inside of label booklet for additional precautionary information including Directions for Use.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.


Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

EPA Reg. No. 62719-192

®Trademark of Dow AgroSciences LLC

**Produced for  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, Indiana 46268**

EPA Est. \_\_\_\_\_



**Net Weight** \_\_\_\_\_

(Page 1 through end):

---

**Precautionary Statements**

---

**Hazards to Humans and Domestic Animals****CAUTION****Harmful If Swallowed or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals**

Avoid breathing dust.

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**First Aid****If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control centre or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

**Environmental Hazards**

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

---

**Directions for Use**

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

**Application Restrictions:** Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.**Entry Restrictions:** Do not allow people or pets to enter the treated areas until dusts have settled. If watering-in is required after the application, do not enter or allow others to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.**Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.****Storage and Disposal**

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.



**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

**How This Product Works**

This product is a granular fertilizer containing Gallery® and Team®, preemergence herbicides for application to established home lawns for control of certain broadleaf weeds and annual grasses. This product also feeds the lawn with a fertilizer designed to provide prolonged, even greening without excess top growth.

This product will not control existing weeds. Existing weeds should be controlled with a foliarly applied postemergence herbicide either before or after application of this product.

**General Use Precautions and Restrictions**

- Do not apply to dichondra.
- Do not apply to newly sprigged or hydromulched areas of Bermudagrass, St. Augustinegrass, centipedegrass or zoysiagrass until turf is well established.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.
- Do not make more than 2 applications of this product per year.
- Apply this product **only** to established turf. Apply to newly seeded turf (including overseeded turf) only after seedlings are established and well rooted (generally after 3 mowings).
- **Reseeding:** Do not reseed treated areas for at least 8 weeks following application of this product.
- Do not apply to non-turf areas such as flower beds or gardens. Take precautions to prevent inadvertent application from rotary-type applicators to such areas. The herbicides in this product prevent seed emergence and may injure bedding plants.
- This product may injure turf that is weakened due to severe winter weather or is not well established.
- Do not apply this product in the spring and summer to tall fescue turfgrass seeded the previous fall.
- Do not apply in the spring to Bermudagrass that has been overseeded with winter annual grasses since thinning may occur.
- Do not overdose or overlap during application in such a way as to apply in excess of labeled rates as turf injury may occur.
- Apply this product only to the grass species listed on this label. **The user assumes responsibility for any plant damage or other liability resulting from use of this product on turf species not recommended on this label.**
- **Maximum Single Application Rate:** 7.2 lb of Turf Fertilizer – Contains Gallery plus Team per 1000 sq ft.
- **Maximum Yearly Rate:** 14.4 lb of Turf Fertilizer – Contains Gallery plus Team per 1000 sq ft.
- **Minimum Re-Treatment Interval:** Cool season turf – 56 to 70 days; warm season turf – 70 to 84 days.

**This product is recommended for use on the following established turfgrass species:**

**Established Cool Season Turfgrass**


- bentgrass, colonial
- bentgrass, creeping
- bluegrass, Kentucky
- fescue, chewings
- fescue, creeping red
- fescue, sheep
- fescue, tall
- ryegrass, perennial

**Established Warm Season Turfgrass**

- bahiagrass
- Bermudagrass
- centipedegrass
- fescue, tall
- St. Augustinegrass
- zoysiagrass

**Weeds Controlled**

**Weeds controlled when applied prior to weed germination:**



31/37

- |                                |                      |
|--------------------------------|----------------------|
| annual bluegrass               | geranium, Carolina   |
| barnyardgrass                  | goosegrass           |
| burweed, lawn                  | henbit               |
| buttercup                      | knotweed, prostrate  |
| carpetweed                     | mallow, little       |
| celery, wild                   | parsley - piert      |
| chickweed, common              | pennywort, lawn      |
| chickweed, mouseear            | pineappleweed        |
| crabgrass                      | speedwell, thymeleaf |
| large crabgrass                | spurge, prostrate    |
| smooth crabgrass               | spurge, spotted      |
| field sandbur (field sandspur) | spurry, corn         |
| foolsparsley                   | woodsorrel, creeping |
| foxtails                       | woodsorrel, yellow   |
| bottlegrass                    |                      |
| bristlegrass                   |                      |
| giant foxtail                  |                      |
| green foxtail                  |                      |
| pigeongrass                    |                      |
| robust foxtail                 |                      |

**Weeds controlled by a repeat application 8 weeks after the first application when applied prior to weed germination:**

- |                            |                     |
|----------------------------|---------------------|
| clover spp. <sup>1</sup>   | medic, black        |
| clover, white <sup>1</sup> | plantain, broadleaf |
| dandelion <sup>1</sup>     | plantain, buckhorn  |
| groundsel, common          | plantain, slender   |

<sup>1</sup>From seed only

**How to Apply**

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution. Optimum weed control performance will be obtained when an application is followed by sprinkler irrigation or rainfall within a few hours. A single rainfall or sprinkler irrigation of 0.5 inch or more is required to activate this product.

**How Much to Apply**

Apply this product to cool season turfgrass at a rate of 5.75 lb per 1000 sq ft and to warm season turfgrass at a rate of 7.2 lb per 1000 sq ft.

**When to Apply**

This product can be applied at any time of the spring or fall. The type of weeds controlled will vary depending upon time of application so other weed control products may be needed to provide a total annual weed control program.

**Early spring (before forsythia blooms):** If applied in early spring, this product will prevent crabgrass and other annual grasses (including *Poa annua*) and broadleaf weeds, such as dandelion, clover, plantain, and spurge, that normally germinate in the spring and summer. When used in this manner, this product may be substituted for a multi-step weed and feed program. For your area, consult the local extension service for optimum timing for control of crabgrass.

**Mid-late spring (after forsythia blooms, before Memorial Day):** This product will prevent annual grasses (including *Poa annua*) and broadleaf weeds, such as dandelion, clover, plantain, and spurge, that have yet to germinate in the spring and summer, ensuring season-long weed control.

**Early fall (before average night temperatures fall below 60°F):** This product will prevent establishment of winter annual grasses (including *Poa annua*) and broadleaf weeds, such as chickweed, henbit, and speedwell, that normally germinate in the fall and winter and broadleaf weeds that germinate the following spring season. A supplemental application of a crabgrass pre-emergence herbicide may be needed in the spring for spring and summer weed control.

**Late fall:** This product will prevent establishment of broadleaf weeds, such as chickweed, henbit, and speedwell, that may germinate over winter and broadleaf weeds that germinate the following spring season. A supplemental application of a crabgrass pre-emergence herbicide may be needed in the spring for spring and summer weed control.

---

### Terms and Conditions of Use

---

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

---

### Warranty Disclaimer

---

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT PERMITTED BY LAW, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

---

### Inherent Risks of Use

---

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risks shall be assumed by Buyer.

---

### Limitation of Remedies

---

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses.



308/2E  
April 9, 2008



Document Processing Desk (**NOTIF**)  
Office of Pesticide Programs (7504P)  
U. S. Environmental Protection Agency  
One Potomac Yard  
2777 S. Crystal Drive  
Arlington, VA 22202

TURF FERTILIZER – CONTAINS GALLERY PLUS TEAM (A.I. BENEFIN, TRIFLURALIN, ISOXABEN)  
EPA REGISTRATION NUMBER: 62719-192  
NOTIFICATION OF MINOR LABEL CHANGE PER PR NOTICE 98-10

Enclosed please find labeling for the notification action of Turf Fertilizer – Contains Gallery® plus Team® herbicide. The following changes have been made by notification:

1. Updated trademark symbol throughout.
2. Added to list of optional label claims.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

**Contents of Submission**

- Transmittal document (this letter)
- Application for Pesticide, EPA Form 8570-1
- Label entitled Turf Fertilizer-Contains Gallery plus Team (Q8B/Turf Fertilizer - Contains Gallery Plus Team/MSTR Notif/04-09-08) (33 Pages plus Registration Notes) (5 Copies)

If you require further information, please contact Cindy Loy, Regulatory Specialist at (317) 337-4655 or Paula McKinnies, Registration Assistant for this product, at (317) 337-4679.

Sincerely,

Kimberly S. Gilbert  
Regulatory Leader  
(317) 337-4685  
(317) 337-4649 (FAX)

Enclosures

