

Turf Fertilizer - Contains Gallery* plus Surflan*

A balanced turf fertilizer containing Gallery and Surflan, selective preemergence herbicides for control of certain broadleaf weeds and annual grasses in established warm season perennial turfgrasses.

Active Ingredients:

isoxaben: N-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]- 2,6-dimethoxybenzamide and isomers.....	0.29%
oryzalin: 3,5-dinitro-N ⁴ ,N ⁴ -dipropylsulfanilamide.....	0.86%
Inert Ingredients	98.85%
Total	100.00%

GUARANTEED FERTILIZER ANALYSIS	
Total Nitrogen.....	18.0%
Available Phosphoric Acid (P2O5)	5.0%
Soluble Potash (K2O).....	9.0%

Contains 0.2875 pounds active ingredient per 25 pound bag.

**Keep Out of Reach of Children
CAUTION**

See back or container/side panel for additional precautionary statements.

Notice: Read the entire label. Use only according to label directions. Before buying this product, read Warranty Disclaimer and Limitation of Remedies Sections elsewhere on this label.

In case of emergency endangering health or the environment involving this product, call collect 517-636-4400.

Do Not Ship or Store with Food, Feeds, Drugs, or Clothing.


EPA Reg. No. 62719-193

EPA Est. No. _____

*Trademark of DowElanco

DowElanco • Indianapolis, Indiana U.S.A.

**Herbicide
Net Contents - 25 Pounds**



BEST AVAILABLE COPY

Precautionary Statements

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

CAUTION

Causes Eye Irritation - Harmful If Swallowed Or Inhaled

Avoid breathing dust and contact with eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

First Aid

If in eyes: Flush eyes with plenty of water. Get medical attention if irritation persists.

If swallowed: Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled: Remove individual to fresh air. Get medical attention if breathing difficulty occurs. If not breathing, give artificial respiration, preferably mouth-to-mouth, and get medical attention immediately.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or inadvertent application to areas adjacent to treated turfgrasses may result in reduced germination or emergence of nontarget plants. Do not contaminate water when disposing of equipment washwaters.

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read all directions carefully before applying this product.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

General Information

This turf fertilizer product contains Gallery[®] and Surflan[®] preemergence herbicides for control of certain broadleaf weeds and annual grasses in established warm season turfgrasses. Apply in late summer to early fall, approximately 3 weeks prior to the expected germination of winter annual weeds, or in early spring, 3 weeks prior to germination of weeds which emerge in spring or early summer.

This product will not control established weeds. Existing weeds should be controlled with postemergence herbicides. To insure optimal fertilizer and herbicidal response, water treated areas as soon as possible after application. A single rainfall or sprinkler irrigation of 0.5 inch or more is required to activate this product.

BEST AVAILABLE COPY

Important Use Precautions:

- Do not apply to non-turf areas such as flower beds or gardens. Take precautions to prevent inadvertent application from rotary-type applicators to such areas. Herbicides in this product prevent seed emergence and may injure bedding plants.
- This product may injure turf that is weakened due to severe winter weather or is not well established. Apply only to healthy established warm season turfgrasses.
- Do not apply this product in the spring and summer to tall fescue turfgrass seeded the previous fall.
- Do not apply in the spring to Bermudagrass which has been overseeded with winter annual grasses since thinning may occur.
- Do not apply to newly sprigged or hydromulched areas of Bermudagrass, St. Augustinegrass, centipede grass or zoysiagrass until turf is well established.
- Do not apply to dichondra.
- Do not apply to golf course putting greens.
- Any cultural practices that disturb the soil, such as aerification or verticutting should be done prior to application of this product.
- Do not overdose or overlap during application in such a way as to apply in excess of labeled rates as turf injury may occur.
- Users who wish to use this product on turfgrass species not recommended on this label may determine the suitability for such uses by treating a small area at a recommended rate. Prior to treating a larger area, the treatment should be observed for any sign of herbicidal injury for 30 to 60 days to determine if the treatment is safe to the target turf species. **THE USER ASSUMES RESPONSIBILITY FOR ANY PLANT DAMAGE OR OTHER LIABILITY RESULTING FROM USE OF GALLERY ON TURF SPECIES NOT RECOMMENDED ON THIS LABEL.**

This Product is Recommended for Use on the Following Established Warm-Season Perennial Turfgrass Species:

<u>Common Name</u>	<u>Scientific Name</u>
Bahiagrass	<i>Paspalum notatum var. Saurae parodi</i>
Bermudagrass	<i>Cynodon dactylon</i>
Centipede grass	<i>Eremochloa ophiuroides</i>
Fescue, tall	<i>Festuca arundinaceae</i>
St. Augustinegrass	<i>Stenotaphrum secundatum</i>
Zoysiagrass	<i>Zoysia japonica</i>
Zoysiagrass	<i>Zoysia tenuifolia</i>

Application

Application Equipment and Application Techniques

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to manufacturer's directions. Check frequently to be sure equipment is operating properly and distributing product uniformly. Avoid use of spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turf injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. A splash board mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

Broadcast Application Rate

Apply at a broadcast rate of 233 pounds per acre (5.3 pounds per 1000 square feet).

BEST AVAILABLE COPY

Weeds Controlled

Summer Annual weeds Controlled by a Single Application:

<u>Common Name</u>	<u>Scientific Name</u>
Barnyardgrass	<i>Echinochloa crus-galli</i>
Burweed, lawn	<i>Soliva pterosperma</i>
Carpetweed	<i>Mollugo verticillata</i>
Celery, wild	<i>Apium leptophyllum</i>
Crabgrass	<i>Digitaria spp.</i>
(Large crabgrass)	
(Smooth crabgrass)	
Crowfootgrass	<i>Dactyloctenium aegyptium</i>
Foxtails	<i>Setaria spp.</i>
(Bottlegrass)	
(Bristlegrass)	
(Giant foxtail)	
(Green foxtail)	
(Pigeongrass)	
(Robust foxtail)	
(Yellow foxtail)	
Goosegrass	<i>Eleusine indica</i>
Johnsongrass	<i>Sorghum halepense</i>
(seedling only)	
Knotweed, prostrate	<i>Polygonum aviculare</i>
Kochia	<i>Kochia scoparia</i>
Lambsquarters, common	<i>Chenopodium album</i>
Mallow, little	<i>Malva parviflora</i>
Pennywort, lawn	<i>Hydrocotyle sibthorpioides</i>
Pigweeds	<i>Amaranthus spp.</i>
Pineappleweed	<i>Matricaria matricarioides</i>
Purslane, common	<i>Portulaca oleracea</i>
Spurge, prostrate	<i>Euphorbia humistrata</i>
Spurge, spotted	<i>Euphorbia maculata</i>
Spurry, corn	<i>Spergula arvensis</i>
Woodsorrel, creeping	<i>Oxalis corniculata</i>
Woodsorrel, yellow	<i>Oxalis stricta</i>

Winter Annual Weeds Controlled by a Single Application:

<u>Common Name</u>	<u>Scientific Name</u>
Buttercup	<i>Ranunculus spp.</i>
Bluegrass, annual	<i>Poa annua</i>
Chickweed, common	<i>Stellaria media</i>
Chickweed, mouseear	<i>Cerastium vulgatum</i>
Foolsparsley	<i>Aethusa cynapium</i>
Geranium, Carolina	<i>Geranium carolinianum</i>
Henbit	<i>Lamium amplexicaule</i>
Parsley - plant	<i>Alchemilla arvensis</i>
Ryegrass, annual	<i>Lolium multiflorum</i>
Speedwell, thymeleaf	<i>Veronica serpyllifolia</i>

BEST AVAILABLE COPY

Weeds partially controlled by a single application; or, controlled by repeat applications:

<u>Common Name</u>	<u>Scientific Name</u>
Clover spp. ¹	<i>Trifolium spp.</i>
Clover, white ¹	<i>Trifolium repens</i>
Dandelion ¹	<i>Taraxacum officinale</i>
Groundsel, common	<i>Senecio vulgaris</i>
Medic, black	<i>Medicago lupulina</i>
Plantain, broadleaf	<i>Plantago major</i>
Plantain, buckhorn	<i>Plantago lanceolata</i>
Plantago, slender	<i>Plantago elongata</i>

¹From seed only

[Editor's note: Subregistrant may insert appropriate Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies statements.]

©Copyright 1994 by DowElanco

BEST AVAILABLE COPY