



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
62719-215

Date of Issuance:
NOV 14 2005

NOTICE OF PESTICIDE:
 Registration
 x Reregistration
(under FIFRA, as amended)

Term of Issuance:
Unconditional

Name of Pesticide Product:
Grandstand R

Name and Address of Registrant (include ZIP Code):

Robert Peterson
DOW AgroSciences
9330 Zionsville Road
Indianapolis, IN 46268

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Based on your response to the Triclopyr Reregistration Eligibility Decision (RED) document, EPA has reregistered the product listed above provided you:

1. Under the subheading Swath Adjustment, replace ...the swath will be displaced downward with **downwind**....
2. Add to the Environmental Hazards section, "This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable particularly where the water table is shallow may result in groundwater contamination.

3. Under the heading "Limitation of Remedies" add the following text:

To the extent allowed by law, all such risks shall be assumed by the buyer.

Signature of Approving Official:

Date:

11-14-05

2/13

(Base Label):

(Logo) Dow AgroSciences LLC

Grandstand* R

For selective postemergence broadleaf weed control in rice.

Active Ingredient(s):

Triclopyr (3,5,6-trichloro-2-pyridinyloxyacetic acid), triethylamine salt	44.4%
Inert Ingredients	55.6%
Total Ingredients	100.0%

Acid Equivalent: Triclopyr - 31.8% - 3 lb/gallon

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

NOV 14 2005

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.**

62719-215

Keep Out of Reach of Children

DANGER PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Corrosive • Causes Irreversible Eye Damage • Harmful If Swallowed Or Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals.

Do not get in eyes, on skin, or on clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Protective eyewear
- Chemical resistant gloves (≥ 14 mils) such as butyl rubber, natural rubber, neoprene rubber or nitrile rubber

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the WPS (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product.
- Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Note to Applicator: Allergic skin reaction is not expected from exposure to spray mixtures of this product when used as directed.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

Environmental Hazards

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark except when treating rice fields as specified in this product label. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Physical or Chemical Hazards

Do not use or store the product near heat or open flame.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to label booklet for Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-215

EPA Est. 00000-XX-00

*Trademark of Dow AgroSciences LLC

4/B

Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Herbicide

Net Contents __ gal

5/13

(Label Booklet Cover):

(Logo) Dow AgroSciences LLC

Grandstand* R

For selective postemergence broadleaf weed control in rice.

Active Ingredient(s):

Triclopyr (3,5,6-trichloro-2-pyridinyloxyacetic acid), triethylamine salt	44.4%
Inert Ingredients	55.6%
Total Ingredients	100.0%

Acid Equivalent: Triclopyr - 31.8% - 3 lb/gallon

Keep Out of Reach of Children

DANGER PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Personal Protective Equipment (PPE), User Safety Recommendations and Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-215

EPA Est. 00000-XX-00

*Trademark of Dow AgroSciences LLC
Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Herbicide

Net Contents __ gal

(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals

DANGER

Corrosive • Causes Irreversible Eye Damage • Harmful If Swallowed Or Absorbed Through Skin • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals.

Do not get in eyes, on skin, or on clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Protective eyewear
- Chemical resistant gloves (\geq 14 mils) such as butyl rubber, natural rubber, neoprene rubber or nitrile rubber

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the WPS (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product.
- Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Note to Applicator: Allergic skin reaction is not expected from exposure to spray mixtures of this product when used as directed.

7/B

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

Environmental Hazards

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark except when treating rice fields as specified in this product label. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Physical or Chemical Hazards

Do not use or store the product near heat or open flame.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Dow AgroSciences does not intend that this product be used for manufacturing or formulating.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Shoes plus socks
- Protective eyewear
- Chemical-resistant gloves (\geq 14 mils) such as butyl rubber, natural rubber, neoprene rubber or nitrile rubber

Storage and Disposal

Do not contaminate water, food or feed by storage and disposal. Open dumping is prohibited.

Pesticide Storage: Store above 28°F or agitate before use. Do not store near heat or open flame.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic containers may also be disposed of by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General: Consult federal, state, or local authorities for approved alternative procedures.

General Information

Grandstand* R is postemergence systemic herbicide for the control of certain broadleaf weeds in rice (including ratoon rice). Grandstand R controls broadleaf weeds through foliar uptake; therefore, thorough coverage of target weeds is important. DO NOT apply under conditions which would allow spray drift to come in contact with adjacent broadleaf crops as crop injury may occur.

Special Use Precautions and Restrictions

1. Apply this product to rice only as specified on this label. Do not apply to any other crop or site.
2. Do not apply this product to upland (non flooded) rice.
3. Do not apply this product prior to the 2 to 3-leaf stage or after the 1/2" internode elongation stage of rice development (see special timing of application instructions for water seeded rice). Do not apply in the booting or subsequent stages of rice development.
4. Direct application to ditches used to transport irrigation water is prohibited.
5. Do not apply more than 1 pint (0.375 lb a.e.)/acre in a single application. Do not make more than two applications or apply more than 2 pints (0.75 lb a.e.) per acre during the growing season. Applications made after planting of rice must be at least 20 days apart.
6. Do not apply Grandstand R directly to, or otherwise permit it to come into contact with, cotton, soybeans, grapes, tobacco, vegetable crops, flowers, ornamental shrubs or trees, or other desirable broadleaf plants, as serious injury may occur. DO NOT permit spray mists containing Grandstand R to drift onto desirable broadleaf plants.
7. Do not rotate treated land to crops other than rice for 4 months following treatment.
8. When using tank mixtures, read and follow the use directions and precautions on each product label.
9. Do not apply less than 20 days prior to draining the field, unless the water is contained within a tailwater recovery system, or other system appropriate for preventing discharge from rice. Discharge is permitted 20 days following the last application of Grandstand R within the system.
10. **Application to fields which have been leveled (except water leveling) within 12 months prior to application may result in serious rice injury in areas that have been cut or filled.**
11. Do not apply where runoff or irrigation water may flow directly onto agricultural land other than rice fields as injury to crops may occur.
12. Dow AgroSciences does not intend that this product be used for manufacturing or formulating.
13. **Preharvest Interval:** Do not apply later than 60 days before harvesting rice.
14. **Chemigation:** Do not apply this product through any type of irrigation system.
15. Do not fish or commercially grow fish, shellfish or crustaceans (except crawfish) on treated acres during the 12 months following treatment. For crawfish production, do not apply Grandstand R later than 3 months prior to crawfish harvest.
16. Do not apply Grandstand R with 32% liquid nitrogen fertilizer or zinc fertilizer.
17. Do not apply Grandstand R following application of Whip herbicides, except in California where Grandstand R may be applied 14 days after application of Whip.
18. Use of this product on rice grown in the state of New York is prohibited.

Mixing Instructions

Mixing Order

When preparing spray mixtures, the recommended order of addition to the spray tank is half the water, drift control agent (if used), additional herbicide (if used), and Grandstand R. Then add the remainder of the water. The nonionic surfactant or crop oil concentrate should be added last unless otherwise specified on the surfactant label. Moderate continuous agitation is also required when Grandstand R is tank mixed with emulsifiable concentrate herbicides. When using any tank mixture, read and follow the use directions and precautions on each product label.

Spray Surfactants

For best broadleaf weed control with Grandstand R alone, use of a nonionic surfactant (registered for agricultural use where required) or a crop oil concentrate (COC) is recommended. The suggested rate of

surfactant addition to the spray mixture is 0.25% to 0.5% by volume (2 to 4 pints per 100 gallons of spray mixture) unless otherwise recommended by the surfactant label. The suggested rate of the COC addition to the spray mixture is 1% by volume (8 pints per 100 gallons of spray mixture). Read and follow all use directions and precautions on the surfactant or COC label.

Application Precautions

Aerial Application: Broadcast apply Grandstand R in a minimum of 5 gallons of spray mixture per acre, except where state regulations specify a higher minimum gallonage. For post-flood applications or when foliage is dense, use a spray volume of 5-10 gallons per acre to ensure uniform coverage. Apply at a height which provides the most effective swath width for the aircraft. Fixed-wing aircraft or helicopters should have a well-designed spray system that produces a uniform spray pattern and minimizes spray drift.

Ground Application: Broadcast apply Grandstand R in a minimum of 10 gallons of spray mixture per acre. Flat fan nozzles are recommended. Utilize a well-designed spray system that produces a uniform spray pattern and minimizes spray drift.

Avoid Injurious Spray Drift

Make applications only when there is little or no hazard from spray drift. Small quantities of spray, which may not be visible, may seriously injure susceptible plants. **Do not** spray when wind is blowing toward susceptible crops or ornamental plants that are near enough to be injured. It is suggested that a continuous smoke column or similar equipment be used at the spray site to detect air movement, lapse conditions, or temperature inversions. If the smoke layers or indicates a potential for hazardous drift, do not spray.

Spray Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications:

1. The distance of the outer most operating nozzles on the boom must not exceed 3/4 the length of the rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the following **Aerial Drift Reduction Advisory**. [This information is advisory in nature and does not supersede mandatory label requirements.]

Aerial Drift Reduction Advisory

Information On Droplet Size: The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

Controlling Droplet Size:

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

- **Pressure** - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released parallel to the airstream produced larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Length: For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height: Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment: When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.)

Wind: Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature And Humidity: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions: Applications should not occur during a local, low level temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of the smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas: The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

Application Directions

Application Timing

Grandstand R may be applied as a preplant burndown treatment prior to planting rice, to newly seeded rice, or ratoon rice following harvest of the first crop.

Preplant Burndown Application: Apply at least 21 days before planting dry seeded rice and 14 days before planting water seeded rice.

Application to Newly Seeded Rice: Apply from the 2 to 3-leaf to 1/2" internode elongation stage of growth. If two applications are made during the 2 to 3-leaf to 1/2" internode elongation stage of growth, they must be at least 20 days apart. (See special timing of application instructions for water seeded rice under "Water Management").

Ratoon Rice: Grandstand R may be applied within two weeks following harvest of the first crop to provide control of susceptible broadleaf weeds in ratoon rice.

Note: Rice is most tolerant to postemergence applications of Grandstand R from the 2 to 3-leaf stage to the 1/2" internode elongation stage of rice development. Postemergence applications of the higher rates of Grandstand R may result in temporary rice injury that appears as leaf chlorosis or stunting. Rice will normally recover from these symptoms in two to four weeks. Treatments applied after the 1/2" internode elongation stage may result in increased rice injury. **Do not apply in the booting or subsequent stages.**

Repeat Applications: Do not make more than two applications during the entire crop growing season. See "Special Use Precautions and Restrictions" for details.

Water Management

Preflood Application: For preflood applications, the rice should be in the 2 to 3-leaf stage or larger. A shallow flood may be applied no sooner than 72 hours following application of Grandstand R. If the weeds are drought stressed, flush the field before applying Grandstand R so that weeds are actively growing at time of treatment.

Postflood Application: For postflood applications, treatments should be made when weeds are well emerged above the water surface. Weeds submerged at the time of application will not be controlled. If water level is dropped to expose weeds prior to application, **do not** raise water level for at least 48 hours after application. The growing points of rice plants at the soil surface (crown) should be covered with water at the time of application.

Water Seeded Rice: In water seeded rice, **do not apply before the 3 to 4 leaf stage** or after the 1/2" internode elongation stage of growth.

Tolerance of Rice Varieties: Grandstand R may be used on all rice varieties except the variety "Millie" when grown in the state of Louisiana. However, because new varieties are introduced frequently, the tolerance of a new rice variety to Grandstand R should be checked before large areas are treated.

Application Rates and Weeds Controlled With Grandstand R Alone

Grandstand R should be applied to actively growing weeds at a rate of 0.67 to 1.0 pint (0.25 to 0.375 lb a.e.) per acre with a nonionic surfactant (0.25 to 0.5% by volume) or crop oil concentrate (1% by volume) (see Spray Surfactants under Mixing Directions). **Apply 1 pint for difficult to control species, when broadleaf weeds are large, or in postflood applications.**

12/13

Weed Control Information

Weeds Controlled	Application Rate	Comments
Common Cocklebur Jointvetch spp. ¹ Morningglory spp. ²	0.67-1.0 pint/acre (0.25-0.375 lb a.e./acre)	General: Best control is achieved with applications prior to weed flowering. Weeds larger than 24" in size may not be adequately controlled. Postflood applications should be made when weeds are well emerged above the water surface. Weeds submerged at the time of application will not be controlled. ¹ Jointvetch species are most susceptible from 10" to flowering stage of growth. ² Apply 1 pint/acre when morningglory runners are greater than 6". ³ Rice flatsedge should be treated when less than 4 inches tall.
Alligatorweed Dayflower Eclipta Hemp sesbania Redstem Rice flatsedge ³ Sicklepod Texasweed/Mexicanweed Water Hysopp Ricefield bulrush	1.0 pint/acre (0.375 lb a.e./acre)	

Application Timing and Water Management for Preflood Application in Drill-Seeded Rice:

Application Rates	Drill-Seeded Rice - Preflood Application		
	Rice Stage of Growth to Apply		Water Management
	2-leaf Stage	3 to 4-leaf Stage	Hours After Application Before Flooding
Grandstand R Alone			
0.5 pt	No	No	--
0.67 pt	No	Yes	72 hours
1.0 pt	No	Yes	72 hours
Grandstand R Plus Arrosolo or Propanil			
0.5 pt	Yes	Yes	72 hours
0.67 pt	No	Yes	72 hours

Tank Mix Recommendations

Grandstand R may be tank mixed with several rice herbicides for broad spectrum weed control in rice. Tank mix applications are to be used only when the rice is well established and in the recommended stage of growth for treatment with Grandstand R and the recommended tank mix product. For best results, weed species should also be in the proper stage of growth as specified on the Grandstand R and tank mix product label. **When tank mixing, always follow the use directions and precautions in accordance with each herbicide label. No label dosage rates may be exceeded.**

1. Preflood Application

Tank Mix with Propanil Herbicides: Grandstand R may be tank mixed with propanil herbicides in a preflood application to control grass and broadleaf weed species. Apply 0.5 to 0.67 pint/acre (0.19 - 0.25 lb a.e./acre) Grandstand R plus 3.0 to 4.0 lb ai/acre of the propanil herbicide. **DO NOT** add a surfactant or crop oil concentrate when using propanil herbicides formulated as emulsifiable concentrates. A nonionic surfactant at 0.25% by volume is recommended when using propanil herbicides formulated as dry products or as flowables.

Tank Mix with Arrosolo Herbicides (Not Registered in California): Grandstand R may be tank mixed with liquid Arrosolo herbicide in a preflood application to control grass and broadleaf weed species. Apply

0.5 to 0.67 pint/acre (0.19 - 0.25 lb a.e./acre) of Grandstand R plus 4.5 to 6 lb ai/acre (3 to 4 quarts/acre) of Arrosolo herbicide. **DO NOT** add a surfactant or crop oil concentrate to this tank mixture.

2. Postflood Application

Grandstand R may be tank mixed with propanil herbicides in a postflood application to control grass and broadleaf weed species. Apply 0.67 to 1.0 pint/acre (0.25 - 0.375 lb a.e./acre) of Grandstand R plus 1.0 to 4.0 lb ai/acre of the propanil herbicide. **DO NOT** add a surfactant or crop oil concentrate when using propanil herbicides formulated as emulsifiable concentrates. A nonionic surfactant at 0.25% by volume is recommended when using propanil herbicides formulated as dry products or as flowables. **When using the 1 lb/acre rate of propanil with Grandstand, use only the liquid propanil herbicide formulation.**

Water Seeded Rice: Grandstand R may be tank mixed with liquid Arrosolo herbicide in a postflood application to water seeded rice to control grass and broadleaf weeds. Apply 0.5 to 0.67 pint/acre (0.19 - 0.25 lb a.e./acre) of Grandstand R plus 4.5 to 6 lb ai/acre (3 to 4 quarts/acre) of Arrosolo herbicide.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences **MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

✓ The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

*Trademark of Dow AgroSciences LLC
EPA-accepted: ___/___/___